



**HAL**  
open science

## Simulating the evolution of soil solutions in irrigated rice soils in the Sahel

Claude Hammecker, P. van Asten, S. Marlet, Jean-Luc Maeght, Roland Poss

► **To cite this version:**

Claude Hammecker, P. van Asten, S. Marlet, Jean-Luc Maeght, Roland Poss. Simulating the evolution of soil solutions in irrigated rice soils in the Sahel. *Geoderma*, 2009, 150 (1-2), pp.129-140. 10.1016/j.geoderma.2008.12.024 . ird-00396823

**HAL Id: ird-00396823**

**<https://ird.hal.science/ird-00396823>**

Submitted on 19 Jun 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Elsevier Editorial System(tm) for Geoderma  
Manuscript Draft

Manuscript Number: GEODER4246R1

Title: Simulating the evolution of soil solutions in irrigated rice soils in the Sahel.

Article Type: Research Paper

Section/Category: Soil Chemistry and Biogeochemistry

Keywords: alkalization; sodication; Phreeqc; plant uptake; water management; West Africa

Corresponding Author: Dr Claude Hammecker, Ph.D.

Corresponding Author's Institution: Institut de Recherche pour le Développement - LDD

First Author: Claude Hammecker, PhD

Order of Authors: Claude Hammecker, PhD; Piet van Asten, PhD; Serge Marlet, PhD; Jean-Luc Maeght, Master Sc; Roland Poss, PhD

Abstract: In sahelian countries, agriculture often relies on irrigation especially for rice cropping which has been introduced these last decades. As most of the water resources used for irrigation (mainly rivers and lakes) have an alkaline composition, rice cropping is often held as responsible for soil alkalization. In order to evaluate the probability for this process to occur, we studied 3 different sites in West Africa where numerical simulation of the geochemical interactions between soil, irrigation water and groundwater were performed with PHREEQC. It was possible to simulate the evaporation of the ponding water, the mixing between groundwater and irrigation water, the lateral drainage (flushing) and the leaching, and finally the management of the crop residues. By performing prospective simulations over 60 cropping cycles it has been shown that soil alkalization and sodication are not inevitable in these area especially if the cropping is properly managed. Moreover, when straw is removed and the plot flushed before the new cropping season, alkaline soils can even be improved with rice cropping.



## Simulating the evolution of soil solutions in irrigated rice soils in the Sahel.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Claude Hammecker<sup>a</sup>, Piet van Asten<sup>b e</sup>, Serge Marlet<sup>c</sup>, Jean-Luc Maeght<sup>a</sup> and Roland Poss<sup>d</sup>.

<sup>a</sup> Institut de Recherche pour le Développement UR 176, Land Development Department,  
Office of Science for Land Development, Pahonyothin Road, Chatuchak, Bangkok 10900, Thailand

<sup>b</sup> WARDA, The Africa Rice Center, BP 96, St.Louis, Senegal.

<sup>c</sup> CIRAD, UMR G-EAU, Avenue Agropolis, 34398 Montpellier Cedex 5, France

<sup>d</sup> Institut de Recherche pour le Développement UR 176, SupAgro, 2 place Viala, 34060 Montpellier  
cedex 1, France

<sup>e</sup> present address : IITA- ESARC P.O.Box 7878 Kampala, Uganda

## Abstract

1  
2  
3 In sahelian countries, agriculture often relies on irrigation especially for rice cropping which has  
4  
5 been introduced these last decades. As most of the water resources used for irrigation (mainly rivers  
6  
7 and lakes) have an alkaline composition, rice cropping is often held as responsible for soil  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

In sahelian countries, agriculture often relies on irrigation especially for rice cropping which has been introduced these last decades. As most of the water resources used for irrigation (mainly rivers and lakes) have an alkaline composition, rice cropping is often held as responsible for soil alkalization. In order to evaluate the probability for this process to occur, we studied 3 different sites in West Africa where numerical simulation of the geochemical interactions between soil, irrigation water and groundwater were performed with PHREEQC. It was possible to simulate the evaporation of the ponding water, the mixing between groundwater and irrigation water, the lateral drainage (flushing) and the leaching, and finally the management of the crop residues. By performing prospective simulations over 60 cropping cycles it has been shown that soil alkalization and sodication are not inevitable in these area especially if the cropping is properly managed. Moreover, when straw is removed and the plot flushed before the new cropping season, alkaline soils can even be improved with rice cropping.

Keywords : alkalization, sodication, PHREEQC, plant uptake, water management, West Africa

## Introduction

1  
2  
3 In the Sahel, irrigated rice (*Oryza Sativa* L.) cropping has been introduced on a wide scale during  
4  
5 the last 20 years. The construction of large (> 100 ha) and costly irrigation schemes were intended  
6  
7 to increase food security and provide income for Sahelian farmers, whose crops and livestock were  
8  
9 devastated during the droughts in the 1970-1980's. Irrigated rice cropping in the hot and dry  
10  
11 Sahelian climate seems like a paradox. However, the major Sahelian rivers and some large artificial  
12  
13 reservoirs provide more than sufficient water to support current irrigation activities throughout the  
14  
15 year. Although most Sahelian irrigation waters contain relatively little salt, they usually show  
16  
17 alkaline compositions, where alkalinity (Alk.) is the dominant aqueous species. It is defined as the  
18  
19 algebraic sum of the strong base cation and strong acid anion equivalents (Stumm and Morgan,  
20  
21 1970), and can be expressed as :  $Alk. = (HCO_3^-) + (CO_3^{2-}) + (OH^-) - (H^+)$  , where ( ) denote the total  
22  
23 concentration in  $mol_c l^{-1}$  . Moreover most of the Sahelian irrigation waters, show also a positive  
24  
25 calcite residual alkalinity ( $RA_{calcite}$ ) which corresponds to the alkalinity of the solution after  
26  
27 precipitation of calcite :  $RA_{calcite} = Alk. - (Ca^{2+})$ , (Valles et al., 1991; Bertrand et al., 1993).  $RA_{calcite}$   
28  
29 has also been shown to be a conservative tracer of the soil solution ( Ribolzi et al., 1996; Barbiero et  
30  
31 al., 2001; Barbiero et al., 2004). Even the residual sodium carbonate, defined as  $RSC = Alk. -$   
32  
33  $(Ca^{2+}) - (Mg^{2+})$  ; which corresponds to the alkalinity of the solution after calcite and Mg-carbonate  
34  
35 or Mg-silicate precipitation, is positive for these solutions and is an indicator of potential  
36  
37 alkalinization and sodication (Eaton, 1950; Hardie and Eugster, 1970; Van Beek and Van Breemen,  
38  
39 1973; Al Droubi et al.,1980). Continuous concentration of such waters in the soil root zone may  
40  
41 lead to the formation of alkaline and sodic soils, showing both high pH values and high  
42  
43 exchangeable sodium percentage :  $ESP = XNa^+ / CEC$  , where  $XNa^+$  is the exchangeable  $Na^+$  and  
44  
45 CEC the total cation exchange capacity. Such soils are less productive because of the pH-induced  
46  
47 loss or low availability of several plant nutrients (e.g. Fe, P, Zn) and poor physical properties of the  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

sodic horizon (Abrol et al., 1988).

1 Whether or not large quantities of alkaline salts will accumulate in the soil root zone depends  
2  
3 largely on the water balance of the root zone. Evapotranspiration in the Sahelian climate can be  
4  
5 very large and regularly exceeds rates of  $10 \text{ mm day}^{-1}$  in the dry season (November – May) when  
6  
7 maximum temperatures reach 40 to 45 °C (Van Asten et al., 2003, Hammecker et al., 2003). In most  
8  
9 irrigation schemes, percolation rates are very low ( $< 4 \text{ mm day}^{-1}$ ) (Condom, 2000 ; Hammecker et  
10  
11 al., 2003 ; Van Asten, unpublished), due to the low hydraulic conductivity of the clayey soils and air  
12  
13 entrapment (Hammecker et al., 2003). In the course of the cropping season, percolation rates can  
14  
15 further decrease if groundwater levels reach the soil surface. The combination of large  
16  
17 evapotranspiration and small percolation can lead to the excessive build up of alkaline salts in the  
18  
19 long run.  
20  
21  
22  
23  
24

25 Recent studies have shown that the current soil alkalinity and sodicity problems in Sahelian  
26  
27 irrigation schemes are primarily caused by salts that originate from the parent material (Barbiero  
28  
29 and Van Vliet-Lanoe, 1998, Van Asten et al., 2004), or that accumulated as a result of long-time  
30  
31 irrigation of non-flooded crops in combination with shallow saline groundwater levels. Hence, these  
32  
33 salts were already in the soils before irrigated rice cropping was introduced and there is little  
34  
35 indication that the recent introduction of irrigated rice cropping contributed to a significant increase  
36  
37 in the salt content of these soils (van Asten, 2003). Moreover, most rice soils have the capacity to  
38  
39 buffer an increase in soil solution alkalinity through desorption of exchangeable  $\text{Ca}^{2+}$  and the  
40  
41 subsequent precipitation of calcite. However, although this buffer capacity is large, it is not infinite.  
42  
43 Consequently, it still remains uncertain whether current irrigation practices will not lead to soil  
44  
45 alkalinization and sodication in the long run (i.e. some decades) in areas where salts added through  
46  
47 irrigation can not sufficiently be leached or evacuated from the soil root zone. Given the high  
48  
49 investment costs of irrigated agriculture and the high costs of reversing alkalinization and  
50  
51 sodication processes, it is important to know whether current or alternative crop and water  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

management will lead to the degradation of one of the Sahel's most productive crop system.

1  
2  
3 In this study, we focus on three Sahelian irrigation situations, inside irrigation schemes, that differ  
4 distinctly in terms of parent material, soil water balance and irrigation water quality. All sites have  
5 earlier been subject of studies on soil and water quality (Hammecker and Maeght, 1999; Boivin et  
6 al., 2002; Van Asten, 2003). Therefore, mean values for basic crop, water and soils parameters are  
7 already available. Taking into account some of the most important chemical and hydrological  
8 features, we tried to simulate how soils and soil solutions would evolve over time, under current  
9 and alternative land management practices at the three sites. The most critical parameter, i.e. the  
10 ratio between leached and irrigated solution volume, is the leaching fraction (LF) whether with  
11 vertical drainage or lateral drainage (flushing). In order to account for spatial variability within the  
12 irrigation schemes, we calculated soil solution evolution for different leaching fractions. This  
13 approach allows for the identification of the minimum leaching fraction that is required, in order to  
14 prevent soil degradation in the long run. Proton excretion (or alkalinity uptake) by the rice plant can  
15 also contribute significantly to the alkalinity balance of the soil root zone (Van Asten et al., 2004)  
16 and was also taken into account.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 The objectives of this study are: (i) to give an overview of basic soil and water parameters for a  
41 range of Sahelian irrigation schemes, (ii) to simulate soil evolution under current and alternative  
42 crop and water management practices, and (iii) to identify the minimum leaching fraction that is  
43 required to prevent soil degradation in the long term.  
44  
45  
46  
47  
48  
49  
50  
51

## 52 2. Material and methods

### 53 54 55 56 57 2.1. Study sites 58 59 60 61 62 63 64 65



1 This study is based on data collected in three different irrigation areas in West Africa located in the  
2 valley of River Senegal, the Gorgol noir and in the valley of river Niger, where irrigated rice  
3 cropping is the main agricultural activity.  
4

5 These sites are representative of most of the conditions where rice-cropping cultivation is observed  
6 in the Sahel, and are representative of different geological, geomorphological and pedological  
7  
8 situations with different soil textures. They are representative of (i) silty clay loam soils, developed  
9  
10 on crystalline bedrock (Foum Gleita), (ii) alluvial levees with loamy sand soils (Niono) and (iii)  
11  
12 alluvial basin with clayey vertisol (Donaye).  
13  
14  
15  
16  
17  
18  
19

#### 20 2.1.1. Foum Gleita

21 The 'Mauritanides' geological region where Foum Gleita is located, is representative of the Pan-  
22 African orogenic belt which has been folded and lifted in the Appalachian era (Carité, 1989) and is  
23 consisting of metamorphic rocks, mainly greenschists, calcshists, micaschists and quartz, and  
24  
25 secondary intrusions of dolerite, on which pedogenesis developed alkaline soils.  
26  
27  
28  
29  
30  
31

32 The Foum Gleita irrigation scheme is located in central southern Mauritania (N16°08'; W12°46').  
33

34 The scheme was constructed between 1985 and 1989, with a surface area of 1950 ha. Irrigation  
35 water originates from a large artificial lake that formed behind a dam in the Gorgol Noir river.  
36  
37

38 Irrigation and drainage are gravimetric but the drainage canals are often not functional due to poor  
39 maintenance. Most soils in Foum Gleita have formed from the schist parent rock that can be found  
40 at shallow depths (< 2.5m) (Van Asten et al., 2003). Soil texture is mostly silty clay loam, with  
41 around 30% clay. Although no specific clay mineralogical studies have been done in the Foum  
42 Gleita schemes, other studies in the Gorgol basin have shown that a large proportion of the clays are  
43 illites and kaolinites (Maignien, 1961; Audry, 1961). The soils have a moderate to strong structure,  
44  
45 varying from subangular blocky to prismatic in the more vertic soils. Biological activity in these  
46  
47 soils is generally high. The porosity of the soils is generally medium to high, but is distinctly lower  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 near soil surface. Although some soils have high ESP values, up to 65%, no evidence of soil  
2 structure degradation due to sodication has been observed in Foug Gleita. (Van Asten et al., 2003).  
3 In this region the cation exchange capacity (CEC) of soils ranges from 8 to 18 cmol.c.kg<sup>-1</sup>, and the  
4 electrical conductivity of the 1:5 paste (EC<sub>1:5</sub>) varies from 0.12 to 1.32 dS.cm<sup>-1</sup>. Percolation rates  
5 are generally below 4 mm day<sup>-1</sup>, and decrease during the rainy season as groundwater approaches  
6 the surface. On a few soils, high percolation rates were measured (> 10 mm day<sup>-1</sup>) and farmers  
7 complained about difficulties in maintaining a submersion needed for rice. After the cropping  
8 season, groundwater quickly drops to below 1.2 m, indicating that deep percolation into the  
9 vertically oriented bedrock occurs, suggesting that horizontal groundwater flow along the slopes is  
10 minimal (van Asten et al., 2003). The absence of a shallow groundwater level during fallow periods  
11 suggests that capillary rise from the ground water can generally be neglected as a source of  
12 (alkaline) salts. The irrigation water is 2 to 4 times more saline than at the other study sites (EC =  
13 0.17 dS.m<sup>-1</sup>), but the sodium adsorption ratio, defined as :  $SAR = Na^+ / (Ca^{2+} + Mg^{2+})^{1/2}$  where the  
14 concentration are in mmol.l<sup>-1</sup>, is still considered as very low (0.42). The residual sodium carbonate  
15 (RSC) is low and consequently suitable for irrigation according to the Riverside's diagram (see  
16 table 1). However, as at the other sites, RSC is positive and may therefore lead to alkalization and  
17 sodication.

### 2.1.2. Niono

21 The site of Niono, is typical of the inner Delta of river Niger, developed on three different  
22 geological units: (i) Infracambrian substratum (-500 to -480 M years) composed of fractured  
23 schists and sandstones, (ii) Continental Terminal of Post Eocene to Pre-Quaternary age (-2.5 to -2 M  
24 years) formed by alternate sedimentary deposits of clays, sandy clays and sand, (iii) recent  
25 quaternary alluvial sediments deposited by the river Niger. In the studied area, the main  
26 morphological units are sandy levees, loamy inter levee depressions and clayey settling basins.

1 The study is based on data obtained in the huge irrigation scheme of the Office du Niger in Mali  
2 near Ségou (14°32 N 05°59 W). The climate is semi-arid (sudano-sahelian) with an average rainfall  
3 of 460 mm/year, distributed from June to October. The annual evaporation reaches 2.5m which  
4 generates soil solution concentration and consequently alkalization processes related to the alkaline  
5 composition of irrigation water (Condom, 2000) . The distribution of the soils in this area is  
6 inherited from ancient delta deposits constituted of clayey depressions delimited by sandy levees.  
7 These soils evolved under hydrologic conditions imposed by the water management during last half  
8 century, especially related to the uprise of the groundwater level.  
9  
10 The soils in this study are located on a little levee. The upper soil layers are composed of 11% clay  
11 and 78% sand, and classified as loamy sand. The identified clay minerals are mainly kaolinite,  
12 smectite and traces of illite. The structure of the soil is polyhedral. The soil displays an important  
13 biological porosity in the upper part of the profile (0-70cm) due to roots and to meso-fauna activity.  
14 The CEC for this soil is 9 cmol<sub>c</sub>/kg but with large proportion of exchangeable sodium (ESP) of  
15 about 16% (table 2) ; the EC<sub>1:1</sub> for the soils in the area range from 0.18 to 1.09 dS.m<sup>-1</sup> . For the last  
16 30 years the fields have been mainly cropped in rice with market gardening as a marginal activity  
17 (Condom, 2000).

18  
19 In this large irrigation scheme plots are equipped with lateral drainage canals receiving mainly  
20 water from superficial flushing. After several decades of intensive irrigation, the water table level  
21 has risen approximatively by 50 m and is now close to the surface. Condom (2000) found that  
22 during the cropping season the average infiltration was about 80 mm (i.e. 9% of the leaching  
23 fraction), whereas during the inter cropping periods an equivalent amount of water from the water  
24 table supplied capillary rise towards the surface. Lateral losses towards drains and leaks represent  
25 26 % of the total water inputs (Condom, 2000).

26  
27 The Niger River that supplies the irrigation water by gravity has very low concentration (EC) and  
28 SAR values (Table 1), whereas the groundwater is 10 times more concentrated with a very high  
29

1 SAR value. According to the Riverside classification (Riverside salinity lab. staff, 1954) the quality  
2 of irrigation water does not constitute any harm to soil conservation but the composition of the  
3 groundwater typically represents a sodic hazard. Nevertheless, as illustrated by the positive  $RA_{\text{calcite}}$   
4 and RSC value (Table 1), these solutions can both become very alkaline and sodic when they  
5 concentrate, as the  $Ca^{2+}$  and  $Mg^{2+}$  might precipitate in minerals with low solubility like calcite and  
6 Mg-carbonates or Mg-silicates (e.g. sepiolite), whereas alkalinity and Na, keep concentrating in the  
7 solution (Vallès, 1991; Bethke, 2007).  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

### 19 2.1.3. Donaye

20 The third site, Donaye is representative of the irrigated soils of the river Senegal, representing a  
21 total surface area of irrigation schemes of 240,000 ha (Barbiero et al., 2001). These soils developed  
22 on marine sediments deposited during the Nouakchottian transgression (4300 BP) which extended  
23 380 km inland, near Bogue.  
24  
25  
26  
27  
28  
29  
30

31 The irrigation scheme is located in northern Senegal (N16°40'; W15°), in the middle valley of the  
32 Senegal River, near Podor. It is a 20-year-old irrigation scheme managed by the farmers, without  
33 drainage network. The soil which has for long time been described as formed by recent alluvial  
34 deposits is a vertisol (vertic Xerofluvent). However in a recent work Barbiero et al. (2005) have  
35 suggested that the soil distribution in lower and middle valley of river Senegal might result from the  
36 pedological transformation of a unique unripe mud, deposited during the last quaternary marine  
37 transgressions. The soils of the Senegal valley can be considered as a chronosequence, where the  
38 most recent soils can be observed downriver in the delta, and the most developed ones upriver, in  
39 the middle valley. The pedological process driven by the acidic conditions associated to the  
40 oxidation of inherited pyrite, led to the dissolution of the parental mud. The protonation and acid  
41 hydrolysis of the aluminosilicates in this material resulted in the formation of a sandy horizon,  
42 representing the quartz skeleton. In the upper part of the soil, the interaction of the river water  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

(through floods and irrigation ) with the acidic, Al and Si rich soil solution contributed to the precipitation of clay minerals, forming vertisols. In the Delta of the river Senegal this process is still active (Barbiero et al. 2005) and the vertisol is still thin whereas in Donaye it represents a two-meter-deep soil showing a vertic structure (slickensides), and constituted by 65 to 70% clay minerals, mainly beidellite and kaolinite (Favre et al., 2002), which shows a high cation exchange capacity of  $22.7 \text{ cmol.kg}^{-1}$ (table 2). The soil overlays a 5 to 6 m thick sandy layer corresponding to the afore mentioned quartz skeleton.

Rice cropping is performed by surge irrigation, and flooding is maintained during the complete plant growth cycle (about 100 days). The presence of a shallow water table at 2m depth contributes to air entrapment and stops vertical infiltration (Hammecker et al. 2003). Consequently, vertical leaching is negligible, and soil solution in the upper part of the profile (above 40 cm) does not connect with the water table during the irrigation period. However, bad maintenance of bunds aimed to hold the irrigation in the paddy plots, generates important leaks contributing to non negligible loss of water, especially during the first irrigation. Irrigation water is pumped in a loop of river Senegal called Doué. This water with a very low concentration ( $\text{EC} = 0.07 \text{ dS.m}^{-1}$ ) and a low SAR value (Table 1), is suitable for irrigation according to Riverside's diagram. Nevertheless, like in the former situations, the  $\text{RA}_{\text{calcite}}$  and RSC of the irrigation water are positive, and the concentration of this solution may lead to sodication.

## 2.2. Numerical simulation.

### 2.2.1. The model and its main parameters.

The evolution of the soil solution and its interactions with the solid phase (precipitation-dissolution and exchange processes) were evaluated with the computer model PHREEQC v2. (Parkhurst and Appelo, 1999). This model, based on aqueous thermodynamics, computes equilibrium chemistry of aqueous solutions interacting with minerals, gases, solid solutions, exchangers, and sorption

1 surfaces, providing a full speciation of an aqueous system. The different databases provided with  
2 the model give a very wide possibility for computing the activities of most of the aqueous species  
3 present in natural solutions, especially for the  $\text{CO}_3\text{-HCO}_3$  solutions. This model is especially useful  
4 as, in addition to the calculation of the chemical equilibria, it allows for the simulation of mixing  
5 different solutions, for concentrating solution due to evaporation, and for adding or taking out  
6 aqueous species from the soil solution. Using these options, it is possible to simulate solution  
7 changes due to flooded irrigation combined with concentration of the soil solution and evaporation.  
8 In surge flooding conditions, successive water inputs are usually performed to maintain a constant  
9 ponding level.

10 Considering the duration of the rice cycle, the average evapotranspiration and the soil porosity,  
11 Hammecker et al. 2003 found that about 17 irrigations were necessary during one growing cycle.  
12 Considering that in the PHREEQC model all the calculations are performed for a given volume of  
13 solution, the Cation Exchange Capacity (CEC) and exchangeable cations were expressed per  
14 volume of liquid phase, depending on the soil porosity. The mass balance for water and solutes was  
15 calculated in the root zone, i.e. the first 20 centimeters of the soil profile (Slaton et al., 1990;  
16 Sharma et al., 1994; Boivin et al.2002). The irrigation water management during the cropping  
17 season, was simulated for one liter of soil solution in the root zone and the corresponding water  
18 volume for the superficial ponding water. As depicted in figure 2, during the first irrigation of the  
19 season 3 liters are introduced into the system : 1 liter for the soil solutions, and two liters for  
20 superficial ponding. During the cropping season, the superficial water level can almost disappear  
21 completely, but farmers usually irrigate before the complete drying-up of the plot.

22 Different hydrological conditions, typical of field situations, were considered to simulate the  
23 geochemical evolution taking into account different leaching fractions, presence or absence of  
24 capillary rise from an aquifer with different composition, and the influence of plant uptake of  
25 nutriment. The actual leaching fraction (LF) being considered is the fraction of applied irrigation

water that drains below the root zone :

$$LF = \frac{D_d}{D_a} \quad [1]$$

Where  $D_d$  [L] is the amount of water drained below the root zone and  $D_a$  [L] the amount of irrigation water applied.

### 2.2.2. Chemical equilibria

Ionic activities in the soil solution were computed according to Debye-Hückel expression, based on the ionic strength of the solution. The soil solution was equilibrated with exchange sites on clay minerals and organic matter following Gaines and Thomas (1953) convention and mass conservation. Equilibrium with gas and minerals were considered for those initially present in soil, and those likely to crystallize or to affect the evolution of soil solution. The minerals usually found in arid soils like Mg-calcite, gypsum, quartz, kaolinite and sepiolite, were allowed to precipitate and dissolve, consequently to modify the composition of the solution. As quartz is initially present in the soil, this mineral has been chosen rather than amorphous silica, though the two options were tested and no differences in the major results of these simulations were found. When Al in solution analysis was not available it was computed considering equilibrium with kaolinite at the pH of the soil solution. The main solid and gaseous phases used in these simulations are reported in table 3, with their dissociation equation, and solubility products (K). Partial CO<sub>2</sub> pressure was considered to be equal to atmospheric value (log pCO<sub>2</sub> = -3.5 atm) whereas in the soil solution it was considered to be higher and constant, with log pCO<sub>2</sub> = -1.8 atm, like in most of the tropical soils (Brook et al. 1983). The crystallization of the different minerals was considered to take place when the solution reached equilibrium, regardless of any eventual super-saturation or kinetics phenomenon. However as the simulation is performed over a very long term, with different water contents, these processes will scarcely affect the final result.

### 2.2.3 The hydrological parameters

1 The adopted procedure for calculation was first, to equilibrate the irrigation solution with the soil  
2 solution, the mineral phases, and the partial CO<sub>2</sub> pressure in soil (10<sup>-1.8</sup> atm). The resulting solution  
3 was then concentrated to 5% of its original volume, as the farmers don't let the plot dry out  
4 completely. This value has been estimated from experimental observations (Hammecker, non  
5 published), though it has been verified that the choice of this value has little influence on the final  
6 result. During this procedure equilibria with the solid phase were computed considering the soils  
7 being initially formed by quartz, kaolinite, beidelite and by their respective exchangeable cations  
8 (table 2).  
9

10 Finally, depending on the leaching fraction (LF), the remaining water content after evaporation  
11 (Q<sub>0</sub>) was added to a new amount of irrigation water (Q<sub>1</sub>), in order to start a new simulation with the  
12 same amount of water (one liter), following this relation :  
13

$$Q_0 = (1-LF) \times 0.05 \quad [2a]$$

$$Q_1 = 1 - Q_0 \quad [2b]$$

14 This procedure was performed 17 times for one cropping cycle, and 60 cropping cycles were  
15 simulated. In order to take into account the capillary rise in these geochemical processes, we  
16 estimated the quantity of solution possibly risen from the water table towards the root zone with  
17 water flows derived from hydraulic gradients measured during the dry period (Condom, 2000). As  
18 capillary rise occurs only during the inter-cropping period, we treated both periods together, by  
19 mixing the two solutions (irrigation and water table) in their respective proportions (90% and 10%)  
20 in the root zone.  
21

22 In addition to vertical drainage, superficial lateral drainage (flushing) is performed intentionally or  
23 accidentally by the farmers in some situations. During this type of water management all salts and  
24 concentrated soils solutions present in the superficial layers of soil are dissolved and partially  
25 evacuated with the first water input at the beginning of the cropping period. As the soil profile dried  
26



out during the inter-cropping season most of the soluble salts have accumulated in surface.

1 Superficial lateral drainage experiment performed in Donaye showed that just with natural leakage  
2  
3 from the bunds, an average of 15% of the total salinity (0-1m) could be removed (Hammecker et al.  
4  
5 1999). In Niono the flushing (lateral drainage) is performed intentionally, at the beginning of each  
6  
7 new cropping season whereas in Donaye it occurs accidentally as, the margins of the plots leak after  
8  
9 the inter cropping period when the soil shrinks and the margins crack. Consequently, we simulated  
10  
11 an input of irrigation before each cropping period. This water was equilibrated with the soil  
12  
13 solution, the minerals and the exchangeable cations, and was then partially evacuated from the  
14  
15 system (Fig. 3). This numerical treatment, simulates the exportation of elements during a flush  
16  
17 (lateral drainage), where flushing rates were defined as the proportion of water exported from the  
18  
19 system, after the first water input had equilibrated with soil. Equilibrium during this procedure is  
20  
21 conditioned by different physico-chemical processes like : (i) the competition between convective  
22  
23 downward flow during infiltration and upward molecular diffusion towards the superficial water  
24  
25 layer, (ii) the kinetics of minerals dissolution , (iii) the lateral losses in the plot. These processes  
26  
27 were not considered individually but integrated into a bulk value of flushing rate. Different  
28  
29 theoretical flushing rates were tested (5%, 10%, 30% and 50%) as no precise measurements for  
30  
31 actual superficial lateral drainage were available in the three conditions. However they are in  
32  
33 concordance with the flushing rate measured by Hammecker et al.1999 and with those determined  
34  
35 in the experiment performed by Haefele et al. (1999) in the same area, who that showed superficial  
36  
37 lateral drainage after submersed ploughing (puddling) at the beginning of the cropping season,  
38  
39 could increase this salt exportation four folds.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

#### 53 2.2.4. Plant uptake

54  
55  
56 The geochemical budget was considered with and without the incidence of the plant uptake, in order  
57  
58 to quantify the contribution of the plant uptake to soil degradation or conservation processes. The  
59  
60

main threat being alkalization due to carbonate concentration (pH increase), the focus has been set on the ability for rice to export alkalinity, and the main related cations (Ca and Mg) possibly affecting the carbonate balance and ESP. Data of Dobermann and Fairhurst (2000) on the average final composition of rice were used (Table 4). We considered an average yield of 5t/ha, which corresponds to 10t/ha of dry biomass(DB), a soil depth (z) of 0.2 m, an average bulk density ( $\rho_b$ ) of 1750 kg.m<sup>-3</sup> and a mineral density ( $\rho_s$ ) of 2650 kg.m<sup>-3</sup>.

The average ratio between the dry biomass (DB) and the volume of the soil solution ( $V_{soil}$ ) has been calculated to be 14.7 kg.m<sup>-3</sup> as follows :



$$\frac{DB}{V_{soil}} = \frac{DB \cdot 10^3}{z \cdot 10^4 \cdot \theta_s}$$

where  $\theta_s$  is the porosity. As the soil profile is saturated porosity is defined as:

$$\theta_s = 1 - \frac{\rho_b}{\rho_s}$$

Plants export alkalinity because they absorb more cations than anions. In order to maintain electrical neutrality both inside and outside of the roots they have to produce organic anions and excrete protons (Pinton et al.,2001 ; Rengel, 2003). The average plant alkalinity was calculated as the difference between the cations and anions absorbed by the plant (Table 4). It was considered that all plant alkalinity was removed from the field, as farmers export the entire plant for threshing.

Moreover, when a crop is considered in the geochemical system , it is also necessary to take into account the N cycle. In paddy fields the plant absorb N as  $NH_4^+$  originating from the deamination of the organic matter and the hydrolysis of the urea used as N fertilizer:



In both cases for each mole of  $NH_4^+$  produced one mole of protons is removed from soil solution.

Part of the ammonium produced is also adsorbed on the exchangeable complex or lost as ammonia.

Therefore we only took into account the ammonium ions absorbed by the plant, by using the total N content of the plant (Table 4), irrespectively of the actual urea management in these sites. The net acidification due to plant uptake and the N cycle was calculated as:

$$\text{proton produced} = \text{plant alkalinity} - \text{N uptake.}$$

In order to simulate the exportation of alkalinity, hence the acidification of the soil solution due to proton excretion, potassium bicarbonate ( $\text{KHCO}_3$ ), magnesium and calcium carbonate ( $\text{MgCO}_3$ ,  $\text{CaCO}_3$ ) and potassium chloride (KCl) were numerically removed from the irrigation solution in proportions, calculated to meet the corresponding alkalinity uptake (Table 4). The total uptake for the different ions was divided by 17 (i.e. the number of irrigation events during a cropping cycle) and distributed evenly over the entire cropping season.

### 3. Results and discussion.

#### 3.1. Evolution of the irrigation solutions with increasing concentration.

Results displayed in Figure 4 show the evolution of the solution composition versus the concentration rate defined as :  $\text{concentration rate} = (V_i - V) / V_i$ , where  $V_i$  is the initial volume of solution and  $V$  the volume of solution reached after concentration.

These results showed an important increase in pH with concentration, up to values around 9, and clearly superior to 8.3 the pK of calcite. Irrigation water of Foug Gleïta, rapidly reached saturation with calcite for its large initial Ca concentration. Consequently Ca concentration in solution decreased constantly whereas Alkalinity, Na and Mg concentrations increased, despite a slight depletion in Mg when sepiolite crystallization started. At the same time pH increased drastically despite sepiolite precipitation involving a release of protons into the solution (Table 3). Finally even for very high concentration rates, Mg increased noticeably in the solution because of its high initial

content, as sepiolite crystallization is controlled by Si, which is less abundant.

1 During this simulation, the irrigation water of Niono rapidly reached saturation with quartz and  
2  
3 while it is the only mineral precipitating, alkalinity and concentrations in all the cations increased  
4  
5 regularly with concentration rate. The solution reached saturation with calcite for a high  
6  
7 concentration rate (0.75) then Ca concentration decreased. For higher concentration rates, Mg  
8  
9 concentration dropped when sepiolite crystallized whereas Na kept concentrating. A similar  
10  
11 behavior was observed in Donaye, when water of river Senegal concentrated. In the three cases the  
12  
13 pH of irrigation water increased and Na became the dominant cation in solution, enhancing the risk  
14  
15 of sodication. Considering the SAR and the Electrical Conductivity of the irrigation water of the  
16  
17 different situations, they can be considered as suitable for irrigation according to the Riverside  
18  
19 diagram (1954). However this static view does not reflect the evolution of the solutions that may  
20  
21 occur when concentrating. Numerical simulation of the concentration of these solutions (Figure 5),  
22  
23 show that their SAR increased drastically with concentration, whereas the total salinity (EC) didn't  
24  
25 increase as much, mainly because of the precipitation of calcite. Barbiero et al. (2004) showed  
26  
27 some results for experimental concentration of irrigation water from River Niger, in the Republic of  
28  
29 Niger, which have similar initial composition as the irrigation water of Niono, where the SAR of  
30  
31 the solutions also increased but with much higher EC values until reaching the C4S4 domain,  
32  
33 instead of C2S4 as in the numerical simulation (Fig. 5). This discrepancy should mainly be  
34  
35 explained by the fact that in the simulation the kinetics of precipitations have not been taken into  
36  
37 account, and especially the retardation induced by the presence of Mg in the same proportions as Ca  
38  
39 (House et al. 1988). However, in both cases the simulation shows that resulting solutions are  
40  
41 extremely hazardous for soil conservation as they tend to promote sodication.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

### 54 3.2. Evolution of the soil solutions over 60 simulated cropping cycles.

55 The previous simulations described the evolution of the irrigation solution without taking into  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

account the interactions with soils, especially with the exchange complex and the minerals.

Moreover, in order to get an overall idea of the geochemical processes involved in field conditions, the vertical drainage and capillary rise was taken into account, following the simulation procedures described previously for the three situations.

The first set of simulation was aimed at quantifying the influence of the presence of a soil with a reactive adsorption complex and minerals, as well as the influence of the vertical leaching towards the water table. As the concentration of the soil varies constantly between two irrigations, the results displayed in the following figures corresponds to the composition of the soil solution and the exchange complex, immediately after the irrigation i.e. when the plots are full of water, in order to represent the actual solution infiltrating into the soil.

In the case of Foug Gleita (Fig. 6) where irrigation water tends to become very alkaline with concentration, an equivalent trend was observed in presence of soil and vertical drainage. However this trend appeared in lesser proportions, as pH reached only a value of 7.8 after 60 cropping cycles. The major ions in solution were  $\text{Na}^+$  and alkalinity. Both increased regularly with the number of cropping cycles whereas  $\text{Ca}^{2+}$  decreased as calcite precipitated. The complex reacted to compensate this trend, providing  $\text{Ca}^{2+}$  to the soil solution. Consequently, the proportion of exchangeable Ca decreased and exchangeable Mg became dominant. At the same time, exchangeable Na increased and consequently ESP rose to 15% after 10 cropping cycles and finally reached 30% after 60 cropping cycles. The evolution of the pH and alkalinity is still positive, but at a noticeably lower rate than for the concentration of the irrigation water alone. In fact, the large CEC of the soil buffered the effects of irrigation solution composition, especially as it imposed the geochemical composition of the soil solution, and supplied sufficient  $\text{Ca}^{2+}$  ions necessary to precipitate calcite and so prevented the accumulation of excessive alkalinity. According to the Residual Alkalinity (RA), which is still positive, the general trend of the system is to evolve continuously, but slowly, towards alkaline conditions.

1 In the case of Niono (Fig. 7), the simulation showed a continuous increase in pH during the first 35  
2 cropping cycles followed by a stabilization at around pH=8.1. Alkalinity and  
3 Na<sup>+</sup> in soil solution and in the soil displayed a similar evolution, namely increasing until reaching  
4 saturation towards sepiolite. RA<sub>calcite</sub> followed the same increasing trend as alkalinity, witnessing an  
5 alkaline evolution. Exchangeable sodium increased drastically during the first cropping cycles and  
6 reached a maximum of 53% of the exchangeable complex after 50 cropping cycles, until sepiolite  
7 started to crystallize. Although extreme values for ESP have not actually been observed in this  
8 region, they are indicative of the type of degradation the soils might undergo in these conditions :  
9 high pH and ESP values are typical of alkalization and sodication processes. Although the irrigation  
10 solution is potentially less alkaline than in Foum Gleita, its use for irrigation is more hazardous,  
11 because the adsorption complex is very vulnerable to soil solution composition due to low CEC ;  
12 the exchange between the adsorbed Ca<sup>2+</sup> and Na<sup>+</sup> in solution will therefore lead to a rapid increase  
13 in ESP.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 Finally, in the case of Donaye (Fig. 8), the pH of the soil solution did not change much with the  
33 number of cropping cycles and mainly remained near neutrality. Sodium and magnesium were the  
34 major cations in the soil solution, and both concentrations increased with the number of cropping  
35 cycles. Alkalinity was the main anion, but both chloride and sulphate increased drastically with  
36 time, as there was no leaching towards the water table in this site. The soil solution temporarily  
37 reached equilibrium with gypsum during the dry season when all the irrigation water was  
38 evaporated, and especially during the inter cropping periods, according field observations (Boivin et  
39 al., 2002; Barbiero et al., 2005).  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

52 In the first instance RA<sub>calcite</sub> dropped slightly but increased again progressively when calcite and  
53 later gypsum started to precipitate. As the soil is very clayey with high CEC, the composition of the  
54 soil solution was determined by the buffering property of the soil. Exchangeable calcium was  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 progressively desorbed and precipitated, exchangeable magnesium increased and became the first  
2 exchangeable cation. At the same time ESP increased linearly up to 9% after 60 cycles. These  
3 extreme values for adsorbed magnesium were not observed yet, but still represent the values  
4 potentially reached for these conditions. Sixty cropping cycles were not sufficient to lead to alkaline  
5 conditions. However these confined conditions promote a very slow alkalization and sodication  
6 process.  
7  
8  
9  
10  
11

### 12 3.3. Importance of plant uptake

13 Results displayed in figure 9 to 11 showed clearly that the ionic plant uptake in the soil solution,  
14 decreased systematically the pH of soil solution in the three sites. Nevertheless, the plant effect was  
15 moderate for the most alkaline conditions, namely Niono where pH still increased with time, up to  
16 7.75 after sixty cropping cycles. In the case of Foum Gleita and Donaye pH decreased until  
17 reaching a final value of 7.2 as plant uptake counterbalanced the alkaline input of irrigation water.  
18 The total concentration of the soil solution, as indicated by EC, decreased by at least 30% , almost  
19 50% for Foum Gleita and Niono. In Donaye, plant uptake played a major role for it is the only  
20 process exporting elements out of the system, as there was no vertical drainage. The same trend was  
21 observed for ESP in the cases of Foum Gleita and Donaye, where plant uptake contributed to a  
22 reduction by 50% of exchangeable sodium and consequently reduced the sodic risk. In Niono,  
23 unlike the former cases, plant uptake had a moderate effect on ESP. In fact, as  $K^+$ ,  $Ca^{2+}$  and  $Mg^{2+}$   
24 were preferentially absorbed by rice to  $Na^+$ , its concentration increased relatively in soil solution  
25 and exchangeable position. In all cases,  $RA_{calcite}$  of the soil solution was significantly reduced, and  
26 even showing a serious decrease for Donaye until reaching  $0.0015 \text{ mol.l}^{-1}$ , though still remaining  
27 positive.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

57 This simulation showed that in alkaline geochemical conditions, total exportation of rice (grains +  
58  
59  
60  
61  
62  
63  
64  
65

1 straw) improved generally the composition of the soil solution and tended to reduce the risks of  
2 salinisation, alkalization and generally sodication except in the case of Niono where the low CEC  
3 was very sensitive to a relative Na increase in the soil solution. This situation is illustrated by  
4  
5 Barbiero et al. 2001 who carried out an experiment in Niger (Lossa) where the cultivation of a local  
6 fodder grass (*Echinochloa stagnina*) and its exportation out of the field contributed to noticeable de-  
7  
8 alkalization of a highly alkaline soil.  
9  
10  
11  
12  
13  
14

### 15 3.4. Effects of superficial lateral drainage (flushing). 16

17 Results displayed in figures 12 to 14 showed distinctly that even for low rates, superficial flushing  
18 reduced significantly EC (i.e. the total salinity) of the soil solution, as well as ESP. The incidence of  
19 lateral drainage on pH and on RAC was less clear, especially for the case of Donaye as both  
20 parameters used to decrease without superficial lateral drainage and whereas they tended to  
21 increase slightly or stay constant when lateral drainage was simulated. However, in the case of  
22 Foug Gleita (Figure 12) the processes of sodication and alkalization were thoroughly reduced even  
23 for the smallest lateral drainage rate, as pH, EC, ESP and  $AR_{\text{calcite}}$  decrease with increasing flushing  
24 rate. For flushing rates greater than 30% the simulation shows that all these parameters reach  
25 constant low values, namely  $pH = 7.3$ ,  $EC = 0.6 \text{ dS.m}^{-1}$ ,  $ESP = 2\%$ ,  $AR_{\text{calcite}} = 0.03 \text{ mol.l}^{-1}$ ,  
26  
27 witnessing stable geochemical conditions, harmless for the quality of soil and solution.  
28  
29

30 In the case of Niono (figure 13), superficial flushing was as effective as in the previous example,  
31 for its incidence was clearly noticeable, even for low flushing rates. The simulation showed that a  
32 flushing rate of 30% was sufficient to control and maintain EC at a constant level, with only a slight  
33 increase in ESP. The incidence of lateral drainage on pH and  $AR_{\text{Calcite}}$  was very important for the  
34 low rates (5% and 10%) during 20 first cropping cycles, as these parameters decreased slightly,  
35 until  $AR_{\text{calcite}}$  becoming negative for 5% flushing rate, and finally increased constantly. For the  
36 higher flushing rates (30 % and 50 %) the incidence on pH and  $AR_{\text{calcite}}$  was less important for the  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 first simulated cropping cycles, but remained constant over the entire period. This special behavior  
2 is probably related to the difference in  $p\text{CO}_2$  between the soil solution ( $10^{-1.8}$  atm) and the  
3 supplementary irrigation water ( $10^{-3.5}$  atm) added to compensate lateral drainage, and their relative  
4 proportions when mixed, promoting whether calcite precipitation or dilution of the soil solution. In  
5 any cases the superficial flushing reduced significantly the risk of alkalisation and sodication,  
6 though ESP values still reached relatively high values (17%).  
7  
8  
9  
10  
11  
12

13 An equivalent behavior was noticed in the simulation for the case of Donaye (figure 14), although  
14 no vertical drainage was considered; EC and ESP showed a significant decrease even for low  
15 flushing rates and reached a low constant value all along the cropping cycles when higher  
16 superficial lateral drainage rates were achieved (EC =  $0.7 \text{ dS}\cdot\text{m}^{-1}$  ; ESP = 1.5 % for 30% flushing  
17 rate and EC =  $0.6 \text{ dS}\cdot\text{m}^{-1}$  ; ESP = 0.8 % for a flushing rate of 50 %). Consequently the soil solution  
18 and soil quality increased. On the other hand, pH and  $\text{AR}_{\text{calcite}}$  showed a significant decrease when  
19 no lateral drainage was simulated whereas they showed a similar behavior for the different flushing  
20 rates as described for the case of Niono. However these variations of pH and  $\text{AR}_{\text{calcite}}$  can be  
21 considered as negligible considering alkalization because for the worst condition (flushing rate of  
22 10 %), pH reached 7.33 and  $\text{AR}_{\text{calcite}}$  reached  $0.004 \text{ mol}_c\cdot\text{l}^{-1}$  after 60 cropping cycles .  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 Moreover these results also show that  $\text{AR}_{\text{calcite}}$  , which is considered as a conservative tracer  
38 (Ribolzi et al. 1996; Barbiero et al. 2001; Barbiero et al. 2004), varies along the simulation, and  
39 probably also in field conditions, as solutions with different  $p\text{CO}_2$  are mixed.  
40  
41  
42  
43  
44

45 In Donaye, mineralogical and chemical experimental studies of the soil have been performed to  
46 assess the modifications of the soil quality after 15 years of intensive irrigation. When compared  
47 inside an irrigated plot and outside the irrigation scheme, the evolution of the soil exchangeable  
48 complex, especially the ESP, has been found not significantly different (Charolais and Weber,  
49 1994). Equally, when the pH of the soil, the EC and the content of accumulated carbonates in the  
50 topsoil have been measured on a 1/50 aqueous extract, (Schiess, 1995; Boivin et al. 1998) only a  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

slight increase in the soil between the two situations was noticeable, though the difference isn't really statistically different between the two groups (figure 15). These experimental results show that in 15 years of irrigation the soil quality has not changed noticeably. In Donaye the rice straw is usually not exported from the field, but left to graze for cattle or burned. However, “natural” leakage throughout the bunds of the plot, due to a lack of maintenance is almost systematically observed. In fact, this disorder actually contributes to the superficial flushing, especially during the first irrigation when the soil shrinkage cracks are still open in bunds, and helps to preserve the soil quality despite the improper quality of irrigation water.

Although the quantitative effect of superficial flushing is difficult to model accurately because of the lack of information, this study demonstrates clearly the crucial role of superficial flushing for soil conservation, and attests the general conclusions of the numerical simulation results.

### 3.5. Effect of soil reduction.

Although this process has not been modelled specifically in this work for the lack of Eh information in some cases, it is still possible to evaluate its effect on the proton balance. In the case of prolonged submersion conditions, like in paddy fields, the soil undergoes reduction as microbial activity consumes the dissolved oxygen, and finally reduces the iron oxi-hydroxides (the most commonly found). The oxidation of organic matter by reducing Fe(III) to Fe(II) produces alkalinity and consequently an increase in pH :



When the system oxidizes at the end of the submersion period an equivalent amount of acidity is produced and the general proton balance is null in confined conditions. However in case of a leaching fraction during reduced conditions, reoxidation leads generally to ferrolysis phenomenon which is a soil transformation involving a protonation of the complex, an attack of the clay crystalline structure, where  $\text{H}^+$  is penetrating the octahedral layers and taking the place of  $\text{Al}^{3+}$  as

described by Brinkman (1979) and revisited by van Ranst and De Coninck (2002). Consequently, soil reduction has either no effect on alkalinity or, on contrary, consumes alkalinity and reduces pH in the studied situations.

## Conclusion

In this study we used PHREEQC to predict the evolution of the soil and soil solution over several years, with different soil and water management. Although the use of Riversides' diagram,  $RA_{\text{calcite}}$  and the RSC index, can be informative about the potential risks, they cannot predict the evolution of the soil nor of the soil solution. Especially as they don't take into account the buffer capacity of soil, the water management involving dilutions and variations in  $p\text{CO}_2$ , and the effect of the plants uptake. We demonstrated that despite the high risk of alkalisation and sodication according to these indicators, in three west African situations of irrigation, soil degradation is not a fate. Considering the buffer effect of the soil (very high in Senegal), the possibilities of removing alkalinity from the plots by straw management, and finally the use of superficial flushing whether naturally (Foum Gleita), artificially with an effective drainage network (Niono) or accidentally with leaks in the bunds (Donaye), alkalisation or sodication is not jeopardizing soil properties in well-conducted rice cropping system. These results using geochemical numerical modelling, attest that in Sahel, although the main rivers (river Senegal, Niger, Gorgol Noir) show an alkaline composition with positive  $RA_{\text{calcite}}$  and RSC values, their use for irrigation in paddy fields cannot be considered as directly responsible for soil alkalisation. Barbiero et al. (1998), in the Republic of Niger, or van Asten et al. (2004) in Foum Gleita, found similar conclusions based on experimental results. The presence of alkaline soils is more probably due to the alteration of an alkaline bed rock (in Foum Gleita) or to the presence of groundwater leaching through a deeper alkaline bed rock (in Niono).

Unlike the usual assumptions, where rice cropping is held responsible for soil salinization or  
alkalinisation, this study shows that in these cases, rice cropping can improve the soil quality when  
irrigation and residue management are performed properly. Namely a superficial lateral drainage  
performed once, at the beginning of the cropping season, although being sometimes seen as a waste  
of water by the local farmers, improves noticeably the sustainability of the system despite adverse  
composition of the irrigation water. Moreover, in the situation of alkalinisation risk, the exportation  
of the straw out of the plots contributes significantly to the control of the average alkalinity of soil,  
especially in the cases of Niono and Donaye.

Table 1: Composition of irrigation water (I) and groundwater (GW) in the different locations

	<b>Ca</b>	<b>Mg</b>	<b>Na</b>	<b>K</b>	<b>Alk.</b>	<b>Cl</b>	<b>SO<sub>4</sub></b>	<b>RSC</b>	<b>pH</b>	<b>SAR</b>	<b>EC</b>
	<b><math>\times 10^{-3} \text{ mol}_e \cdot \text{l}^{-1}</math></b>					<b><math>\text{dS} \cdot \text{m}^{-1}</math></b>					
Foum <sup>I</sup>	0.78	0.38	0.32	0.12	1.4	0.11	0.09	0.24	7.54	0.42	0.17
Niono <sup>I</sup>	0.11	0.1	0.17	0.05	0.38	0.08	0	0.17	6.52	0.62	0.04
Niono <sup>GW</sup>	0.62	0.36	6.05	0.09	6.91	0.13	0.08	5.93	7.69	8.62	0.58
Donaye <sup>I</sup>	0.23	0.12	0.1	0.06	0.61	0.06	0.07	0.26	7.28	0.25	0.07

1 Table 2: Adsorbed cations of the different soils. ESP : exchangeable sodium percentage.  
 2  
 3  
 4

	<b>XCa</b>	<b>XMg</b>	<b>XNa</b>	<b>XK</b>	<b>Σcat</b>	<b>ESP</b>
	<b>cmol<sub>c</sub>.kg<sup>-1</sup></b>					<b>%</b>
Foum Gleita	11.8	4.8	0.3	0.6	17,5	1.71
Niono	5.5	2	1.5	0.1	9.1	16.48
Donaye	13	9	0.4	0.3	22.7	1.76

Table 3 : Minerals dissociation equations and solubility product values at 10<sup>5</sup> Pa and 298.15K (1) :  
 PHREEQC data base, (2) : estimated as a solid solution with data from PHREEQC data base, (3) :  
 Kindis data base (Made, 1991) , (4) : fixed (Brooks et al. 1983)

Minerals	equations	log K (25 °C)
CO <sub>2</sub>		-1.8 <sup>(4)</sup>
Mg-Calcite	$\text{Ca}_{.98}\text{Mg}_{.02}\text{CO}_3 = .98 \text{Ca}^{2+} + .02\text{Mg}^{2+} + \text{CO}_3^{-2}$	-8.46608 <sup>(2)</sup>
Sepiolite	$\text{Mg}_2\text{Si}_3\text{O}_7.5\text{OH} \cdot 3\text{H}_2\text{O} + 4 \text{H}^+ + 0.5\text{H}_2\text{O} = 2 \text{Mg}^{2+} + 3 \text{H}_4\text{SiO}_4$	15.76 <sup>(1)</sup>
Kaolinite	$\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4 + 6 \text{H}^+ = \text{H}_2\text{O} + 2 \text{H}_4\text{SiO}_4 + 2 \text{Al}^{3+}$	7.44 <sup>(1)</sup>
Gypsum	$\text{CaSO}_4 \cdot 2\text{H}_2\text{O} = \text{Ca}^{2+} + \text{SO}_4^{-2} + 2 \text{H}_2\text{O}$	-4.58 <sup>(1)</sup>
quartz	$\text{SiO}_2 + 2 \text{H}_2\text{O} = \text{H}_4\text{SiO}_4$	-3.98 <sup>(1)</sup>
beidellite	$\text{Si}_{3.8}\text{Al}_{1.7}\text{Fe}_{.3}\text{Mg}_{.2}\text{O}_{10}(\text{OH})_2\text{Ca}_{.2} + 6.8\text{H}^+ + 3.2 \text{H}_2\text{O} =$ $3.8\text{H}_4\text{SiO}_4 + 1.7\text{Al}^{3+} + .3\text{Fe}^{3+} + .2\text{Mg}^{2+} + .2\text{Ca}^{2+}$	-33.98 <sup>(3)</sup>

Table 4 : Average paddy uptake (grain + straw) and equivalent uptake in soil solution (average yield of 5t/ha, and average bulk density of 1750kg.m<sup>-3</sup>)

	Dobermann & Fairhurst, 2000	equivalent ionic	
	%	species	mmol/l
N	0.65	NH <sub>4</sub> <sup>+</sup>	6.62
K	1.4	K <sup>+</sup>	5.1
Mg	0.2	Mg <sup>2+</sup>	2.35
Ca	0.3	Ca <sup>2+</sup>	2.13
Na	0.2	Na <sup>+</sup>	1.24
Σc			18.01
P	0.1	H <sub>2</sub> PO <sub>4</sub> <sup>-</sup>	0.46
Cl	0.5	Cl <sup>-</sup>	2.01
S	0.075	SO <sub>4</sub> <sup>2-</sup>	0.66
Σa			3.24
Σc - Σa		Alkalinity	14.3



## References

- 1  
2  
3 Al Droubi, A., Fritz, B., Gac, J-Y. and Tardy, Y., 1980. Generalized residual alkalinity concept;  
4  
5 application to prediction of the chemical evolution of natural waters by evaporation. Am. J.  
6  
7  
8  
9  
10  
11 Arbol, I.P., Yadav, J.S.P. and Massoud, F.I., 1988. Salt-affected soils and their management. Food  
12  
13 and Agricultural Organization of the United Nations, Soils Bulletin 39, Rome 131 pp.  
14  
15  
16 Audry, P., 1961. Etude pédologique du cercle de Guidimaka (République Islamique de Mauritanie).  
17  
18 Etude du milieu naturel et étude pédologique. - Rapp. Off. Rech. sci. tech. O-mer, Dakar, 173  
19  
20  
21 p. multigr.,  
22  
23 Barbiero, L. and Van-Vliet-Lanoe, B., 1998. The alkali soils of the middle Niger Valley; origins,  
24  
25 formation and present evolution. Geoderma, 84: 323-343.  
26  
27  
28 Barbiero, L.; Valles, V.; Regeard, A., Cheverry, C., 2001 Residual alkalinity as tracer to estimate the  
29  
30 changes induced by forage cultivation in a non-saline irrigated sodic soil Agricultural Water  
31  
32 Management. 50 : 229-241  
33  
34  
35 Barbiero L., Ould Mohamedou A., Laperrousaz C., Furian S., Cunnac S., 2004. Polyphasic origin of  
36  
37 salinity in the Senegal delta and middle valley. Catena 58, 101-124.  
38  
39  
40 Barbiero L., Ould Mohamedou A., Roger L., Furian S., Aventurier A., Rémy J.C., Marlet S., 2005.  
41  
42 The origin of vertisols and their relationship to Acid Sulfate Soils in the Senegal Valley.  
43  
44 Catena 59, 93-116.  
45  
46  
47 Bertrand, R., Keita, B., Ndiaye, K.M., 1993. La dégradation des sols des périmètres irrigués des  
48  
49 grandes vallées sud-sahariennes: Cas de l'Office du Niger au Mali). Cahiers Agricultures 2,  
50  
51 318 329.  
52  
53  
54 Bethke, C.M. 2007. Geochemical and Biogeochemical Reaction Modeling. Cambridge University  
55  
56 Press. 564 pp.  
57  
58  
59  
60  
61  
62  
63  
64  
65

Boivin, P., Favre, F. and Maeght, J-L., 1998. Les sols de la moyenne vallée du fleuve Sénégal :

caractéristiques et évolution sous irrigation. *Étude et Gestion des Sols*, 5: 235- 246

Boivin, P., Favre, F., Hammecker, C., Maeght, J-L., Delarivière, J., Poussin, J-C. and Wopereis,

M.C.S., 2002. Processes driving soil solution chemistry in a flooded rice-cropped vertisol:

analysis of long-time monitoring data. *Geoderma*, 110: 87-107

Bolan N.S. and Hedley M.J., 2003. Role of carbon, nitrogen, and sulfur cycles in soil acidification.

In: Z. Rengel (Editor), *Handbook of soil acidity*. Marcel Dekker, New York, pp. 29-56.

Brinkman, R. 1979. Ferrollysis, a soil-forming process in hydromorphic conditions. Doctoral thesis.

Agricultural Research Reports 887, Centre for Agricultural Publishing and Documentation,

Wageningen.

Brooks, G.A., Folkoff, M.E., Box, E.O. 1983. A global model of soil carbon dioxide. *Earth Surface*

*Processes and Land/onus* 8:79-88.

Carité, D. 1989. *Géologie en Mauritanie*. Edisud. 284 pp.

Charolais, M., Weber, V., 1994. Evolution des sols irrigués dans la moyenne vallée du fleuve

Sénégal. M.Sc. Thesis, EPFL, Lausanne, Switzerland.

Condom, N., 2000. Analyse et modélisation des processus hydro-géochimiques de la salinisation

des sols. Application aux sols rizicoles de l'Office du Niger (Mali). PhD Thesis. Ecole

Nationale Supérieure d'Agronomie de Montpellier. 183 pp.

Dobermann, A. and Fairhurst, T.H., 2000. Rice: nutritional disorders and nutrient management.

International Rice Research Institute & Potash and Phosphate Institute, Singapore, 191 pp.

Eaton, F.M., 1950. Significance of carbonates in irrigation waters. *Soil Science*, 69:123-133

Favre, F., Tessier, D., Abdelmoula M., Génin, J. M. ,Gates, W. P. , Boivin, P. , 2002. Iron reduction

and changes in cation exchange capacity in intermittently waterlogged soil. *European Journal*

*of Soil Science*, 53: 175-183

- 1 thermodynamics of exchange adsorption. *Journal of Chemical Physics*, 21 :714-718.  
2
- 3 Haefele, S.; Wopereis, M. C. S.; Boivin, P. and N'Diaye, A. M. 1999. Effect of puddling on soil  
4 desalinization and rice seedling survival in the Senegal River Delta. *Soil and Tillage Research*,  
5  
6 51, 35-46  
7  
8  
9
- 10 Hammecker, C. and Maeght, J-L., 1999. Synthèse sur l'évolution des sols sous culture (Sénégal).  
11  
12 Lutte contre la dégradation des sols dans la vallée du fleuve Sénégal. Document PSI, technical  
13  
14 report.  
15  
16  
17
- 18 Hammecker, C., Maeght, J-L. And Wade M. 1999. Essai de drainage sur le périmètre de Donaye 8.  
19  
20 Lutte contre la dégradation des sols dans la vallée du fleuve Sénégal. Document PSI, technical  
21  
22 report.  
23  
24
- 25 Hammecker, C., Antonino, A.C.D., Maeght, J-L. and Boivin, P. 2003. Experimental study and  
26  
27 numerical modelling of the water transfers in an irrigated plot in Northern Senegal: evidences  
28  
29 of air entrapment. *European Journal of Soil Science*. 54: 491-503  
30  
31
- 32 Hardie, L. A. and Eugster, H. 1970. The evolution of closed basin brines. *Mineral. Soc. Amer. Spec.*  
33  
34 *Pap.*, 3, 273-290.  
35  
36
- 37 Helyar, K.R. and Porter, W.M., 1989. Soil acidification, its measurement and the processes  
38  
39 involved. In: A.D. Robson (Editor), *Soil acidity and plant growth*. Academic press, Sydney,  
40  
41 pp. 61-101.  
42  
43  
44
- 45 House, W.A, Howson, M. R., Pethybridge, A. D., 1988. Crystallisation kinetics of calcite in the  
46  
47 presence of magnesium ions. *J. Chem. Soc., Faraday Trans. I*, 84 : 2723 - 2734,  
48  
49
- 50 Madé, B., 1991. Modélisation thermodynamique et cinétique des réaction géochimiques dans les  
51  
52 interactions eau-roche. PhD thesis, Université Louis Pasteur, Strasbourg.  
53  
54  
55
- 56 Maignien R., 1961. Contribution à l'étude des sols à pâturage du cercle du Gorgol. *Rapp. Off.*  
57  
58 *Rech. sci. tech. O-mer, Dakar*, 125 p. multigr.  
59  
60  
61  
62  
63  
64  
65

- 1 for speciation, batch-reaction, one-dimensional transport, and inverse geochemical  
2  
3 calculations. Water-Resources investigations report 99-4256 USGS, Denver, Colorado  
4  
5
- 6 Pinton, R., Varanini, Z., Nannipieri, P. 2001. The Rhizosphere: Biochemistry and Organic  
7  
8 Substances at the Soil-plant Interface. CRC Press. 424 pp.  
9
- 10  
11 Valles, V., Pachepsky, Y.A., Ponizovsky, A.A. 1991. Invariant criteria for irrigation water quality  
12  
13 assessment in arid and semi arid regions. In: Genesis and Control of Fertility of Salt Affected  
14  
15 Soils. ISSS Sub-Commission on Salt Affected Soils Science, USSR, V.V. Dokuchaev Soil  
16  
17 Institute, Moscow, pp. 330-333.  
18  
19
- 20  
21  
22 van Asten, P.J.A. 2003. Soil Quality and Rice Productivity Problems in Sahelian Irrigation  
23  
24 Schemes, PhD thesis, Wageningen University, the Netherlands, 143 pp.  
25  
26
- 27  
28 van Asten, P.J.A., Barbiéro L., Wopereis M.C.S. and van der Zee, S.E.A.T.M. 2003. Actual and  
29  
30 potential salt-related soil degradation in an irrigated rice scheme in the Sahelian zone of  
31  
32 Mauritania. Agricultural Water Management. 60:13-32  
33  
34
- 35  
36 van Asten, P. J. A., van 't Zelfde, J. A., van der Zee, S. E. A. T. M. and Hammecker, C. 2004. The  
37  
38 effect of irrigated rice cropping on the alkalinity of two alkaline rice soils in the Sahel,  
39  
40 Geoderma, 119: 233-247  
41  
42
- 43  
44 van Beek, C.G.E. and Van Breemen, N. 1973. The alkalinity of alcali soils. J. Soil Sci. 24:129-136  
45
- 46  
47 van Ranst, E. and de Coninck, F. 2002. Evaluation of ferrollysis in soil formation. European Journal  
48  
49 of Soil Science. 53:513-519  
50
- 51  
52 Rengel, Z. 2003. Handbook of soil acidity , CRC Press, 496 pp  
53
- 54  
55 Ribolzi, O., Valles, V., Bariac, T., 1996. Comparison of hydrograph deconvolutions using residual  
56  
57 alkalinity, chloride and oxygen 18 as hydrochemical tracers. Water Res. Resour. 32 (4), 1051-  
58  
59  
60  
61  
62  
63  
64  
65

1059.

1 Riverside Salinity Laboratory Staff, 1954. Diagnosis and Improvement of Saline and Alkali Soils.

2  
3  
4 Agriculture Handbook n°60, United States Department of Agriculture

5  
6  
7 Schiess, S. 1995. Alcalinisation des sols irrigués dans la moyenne vallée du fleuve Sénégal :

8  
9 diagnostique (sic) par évaluation du stock de carbonates. M.Sc. Thesis, EPFL, Lausanne,  
10  
11 Switzerland.

12  
13  
14 Sharma, P., Pantuwan, G., Ingram, K., DeDatta, S. 1994. Rainfed lowland rice roots: soil and

15 hydrological effects. In: GJD, K. (Ed.), Rice Roots Nutrient and Water Use. IRRI, Manila, pp.  
16  
17 55 66.  
18  
19

20  
21  
22 Slaton, N.A., Beyrouy, C.A., Wells, B.R., Norman, R.J., Gbur, E.E. 1990. Root growth and

23 distribution of two short-season rice genotypes. Plant and Soils. 121: 269-278  
24

25  
26  
27 Stumm, W., Morgan, J.J., 1970. Aquatic Chemistry : An Introduction Emphasizing Chemical

28 Equilibria in Natural Waters. Wiley, New York.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure 1** : Location of the studied sites

**Figure 2** : Schematic representation water use dynamics in simulation : **a.** after irrigation, 2 liters of superficial ponding water + 1 liter soil solution in the root zone, **b.** : after evaporation and just before the next irrigation, 1 liter of soil solution in the root zone.

**Figure 3**: Schematic representation of the procedure adopted to simulate the effects of the leaching fraction (**L.F.**) and of the superficial flushing (**S.F.**). **a. -1**: the soil is saturated, **2** : at the end of the concentration period, part of the remaining solution is evacuated from the system (**LF**). **b. -1** : at the end of an inter-cropping period precipitated minerals and concentrated solutions are close to the soil surface, **2**: addition of a superficial layer of irrigation water, dissolving the minerals and equilibrating with soil solution and and exchangeable cations, **3**: part of the superficial layer is evacuated.

**Figure 4** : Evolution of irrigation water composition during concentration for  $p\text{CO}_2 = 10^{1.5}$  Pa and possibility for common minerals (calcite, quartz, sepiolite, kaolinite, gypsum) to precipitate.

**Figure 5** : Evolution of irrigation water composition during concentration in the Riverside diagram.

**Figure 6** : Evolution of soil solution and adsorption complex for Foum Gleita.  $\text{RAC} : \text{RA}_{\text{calcite}}$  , the straight horizontal solid line states for the precipitation of calcite.

**Figure 7** : Evolution of soil solution and adsorption complex for Niono.  $\text{RAC} : \text{RA}_{\text{calcite}}$  , the straight horizontal solid line states for the precipitation of calcite and the dotted line for the precipitation of sepiolite.

**Figure 8** : Evolution of soil solution and adsorption complex for Donaye,  $RAC : RA_{calcite}$ , the straight horizontal solid line states for the precipitation of calcite and the dotted line for the precipitation of gypsum.

**Figure 9** : Evolution of pH, EC of soil solution, ESP and  $RA_{calcite}$  of the soil solution for Fourn Gleïta considering presence (solid line) and absence (dotted line) of plant uptake.

**Figure 10** : Evolution of pH, EC of soil solution, ESP and  $RA_{calcite}$  of the soil solution for Niono considering presence (solid line) and absence (dotted line) of plant uptake.

**Figure 11** : Evolution of pH, EC of soil solution, ESP and  $RA_{calcite}$  of the soil solution for Donaye considering presence (solid line) and absence (dotted line) of plant uptake.

**Figure 12** : Evolution of pH, EC of soil solution, ESP and  $RA_{calcite}$  of the soil solution for Fourn Gleïta, considering different superficial flushing rates, vertical drainage and plant uptake.

**Figure 13** : Evolution of pH, EC of soil solution, ESP and  $RA_{calcite}$  of the soil solution for Niono, considering different superficial flushing rates, vertical drainage and plant uptake.

**Figure 14** : Evolution of pH, EC of soil solution, ESP and  $RA_{calcite}$  of the soil solution for Donaye, considering different superficial flushing rates and plant uptake.

**Figure 15** : Comparison of soil quality between irrigated rice cropped plots and uncultivated soil in Donaye. The diagrams were obtained with superficial (0-10cm) soil samples,  $n=130$  in irrigated plots and  $n=95$  in uncultivated soils.

## Revision notes for the manuscript GEODER 4246 :

Simulating the evolution of soil solutions in irrigated rice soils in the Sahel.

By

Claude Hammecker<sup>a</sup>, Piet van Asten<sup>b e</sup>, Serge Marlet<sup>c</sup>, Jean-Luc Maeght<sup>a</sup> and Roland Poss<sup>d</sup>.

The comments and corrections suggested by the reviewers have been taken into account in the new version of the manuscript. Considering the numerous items listed by the reviewers, they will not be commented individually.

Reviewer 1 :

- The different sites studied in this paper have been put into a regional context demonstrating their relevance according to specific geological, geomorphological, pedological, and textural conditions. Moreover their specific water management and hydrological conditions have been highlighted.
- The evolution of the Residual Alkalinity Calcite has been introduced into the new version of the manuscript, according to the reviewers recommendations. However this simulation study showed that  $AR_{\text{calcite}}$  is dependent on the mixing of solutions where  $pCO_2$  is different, and therefore partially fails to its conservative tracer property.
- The typos and missing references have been corrected.
- As during the simulated process, the volume of solution varies between two consecutive irrigations, the results presented for the soil solution compositions and the exchangeable cations are those computed for the situation where the water level in the plot is at the maximum, that is just before evaporation starts. Therefore the results presented in the manuscript are not exactly thoses for which the most soluble minerals like sepiolite or gypsum precipitate, though they do precipitate just before the next irrigation.



- The authors chose to keep the results and the discussion in the same section, in order to avoid repeating the same statements. However as recommended by the reviewer, the results have been discussed more thoroughly and documented.
- Extreme ESP values or Mg concentrations in soil solution and on the exchangeable complex, computed by the model in absence of plant or drainage have not been observed in field conditions, attesting that these processes have an important influence on the evolution of the soil and soil solution. When taken into account in the simulations they contribute in mitigating the influence of adverse irrigation water quality, displaying results closer to reality.
- The functioning of the model PHREEQC has been described more clearly in the revised version.
- The units for pCO<sub>2</sub> have been homogenized and the choice of pCO<sub>2</sub> in tropical soils has been documented.
- The reference to soil reduction and specially to ferrollysis had been precised according to the reviewers recommendations.
- In the conclusion we drew recommendations for a long lasting management of irrigation schemes according to the specificity of each site.

Reviewer 2 :

- As suggested, it has been stated more precisely that the results presented in the manuscript are simulations and not experimental results. However each it was possible the simulated results were validated by field data or references.
- The different precisions about the the definitions of alkalinity, ESP, SAR, EC, Ece, SARe required by the reviewer have been introduced in the revised version, when available.
- Supersaturation of the soil solution has not been taken into account in these study as it is usually governed by kinetic processes. Considering that the simulations are aimed to describe long term evolutions, we felt that the it would not have improved the simulation and the final results.
- The use of abbreviations has been improved and homogenized in the new version of the manuscript.
- Few data were available about the irrigation efficiencies in the different sites. Therefore this subject has not been discussed in the paper, though we understand it's importance.

## Simulating the evolution of soil solutions in irrigated rice soils in the Sahel.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Claude Hammecker<sup>a</sup>, Piet van Asten<sup>b e</sup>, Serge Marlet<sup>c</sup>, Jean-Luc Maeght<sup>a</sup> and Roland Poss<sup>d</sup>.

<sup>a</sup> Institut de Recherche pour le Développement UR 176, Land Development Department,  
Office of Science for Land Development, Pahonyothin Road, Chatuchak, Bangkok 10900,  
Thailand.

Fax : +66 2561 -2186

E-mail : [Claude.Hammecker@ird.fr](mailto:Claude.Hammecker@ird.fr)

<sup>b</sup> WARDA, The Africa Rice Center, BP 96, St.Louis, Senegal.

<sup>c</sup> CIRAD, UMR G-EAU, Avenue Agropolis, 34398 Montpellier Cedex 5, France

<sup>d</sup> Institut de Recherche pour le Développement UR 176, SupAgro, 2 place Viala, 34060  
Montpellier cedex 1, France

<sup>e</sup> present address : IITA- ESARC P.O.Box 7878 Kampala, Uganda

## Abstract

1  
2  
3  
4 In sahelian countries, agriculture often relies on irrigation especially for rice cropping which  
5  
6 has been introduced these last decades. As most of the water resources used for irrigation  
7  
8 (mainly rivers and lakes) have an alkaline composition, rice cropping is often held as  
9  
10 responsible for soil alkalinization. In order to evaluate the probability for this process to  
11  
12 occur, we studied 3 different sites in West Africa where numerical simulation of the  
13  
14 geochemical interactions between soil, irrigation water and groundwater were performed  
15  
16 with PHREEQC. It was possible to simulate the evaporation of the ponding water, the mixing  
17  
18 between groundwater and irrigation water, the lateral drainage (flushing) and the leaching,  
19  
20 and finally the management of the crop residues. By performing prospective simulations over  
21  
22 60 cropping cycles it has been shown that soil alkalinization and sodication are not inevitable  
23  
24 in these area especially if the cropping is properly managed. Moreover, when straw is  
25  
26 removed and the plot flushed before the new cropping season, alkaline soils can even be  
27  
28 improved with rice cropping.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Keywords : alkalinization, sodication, PHREEQC, plant uptake, water management, West  
40  
41 Africa  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

1  
2  
3  
4 In the Sahel, irrigated rice (*Oryza Sativa* L.) cropping has been introduced on a wide scale  
5  
6 during the last 20 years. The construction of large (> 100 ha) and costly irrigation schemes  
7  
8 were intended to increase food security and provide income for Sahelian farmers, whose  
9  
10 crops and livestock were devastated during the droughts in the 1970-1980's. Irrigated rice  
11  
12 cropping in the hot and dry Sahelian climate seems like a paradox. However, the major  
13  
14 Sahelian rivers and some large artificial reservoirs provide more than sufficient water to  
15  
16 support current irrigation activities throughout the year. Although most Sahelian irrigation  
17  
18 waters contain relatively little salt, they usually show alkaline compositions, where alkalinity  
19  
20 (Alk.) is the dominant aqueous species. It is defined as the algebraic sum of the strong base  
21  
22 cation and strong acid anion equivalents (Stumm and Morgan, 1970), and can be expressed  
23  
24 as :  $Alk. = (HCO_3^-) + (CO_3^{2-}) + (OH^-) - (H^+)$  , where ( ) denote the total concentration in  
25  
26  $mol_c l^{-1}$  . Moreover most of the Sahelian irrigation waters, show also a positive calcite residual  
27  
28 alkalinity ( $RA_{calcite}$ ) which corresponds to the alkalinity of the solution after precipitation of  
29  
30 calcite :  $RA_{calcite} = Alk. - (Ca^{2+})$ , (Valles et al., 1991; Bertrand et al., 1993).  $RA_{calcite}$  has also  
31  
32 been shown to be a conservative tracer of the soil solution ( Ribolzi et al., 1996; Barbiero et  
33  
34 al., 2001; Barbiero et al., 2004). Even the residual sodium carbonate, defined as  $RSC = Alk.$   
35  
36  $- (Ca^{2+}) - (Mg^{2+})$  ; which corresponds to the alkalinity of the solution after calcite and Mg-  
37  
38 carbonate or Mg-silicate precipitation, is positive for these solutions and is an indicator of  
39  
40 potential alkalinization and sodication (Eaton, 1950; Hardie and Eugster, 1970; Van Beek and  
41  
42 Van Breemen, 1973; Al Droubi et al.,1980). Continuous concentration of such waters in the  
43  
44 soil root zone may lead to the formation of alkaline and sodic soils, showing both high pH  
45  
46 values and high exchangeable sodium percentage :  $ESP = XNa^+ / CEC$  , where  $XNa^+$  is the  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

exchangeable Na<sup>+</sup> and CEC the total cation exchange capacity. Such soils are less productive

1 because of the pH-induced loss or low availability of several plant nutrients (e.g. Fe, P, Zn)  
2  
3 and poor physical properties of the sodic horizon (Abrol et al., 1988).  
4  
5

6 Whether or not large quantities of alkaline salts will accumulate in the soil root zone depends  
7  
8 largely on the water balance of the root zone. Evapotranspiration in the Sahelian climate can  
9  
10 be very large and regularly exceeds rates of 10 mm day<sup>-1</sup> in the dry season (November – May)  
11  
12 when maximum temperatures reach 40 to 45 °C (Van Asten et al., 2003, Hammecker et al.,  
13  
14 2003). In most irrigation schemes, percolation rates are very low (< 4 mm day<sup>-1</sup>) (Condom,  
15  
16 2000 ; Hammecker et al., 2003 ; Van Asten, unpublished), due to the low hydraulic  
17  
18 conductivity of the clayey soils and air entrapment (Hammecker et al., 2003). In the course of  
19  
20 the cropping season, percolation rates can further decrease if groundwater levels reach the  
21  
22 soil surface. The combination of large evapotranspiration and small percolation can lead to  
23  
24 the excessive build up of alkaline salts in the long run.  
25  
26  
27  
28  
29  
30

31 Recent studies have shown that the current soil alkalinity and sodicity problems in Sahelian  
32  
33 irrigation schemes are primarily caused by salts that originate from the parent material  
34  
35 (Barbiero and Van Vliet-Lanoë, 1998, Van Asten et al., 2004), or that accumulated as a result  
36  
37 of long-time irrigation of non-flooded crops in combination with shallow saline groundwater  
38  
39 levels. Hence, these salts were already in the soils before irrigated rice cropping was  
40  
41 introduced and there is little indication that the recent introduction of irrigated rice cropping  
42  
43 contributed to a significant increase in the salt content of these soils (van Asten, 2003).  
44  
45  
46  
47  
48

49 Moreover, most rice soils have the capacity to buffer an increase in soil solution alkalinity  
50  
51 through desorption of exchangeable Ca<sup>2+</sup> and the subsequent precipitation of calcite.  
52  
53

54 However, although this buffer capacity is large, it is not infinite. Consequently, it still remains  
55  
56 uncertain whether current irrigation practices will not lead to soil alkalization and  
57  
58  
59  
60  
61

sodication in the long run (i.e. some decades) in areas where salts added through irrigation can not sufficiently be leached or evacuated from the soil root zone. Given the high investment costs of irrigated agriculture and the high costs of reversing alkalization and sodication processes, it is important to know whether current or alternative crop and water management will lead to the degradation of one of the Sahel's most productive crop system.

In this study, we focus on three Sahelian irrigation situations, inside irrigation schemes, that differ distinctly in terms of parent material, soil water balance and irrigation water quality.

All sites have earlier been subject of studies on soil and water quality (Hammecker and Maeght, 1999; Boivin et al., 2002; Van Asten, 2003). Therefore, mean values for basic crop, water and soils parameters are already available. Taking into account some of the most important chemical and hydrological features, we tried to simulate how soils and soil solutions would evolve over time, under current and alternative land management practices at the three sites. The most critical parameter; i.e. the ratio between leached and irrigated solution volume, is the leaching fraction (LF) whether with vertical drainage or lateral drainage (flushing). In order to account for spatial variability within the irrigation schemes, we calculated soil solution evolution for different leaching fractions. This approach allows for the identification of the minimum leaching fraction that is required, in order to prevent soil degradation in the long run. Proton excretion (or alkalinity uptake) by the rice plant can also contribute significantly to the alkalinity balance of the soil root zone (Van Asten et al., 2004) and was also taken into account.

The objectives of this study are: (i) to give an overview of basic soil and water parameters for a range of Sahelian irrigation schemes, (ii) to simulate soil evolution under current and

alternative crop and water management practices, and (iii) to identify the minimum leaching fraction that is required to prevent soil degradation in the long term.

## 2. Material and methods

### 2.1. Study sites

This study is based on data collected in three different irrigation areas in West Africa located in the valley of River Senegal, the Gorgol noir and in the valley of river Niger, where irrigated rice cropping is the main agricultural activity.

These sites are representative of most of the conditions where rice-cropping cultivation is observed in the Sahel, and are representative of different geological, geomorphological and pedological situations with different soil textures. They are representative of (i) silty clay loam soils, developed on crystalline bedrock (Foum Gleita), (ii) alluvial levees with loamy sand soils (Niono) and (iii) alluvial basin with clayey vertisol (Donaye).

#### 2.1.1. Foum Gleita

The 'Mauritanides' geological region where Foum Gleita is located, is representative of the Pan-African orogenic belt which has been folded and lifted in the Appalachian era (Carité, 1989) and is consisting of metamorphic rocks, mainly greenschists, calcshists, micaschists and quartz, and secondary intrusions of dolerite, on which pedogenesis developed alkaline soils.

The Foum Gleita irrigation scheme is located in central southern Mauritania (N16°08'; W12°46'). The scheme was constructed between 1985 and 1989, with a surface area of 1950

ha. Irrigation water originates from a large artificial lake that formed behind a dam in the Gorgol Noir river. Irrigation and drainage are gravimetric but the drainage canals are often



not functional due to poor maintenance. Most soils in Foug Gleita have formed from the schist parent rock that can be found at shallow depths (< 2.5m) (Van Asten et al., 2003). Soil texture is mostly silty clay loam, with around 30% clay. Although no specific clay mineralogical studies have been done in the Foug Gleita schemes, other studies in the Gorgol basin have shown that a large proportion of the clays are illites and kaolinites (Maignien, 1961; Audry, 1961). The soils have a moderate to strong structure, varying from subangular blocky to prismatic in the more vertic soils. Biological activity in these soils is generally high. The porosity of the soils is generally medium to high, but is distinctly lower near soil surface. Although some soils have high ESP values, up to 65%, no evidence of soil structure degradation due to sodication has been observed in Foug Gleita. (Van Asten et al., 2003). In this region the cation exchange capacity (CEC) of soils ranges from 8 to 18 cmol<sub>c</sub>.kg<sup>-1</sup>, and the electrical conductivity of the 1:5 paste (EC<sub>1:5</sub>) varies from 0.12 to 1.32 dS.cm<sup>-1</sup>. Percolation rates are generally below 4 mm day<sup>-1</sup>, and decrease during the rainy season as groundwater approaches the surface. On a few soils, high percolation rates were measured (> 10 mm day<sup>-1</sup>) and farmers complained about difficulties in maintaining a submersion needed for rice. After the cropping season, groundwater quickly drops to below 1.2 m, indicating that deep percolation into the vertically oriented bedrock occurs, suggesting that horizontal groundwater flow along the slopes is minimal (van Asten et al., 2003). The absence of a shallow groundwater level during fallow periods suggests that capillary rise from the ground water can generally be neglected as a source of (alkaline) salts. The irrigation water is 2 to 4 times more saline than at the other study sites (EC = 0.17 dS.m<sup>-1</sup>), but the sodium adsorption ratio, defined as :  $SAR = Na^+ / (Ca^{2+} + Mg^{2+})^{1/2}$  where the concentration are in mmol.l<sup>-1</sup>, is still considered as very low (0.42). The residual sodium carbonate (RSC) is low and consequently suitable for irrigation according to the Riverside's diagram (see table

1) However, as at the other sites, RSC is positive and may therefore lead to alkalinization and sodication.

### 2.1.2. Niono

The site of Niono, is typical of the inner Delta of river Niger, developed on three different geological units: (i) Infracambrian substratum (-500 to -480 M years) composed of fractured schists and sandstones, (ii) Continental Terminal of Post Eocene to Pre-Quaternary age (-2.5 to -2 M years) formed by alternate sedimentary deposits of clays, sandy clays and sand , (iii) recent quaternary alluvial sediments deposited by the river Niger. In the studied area, the main morphological units are sandy levees, loamy inter levee depressions and clayey settling basins.

The study is based on data obtained in the huge irrigation scheme of the Office du Niger in Mali near Ségou (14°32 N 05°59 W). The climate is semi-arid (sudano-sahelian) with an average rainfall of 460 mm/year, distributed from June to October. The annual evaporation reaches 2.5m which generates soil solution concentration and consequently alkalization processes related to the alkaline composition of irrigation water (Condom, 2000) . The distribution of the soils in this area is inherited from ancient delta deposits constituted of clayey depressions delimited by sandy levees. These soils evolved under hydrologic conditions imposed by the water management during last half century, especially related to the uprise of the groundwater level.

The soils in this study are located on a little levee. The upper soil layers are composed of 11% clay and 78% sand, and classified as loamy sand. The identified clay minerals are mainly kaolinite, smectite and traces of illite. The structure of the soil is polyhedral. The soil displays an important biological porosity in the upper part of the profile (0-70cm) due to roots and to

meso-fauna activity. The CEC for this soil is 9 cmol<sub>c</sub>/kg but with large proportion of exchangeable sodium (ESP) of about 16% (table 2) ; the EC<sub>1:1</sub> for the soils in the area range from 0.18 to 1.09 dS.m<sup>-1</sup> . For the last 30 years the fields have been mainly cropped in rice with market gardening as a marginal activity (Condom, 2000).

In this large irrigation scheme plots are equipped with lateral drainage canals receiving mainly water from superficial flushing. After several decades of intensive irrigation, the water table level has risen approximatively by 50 m and is now close to the surface. Condom (2000) found that during the cropping season the average infiltration was about 80 mm (i.e. 9% of the leaching fraction), whereas during the inter cropping periods an equivalent amount of water from the water table supplied capillary rise towards the surface. Lateral losses towards drains and leaks represent 26 % of the total water inputs (Condom, 2000).

The Niger River that supplies the irrigation water by gravity has very low concentration (EC) and SAR values (Table 1), whereas the groundwater is 10 times more concentrated with a very high SAR value. According to the Riverside classification (Riverside salinity lab. staff, 1954) the quality of irrigation water does not constitute any harm to soil conservation but the composition of the groundwater typically represents a sodic hazard. Nevertheless, as illustrated by the positive RA<sub>calcite</sub> and RSC value (Table 1), these solutions can both become very alkaline and sodic when they concentrate, as the Ca<sup>2+</sup> and Mg<sup>2+</sup> might precipitate in minerals with low solubility like calcite and Mg-carbonates or Mg-silicates (e.g. sepiolite), whereas alkalinity and Na, keep concentrating in the solution (Vallès, 1991; Bethke, 2007).

### 2.1.3. Donaye

The third site, Donaye is representative of the irrigated soils of the river Senegal, representing

1 a total surface area of irrigation schemes of 240,000 ha (Barbiero et al., 2001). These soils  
2 developed on marine sediments deposited during the Nouakchottian transgression (4300 BP)  
3 which extended 380 km inland, near Bogue.

4  
5  
6 The irrigation scheme is located in northern Senegal (N16°40'; W15°), in the middle valley of  
7  
8 the Senegal River, near Podor. It is a 20-year-old irrigation scheme managed by the farmers,  
9  
10 without drainage network. The soil which has for long time been described as formed by  
11  
12 recent alluvial deposits is a vertisol (vertic Xerofluvent). However in a recent work Barbiero  
13  
14 et al. (2005) have suggested that the soil distribution in lower and middle valley of river  
15  
16 Senegal might result from the pedological transformation of a unique unripe mud, deposited  
17  
18 during the last quaternary marine transgressions. The soils of the Senegal valley can be  
19  
20 considered as a chronosequence, where the most recent soils can be observed downriver in  
21  
22 the delta, and the most developed ones upriver, in the middle valley. The pedological process  
23  
24 driven by the acidic conditions associated to the oxidation of inherited pyrite, led to the  
25  
26 dissolution of the parental mud. The protonation and acid hydrolysis of the aluminosilicates  
27  
28 in this material resulted in the formation of a sandy horizon, representing the quartz  
29  
30 skeleton. In the upper part of the soil, the interaction of the river water (through floods and  
31  
32 irrigation ) with the acidic, Al and Si rich soil solution contributed to the precipitation of clay  
33  
34 minerals, forming vertisols. In the Delta of the river Senegal this process is still active  
35  
36 (Barbiero et al. 2005) and the vertisol is still thin whereas in Donaye it represents a two-  
37  
38 meter-deep soil showing a vertic structure (slickensides), and constituted by 65 to 70% clay  
39  
40 minerals, mainly beidellite and kaolinite (Favre et al., 2002), which shows a high cation  
41  
42 exchange capacity of 22.7 cmol.kg<sup>-1</sup>(table 2). The soil overlays a 5 to 6 m thick sandy layer  
43  
44 corresponding to the afore mentioned quartz skeleton.

45  
46  
47 Rice cropping is performed by surge irrigation, and flooding is maintained during the  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

complete plant growth cycle (about 100 days). The presence of a shallow water table at 2m depth contributes to air entrapment and stops vertical infiltration (Hammecker et al. 2003). Consequently, vertical leaching is negligible, and soil solution in the upper part of the profile (above 40 cm) does not connect with the water table during the irrigation period. However, bad maintenance of bunds aimed to hold the irrigation in the paddy plots, generates important leaks contributing to non negligible loss of water, especially during the first irrigation. Irrigation water is pumped in a loop of river Senegal called Doué. This water with a very low concentration ( $EC = 0.07 \text{ dS.m}^{-1}$ ) and a low SAR value (Table 1), is suitable for irrigation according to Riverside's diagram. Nevertheless, like in the former situations, the  $RA_{\text{calcite}}$  and RSC of the irrigation water are positive, and the concentration of this solution may lead to sodication.

## 2.2. Numerical simulation.

### 2.2.1. The model and its main parameters.

The evolution of the soil solution and its interactions with the solid phase (precipitation-dissolution and exchange processes) were evaluated with the computer model PHREEQC v2. (Parkhurst and Appelo, 1999). This model, based on aqueous thermodynamics, computes equilibrium chemistry of aqueous solutions interacting with minerals, gases, solid solutions, exchangers, and sorption surfaces, providing a full speciation of an aqueous system. The different databases provided with the model give a very wide possibility for computing the activities of most of the aqueous species present in natural solutions, especially for the  $\text{CO}_3^-$ - $\text{HCO}_3^-$  solutions. This model is especially useful as, in addition to the calculation of the chemical equilibria, it allows for the simulation of mixing different solutions, for concentrating solution due to evaporation, and for adding or taking out aqueous species from

1 the soil solution. Using these options, it is possible to simulate solution changes due to  
2 flooded irrigation combined with concentration of the soil solution and evaporation. In surge  
3 flooding conditions, successive water inputs are usually performed to maintain a constant  
4 ponding level.  
5

6  
7  
8 Considering the duration of the rice cycle, the average evapotranspiration and the soil  
9 porosity, Hammecker et al. 2003 found that about 17 irrigations were necessary during one  
10 growing cycle. Considering that in the PHREEQC model all the calculations are performed  
11 for a given volume of solution, the Cation Exchange Capacity (CEC) and exchangeable  
12 cations were expressed per volume of liquid phase, depending on the soil porosity. The mass  
13 balance for water and solutes was calculated in the root zone, i.e. the first 20 centimeters of  
14 the soil profile (Slaton et al., 1990; Sharma et al., 1994; Boivin et al.2002). The irrigation  
15 water management during the cropping season, was simulated for one liter of soil solution in  
16 the root zone and the corresponding water volume for the superficial ponding water. As  
17 depicted in figure 2, during the first irrigation of the season 3 liters are introduced into the  
18 system : 1 liter for the soil solutions, and two liters for superficial ponding. During the  
19 cropping season, the superficial water level can almost disappear completely, but farmers  
20 usually irrigate before the complete drying-up of the plot.  
21

22 Different hydrological conditions, typical of field situations, were considered to simulate the  
23 geochemical evolution taking into account different leaching fractions, presence or absence of  
24 capillary rise from an aquifer with different composition, and the influence of plant uptake of  
25 nutriment. The actual leaching fraction (LF) being considered is the fraction of applied  
26 irrigation water that drains below the root zone :  
27

$$LF = \frac{D_d}{D_a} \quad [1]$$

28 Where  $D_d$  [L] is the amount of water drained below the root zone and  $D_a$  [L] the amount of  
29

irrigation water applied.

### 2.2.2. Chemical equilibria

Ionic activities in the soil solution were computed according to Debye-Hückel expression, based on the ionic strength of the solution. The soil solution was equilibrated with exchange sites on clay minerals and organic matter following Gaines and Thomas (1953) convention and mass conservation. Equilibrium with gas and minerals were considered for those initially present in soil, and those likely to crystallize or to affect the evolution of soil solution. The minerals usually found in arid soils like Mg-calcite, gypsum, quartz, kaolinite and sepiolite, were allowed to precipitate and dissolve, consequently to modify the composition of the solution. As quartz is initially present in the soil, this mineral has been chosen rather than amorphous silica, though the two options were tested and no differences in the major results of these simulations were found. When Al in solution analysis was not available it was computed considering equilibrium with kaolinite at the pH of the soil solution. The main solid and gaseous phases used in these simulations are reported in table 3, with their dissociation equation, and solubility products (K). Partial CO<sub>2</sub> pressure was considered to be equal to atmospheric value ( $\log p\text{CO}_2 = -3.5$  atm) whereas in the soil solution it was considered to be higher and constant, with  $\log p\text{CO}_2 = -1.8$  atm, like in most of the tropical soils (Brook et al. 1983). The crystallization of the different minerals was considered to take place when the solution reached equilibrium, regardless of any eventual super-saturation or kinetics phenomenon. However as the simulation is performed over a very long term, with different water contents, these processes will scarcely affect the final result.

### 2.2.3 The hydrological parameters

1 The adopted procedure for calculation was first, to equilibrate the irrigation solution with the  
2 soil solution, the mineral phases, and the partial CO<sub>2</sub> pressure in soil (10<sup>-1.8</sup> atm). The  
3 resulting solution was then concentrated to 5% of its original volume, as the farmers don't let  
4 the plot dry out completely. This value has been estimated from experimental observations  
5 (Hammecker, non published), though it has been verified that the choice of this value has  
6 little influence on the final result. During this procedure equilibria with the solid phase were  
7 computed considering the soils being initially formed by quartz, kaolinite, beidelite and by  
8 their respective exchangeable cations (table 2).  
9

10 Finally, depending on the leaching fraction (LF), the remaining water content after  
11 evaporation (Q<sub>0</sub>) was added to a new amount of irrigation water (Q<sub>1</sub>), in order to start a new  
12 simulation with the same amount of water (one liter), following this relation :

$$Q_0 = (1-LF) \times 0.05 \quad [2a]$$

$$Q_1 = 1 - Q_0 \quad [2b]$$

13 This procedure was performed 17 times for one cropping cycle, and 60 cropping cycles were  
14 simulated. In order to take into account the capillary rise in these geochemical processes, we  
15 estimated the quantity of solution possibly risen from the water table towards the root zone  
16 with water flows derived from hydraulic gradients measured during the dry period (Condom,  
17 2000). As capillary rise occurs only during the inter-cropping period, we treated both periods  
18 together, by mixing the two solutions (irrigation and water table) in their respective  
19 proportions (90% and 10%) in the root zone.  
20

21 In addition to vertical drainage, superficial lateral drainage (flushing) is performed  
22 intentionally or accidentally by the farmers in some situations. During this type of water  
23 management all salts and concentrated soils solutions present in the superficial layers of soil  
24 are dissolved and partially evacuated with the first water input at the beginning of the  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



cropping period. As the soil profile dried out during the inter-cropping season most of the soluble salts have accumulated in surface. Superficial lateral drainage experiment performed in Donaye showed that just with natural leakage from the bunds, an average of 15% of the total salinity (0-1m) could be removed (Hammecker et al. 1999). In Niono the flushing (lateral drainage) is performed intentionally, at the beginning of each new cropping season whereas in Donaye it occurs accidentally as, the margins of the plots leak after the inter cropping period when the soil shrinks and the margins crack. Consequently, we simulated an input of irrigation before each cropping period. This water was equilibrated with the soil solution, the minerals and the exchangeable cations, and was then partially evacuated from the system (Fig. 3). This numerical treatment, simulates the exportation of elements during a flush (lateral drainage), where flushing rates were defined as the proportion of water exported from the system, after the first water input had equilibrated with soil. Equilibrium during this procedure is conditioned by different physico-chemical processes like : (i) the competition between convective downward flow during infiltration and upward molecular diffusion towards the superficial water layer, (ii) the kinetics of minerals dissolution , (iii) the lateral losses in the plot. These processes were not considered individually but integrated into a bulk value of flushing rate. Different theoretical flushing rates were tested (5%, 10%, 30% and 50%) as no precise measurements for actual superficial lateral drainage were available in the three conditions. However they are in concordance with the flushing rate measured by Hammecker et al.1999 and with those determined in the experiment performed by Haefele et al. (1999) in the same area, who that showed superficial lateral drainage after submersed ploughing (puddling) at the beginning of the cropping season, could increase this salt exportation four folds.

### 2.2.3

#### 2.2.4. Plant uptake

The geochemical budget was considered with and without the incidence of the plant uptake, in order to quantify the contribution of the plant uptake to soil degradation or conservation processes. The main threat being alkalization due to carbonate concentration (pH increase), the focus has been set on the ability for rice to export alkalinity, and the main related cations (Ca and Mg) possibly affecting the carbonate balance and ESP. Data of Dobermann and Fairhurst (2000) on the average final composition of rice were used (Table 4). We considered an average yield of 5t/ha, which corresponds to 10t/ha of dry biomass(DB), a soil depth (z) of 0.2 m, an average bulk density ( $\rho_b$ ) of 1750 kg.m<sup>-3</sup> and a mineral density ( $\rho_s$ ) of 2650 kg.m<sup>-3</sup>.

The average ratio between the dry biomass (DB) and the volume of the soil solution ( $V_{soil}$ ) has been calculated to be 14.7 kg.m<sup>-3</sup> as follows :

$$\frac{DB}{V_{soil}} = \frac{DB \cdot 10^3}{z \cdot 10^4 \cdot \theta_s}$$


where  $\theta_s$  is the porosity. As the soil profile is saturated porosity is defined as:

$$\theta_s = 1 - \frac{\rho_b}{\rho_s}$$

Plants export alkalinity because they absorb more cations than anions. In order to maintain electrical neutrality both inside and outside of the roots they have to produce organic anions and excrete protons (Pinton et al.,2001 ; Rengel, 2003). The average plant alkalinity was calculated as the difference between the cations and anions absorbed by the plant (Table 4). It was considered that all plant alkalinity was removed from the field, as farmers export the entire plant for threshing.

Moreover, when a crop is considered in the geochemical system , it is also necessary to take

into account the N cycle. In paddy fields the plant absorb N as  $\text{NH}_4^+$  originating from the deamination of the organic matter and the hydrolysis of the urea used as N fertilizer:



In both cases for each mole of  $\text{NH}_4^+$  produced one mole of protons is removed from soil solution. Part of the ammonium produced is also adsorbed on the exchangeable complex or lost as ammonia. Therefore we only took into account the ammonium ions absorbed by the plant, by using the total N content of the plant (Table 4), irrespectively of the actual urea management in these sites. The net acidification due to plant uptake and the N cycle was calculated as:

$$\text{proton produced} = \text{plant alkalinity} - \text{N uptake.}$$

In order to simulate the exportation of alkalinity, hence the acidification of the soil solution due to proton excretion, potassium bicarbonate ( $\text{KHCO}_3$ ), magnesium and calcium carbonate ( $\text{MgCO}_3$ ,  $\text{CaCO}_3$ ) and potassium chloride (KCl) were numerically removed from the irrigation solution in proportions, calculated to meet the corresponding alkalinity uptake (Table 4). The total uptake for the different ions was divided by 17 (i.e. the number of irrigation events during a cropping cycle) and distributed evenly over the entire cropping season.

### 3. Results and discussion.

#### 3.1. Evolution of the irrigation solutions with increasing concentration.

Results displayed in Figure 4 show the evolution of the solution composition versus the concentration rate defined as :  $\text{concentration rate} = (V_i - V) / V_i$ , where  $V_i$  is the initial volume of solution and  $V$  the volume of solution reached after concentration.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

These results showed an important increase in pH with concentration, up to values around 9, and clearly superior to 8.3 the pK of calcite. Irrigation water of Foug Gleïta, rapidly reached saturation with calcite for its large initial Ca concentration. Consequently Ca concentration in solution decreased constantly whereas Alkalinity, Na and Mg concentrations increased, despite a slight depletion in Mg when sepiolite crystallization started. At the same time pH increased drastically despite sepiolite precipitation involving a release of protons into the solution (Table 3). Finally even for very high concentration rates, Mg increased noticeably in the solution because of its high initial content, as sepiolite crystallization is controlled by Si, which is less abundant.

During this simulation, the irrigation water of Niono rapidly reached saturation with quartz and while it is the only mineral precipitating, alkalinity and concentrations in all the cations increased regularly with concentration rate. The solution reached saturation with calcite for a high concentration rate (0.75) then Ca concentration decreased. For higher concentration rates, Mg concentration dropped when sepiolite crystallized whereas Na kept concentrating. A similar behavior was observed in Donaye, when water of river Senegal concentrated. In the three cases the pH of irrigation water increased and Na became the dominant cation in solution, enhancing the risk of sodication. Considering the SAR and the Electrical Conductivity of the irrigation water of the different situations, they can be considered as suitable for irrigation according to the Riverside diagram (1954). However this static view does not reflect the evolution of the solutions that may occur when concentrating. Numerical simulation of the concentration of these solutions (Figure 5), show that their SAR increased drastically with concentration, whereas the total salinity (EC) didn't increase as much, mainly because of the precipitation of calcite. Barbiero et al. (2004) showed some results for experimental concentration of irrigation water from River Niger, in the Republic of Niger,

1 which have similar initial composition as the irrigation water of Niono, where the SAR of the  
2 solutions also increased but with much higher EC values until reaching the C4S4 domain,  
3 instead of C2S4 as in the numerical simulation (Fig. 5). This discrepancy should mainly be  
4 explained by the fact that in the simulation the kinetics of precipitations have not been taken  
5 into account, and especially the retardation induced by the presence of Mg in the same  
6 proportions as Ca (House et al. 1988). However, in both cases the simulation shows that  
7 resulting solutions are extremely hazardous for soil conservation as they tend to promote  
8 sodication.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

### 21 3.2. Evolution of the soil solutions over 60 simulated cropping cycles.

22 The previous simulations described the evolution of the irrigation solution without taking  
23 into account the interactions with soils, especially with the exchange complex and the  
24 minerals. Moreover, in order to get an overall idea of the geochemical processes involved in  
25 field conditions, the vertical drainage and capillary rise was taken into account, following the  
26 simulation procedures described previously for the three situations.  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 The first set of simulation was aimed at quantifying the influence of the presence of a soil  
37 with a reactive adsorption complex and minerals, as well as the influence of the vertical  
38 leaching towards the water table. As the concentration of the soil varies constantly between  
39 two irrigations, the results displayed in the following figures corresponds to the composition  
40 of the soil solution and the exchange complex, immediately after the irrigation i.e. when the  
41 plots are full of water, in order to represent the actual solution infiltrating into the soil.  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 In the case of Foug Gleita (Fig. 6) where irrigation water tends to become very alkaline with  
52 concentration, an equivalent trend was observed in presence of soil and vertical drainage.  
53  
54  
55

56 However this trend appeared in lesser proportions, as pH reached only a value of 7.8 after 60  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 cropping cycles. The major ions in solution were  $\text{Na}^+$  and alkalinity. Both increased regularly  
2 with the number of cropping cycles whereas  $\text{Ca}^{2+}$  decreased as calcite precipitated. The  
3 complex reacted to compensate this trend, providing  $\text{Ca}^{2+}$  to the soil solution. Consequently,  
4 the proportion of exchangeable Ca decreased and exchangeable Mg became dominant. At the  
5 same time, exchangeable Na increased and consequently ESP rose to 15% after 10 cropping  
6 cycles and finally reached 30% after 60 cropping cycles. The evolution of the pH and  
7 alkalinity is still positive, but at a noticeably lower rate than for the concentration of the  
8 irrigation water alone. In fact, the large CEC of the soil buffered the effects of irrigation  
9 solution composition, especially as it imposed the geochemical composition of the soil  
10 solution, and supplied sufficient  $\text{Ca}^{2+}$  ions necessary to precipitate calcite and so prevented  
11 the accumulation of excessive alkalinity. According to the Residual Alkalinity (RA), which is  
12 still positive, the general trend of the system is to evolve continuously, but slowly, towards  
13 alkaline conditions.

14 In the case of Niono (Fig. 7), the simulation showed a continuous increase in pH during the  
15 first 35 cropping cycles followed by a stabilization at around  $\text{pH}=8.1$ . Alkalinity and  
16  $\text{Na}^+$  in soil solution and in the soil displayed a similar evolution, namely increasing until  
17 reaching saturation towards sepiolite.  $\text{RA}_{\text{calcite}}$  followed the same increasing trend as alkalinity,  
18 witnessing an alkaline evolution. Exchangeable sodium increased drastically during the first  
19 cropping cycles and reached a maximum of 53% of the exchangeable complex after 50  
20 cropping cycles, until sepiolite started to crystallize. Although extreme values for ESP have  
21 not actually been observed in this region, they are indicative of the type of degradation the  
22 soils might undergo in these conditions : high pH and ESP values are typical of alkalization  
23 and sodication processes. Although the irrigation solution is potentially less alkaline than in  
24 Fom Gleita, its use for irrigation is more hazardous, because the adsorption complex is very

vulnerable to soil solution composition due to low CEC ; the exchange between the adsorbed  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  in solution will therefore lead to a rapid increase in ESP.

Finally, in the case of Donaye (Fig. 8), the pH of the soil solution did not change much with the number of cropping cycles and mainly remained near neutrality. Sodium and magnesium were the major cations in the soil solution, and both concentrations increased with the number of cropping cycles. Alkalinity was the main anion, but both chloride and sulphate increased drastically with time, as there was no leaching towards the water table in this site.

The soil solution temporarily reached equilibrium with gypsum during the dry season when all the irrigation water was evaporated, and especially during the inter cropping periods, according field observations (Boivin et al., 2002; Barbiero et al., 2005).

In the first instance  $\text{RA}_{\text{calcite}}$  dropped slightly but increased again progressively when calcite and later gypsum started to precipitate. As the soil is very clayey with high CEC, the composition of the soil solution was determined by the buffering property of the soil.

Exchangeable calcium was progressively desorbed and precipitated, exchangeable magnesium increased and became the first exchangeable cation. At the same time ESP increased linearly up to 9% after 60 cycles. These extreme values for adsorbed magnesium were not observed yet, but still represent the values potentially reached for these conditions.

Sixty cropping cycles were not sufficient to lead to alkaline conditions. However these confined conditions promote a very slow alkalization and sodication process.

### 3.3. Importance of plant uptake

Results displayed in figure 9 to 11 showed clearly that the ionic plant uptake in the soil solution, decreased systematically the pH of soil solution in the three sites. Nevertheless, the

1 plant effect was moderate for the most alkaline conditions, namely Niono where pH still  
2 increased with time, up to 7.75 after sixty cropping cycles. In the case of Foun Gleita and  
3 Donaye pH decreased until reaching a final value of 7.2 as plant uptake counterbalanced the  
4 alkaline input of irrigation water. The total concentration of the soil solution, as indicated by  
5 EC, decreased by at least 30% , almost 50% for Foun Gleita and Niono. In Donaye, plant  
6 uptake played a major role for it is the only process exporting elements out of the system, as  
7 there was no vertical drainage. The same trend was observed for ESP in the cases of Foun  
8 Gleita and Donaye, where plant uptake contributed to a reduction by 50% of exchangeable  
9 sodium and consequently reduced the sodic risk. In Niono, unlike the former cases, plant  
10 uptake had a moderate effect on ESP. In fact, as  $K^+$ ,  $Ca^{2+}$  and  $Mg^{2+}$  were preferentially  
11 absorbed by rice to  $Na^+$ , its concentration increased relatively in soil solution and  
12 exchangeable position. In all cases,  $RA_{calcite}$  of the soil solution was significantly reduced, and  
13 even showing a serious decrease for Donaye until reaching  $0.0015 \text{ mol.l}^{-1}$  , though still  
14 remaining positive.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37 This simulation showed that in alkaline geochemical conditions, total exportation of rice  
38 (grains + straw) improved generally the composition of the soil solution and tended to  
39 reduce the risks of salinisation, alkalization and generally sodication except in the case of  
40 Niono where the low CEC was very sensitive to a relative Na increase in the soil solution. This  
41 situation is illustrated by Barbiero et al. 2001 who carried out an experiment in Niger (Lossa)  
42 where the cultivation of a local fodder grass (*Echinochloa stagnina*) and its exportation out  
43 of the field contributed to noticeable de-alkalization of a highly alkaline soil.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

### 57 3.4. Effects of superficial lateral drainage (flushing). 58 59 60 61 62 63 64 65



Results displayed in figures 12 to 14 showed distinctly that even for low rates, superficial flushing reduced significantly EC (i.e. the total salinity) of the soil solution, as well as ESP. The incidence of lateral drainage on pH and on RAC was less clear, especially for the case of Donaye as both parameters used to decrease without superficial lateral drainage and whereas they tended to increase slightly or stay constant when lateral drainage was simulated. However, in the case of Foug Gleita (Figure 12) the processes of sodication and alkalization were thoroughly reduced even for the smallest lateral drainage rate, as pH, EC, ESP and  $AR_{\text{calcite}}$  decrease with increasing flushing rate. For flushing rates greater than 30% the simulation shows that all these parameters reach constant low values, namely pH = 7.3, EC = 0.6 dS.m<sup>-1</sup>, ESP = 2%,  $AR_{\text{calcite}}$  = 0.03 mol.l<sup>-1</sup>, witnessing stable geochemical conditions, harmless for the quality of soil and solution.

In the case of Niono (figure 13), superficial flushing was as effective as in the previous example, for its incidence was clearly noticeable, even for low flushing rates. The simulation showed that a flushing rate of 30% was sufficient to control and maintain EC at a constant level, with only a slight increase in ESP. The incidence of lateral drainage on pH and  $AR_{\text{calcite}}$  was very important for the low rates (5% and 10%) during 20 first cropping cycles, as these parameters decreased slightly, until  $AR_{\text{calcite}}$  becoming negative for 5% flushing rate, and finally increased constantly. For the higher flushing rates (30 % and 50 %) the incidence on pH and  $AR_{\text{calcite}}$  was less important for the first simulated cropping cycles, but remained constant over the entire period. This special behavior is probably related to the difference in pCO<sub>2</sub> between the soil solution (10<sup>-1.8</sup> atm) and the supplementary irrigation water (10<sup>-3.5</sup> atm) added to compensate lateral drainage, and their relative proportions when mixed, promoting whether calcite precipitation or dilution of the soil solution. In any cases the superficial flushing reduced significantly the risk of alkalisation and sodication, though ESP

values still reached relatively high values (17%).

1 An equivalent behavior was noticed in the simulation for the case of Donaye (figure 14),  
2  
3 although no vertical drainage was considered; EC and ESP showed a significant decrease  
4  
5 even for low flushing rates and reached a low constant value all along the cropping cycles  
6  
7 when higher superficial lateral drainage rates were achieved (EC = 0.7 dS.m<sup>-1</sup> ; ESP = 1.5 %  
8  
9 for 30% flushing rate and EC = 0.6 dS.m<sup>-1</sup> ; ESP = 0.8 % for a flushing rate of 50 %).  
10  
11 Consequently the soil solution and soil quality increased. On the other hand, pH and AR<sub>calcite</sub>  
12  
13 showed a significant decrease when no lateral drainage was simulated whereas they showed  
14  
15 a similar behavior for the different flushing rates as described for the case of Niono.  
16  
17 However these variations of pH and AR<sub>calcite</sub> can be considered as negligible considering  
18  
19 alkalization because for the worst condition (flushing rate of 10 %), pH reached 7.33 and  
20  
21 AR<sub>calcite</sub> reached 0.004 mol<sub>c</sub>.l<sup>-1</sup> after 60 cropping cycles .  
22  
23

24 Moreover these results also show that AR<sub>calcite</sub> , which is considered as a conservative tracer  
25  
26 (Riboldi et al. 1996; Barbiero et al. 2001; Barbiero et al. 2004), varies along the simulation,  
27  
28 and probably also in field conditions, as solutions with different pCO<sub>2</sub> are mixed.  
29  
30


31 In Donaye, mineralogical and chemical experimental studies of the soil have been performed  
32  
33 to assess the modifications of the soil quality after 15 years of intensive irrigation. When  
34  
35 compared inside an irrigated plot and outside the irrigation scheme, the evolution of the soil  
36  
37 exchangeable complex, especially the ESP, has been found not significantly different  
38  
39 (Charolais and Weber, 1994). Equally, when the pH of the soil, the EC and the content of  
40  
41 accumulated carbonates in the topsoil have been measured on a 1/50 aqueous extract,  
42  
43 (Schiess, 1995; Boivin et al. 1998) only a slight increase in the soil between the two  
44  
45 situations was noticeable, though the difference isn't really statistically different between the  
46  
47 two groups (figure 15). These experimental results show that in 15 years of irrigation the soil  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 quality has not changed noticeably. In Donaye the rice straw is usually not exported from the  
2 field, but left to graze for cattle or burned. However, “natural” leakage throughout the bunds  
3 of the plot, due to a lack of maintenance is almost systematically observed. In fact, this  
4 disorder actually contributes to the superficial flushing, especially during the first irrigation  
5 when the soil shrinkage cracks are still open in bunds, and helps to preserve the soil quality  
6 despite the improper quality of irrigation water.  
7  
8  
9  
10  
11  
12

13 Although the quantitative effect of superficial flushing is difficult to model accurately because  
14 of the lack of information, this study demonstrates clearly the crucial role of superficial  
15 flushing for soil conservation, and attests the general conclusions of the numerical simulation  
16 results.  
17  
18  
19  
20  
21  
22  
23  
24  
25

### 26 3.5. Effect of soil reduction.

27 Although this process has not been modelled specifically in this work for the lack of Eh  
28 information in some cases, it is still possible to evaluate its effect on the proton balance. In  
29 the case of prolonged submersion conditions, like in paddy fields, the soil undergoes  
30 reduction as microbial activity consumes the dissolved oxygen, and finally reduces the iron  
31 oxi-hydroxides (the most commonly found). The oxidation of organic matter by reducing  
32 Fe(III) to Fe(II) produces alkalinity and consequently an increase in pH :  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43



45  
46 When the system oxidizes at the end of the submersion period an equivalent amount of acidity  
47 is produced and the general proton balance is null in confined conditions. However in case of  
48 a leaching fraction during reduced conditions, reoxidation leads generally to ferrolysis  
49 phenomenon which is a soil transformation involving a protonation of the complex, an  
50 attack of the clay crystalline structure, where  $\text{H}^+$  is penetrating the octahedral layers and  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 taking the place of  $Al^{3+}$  as described by Brinkman (1979) and revisited by van Ranst and De  
2 Coninck (2002). Consequently, soil reduction has either no effect on alkalinity or, on contrary,  
3 consumes alkalinity and reduces pH in the studied situations.  
4  
5  
6  
7  
8  
9

## 10 Conclusion

11  
12  
13  
14  
15  
16 In this study we used PHREEQC to predict the evolution of the soil and soil solution over  
17 several years, with different soil and water management. Although the use of Riversides'  
18 diagram,  $RA_{calcite}$  and the RSC index, can be informative about the potential risks, they cannot  
19 predict the evolution of the soil nor of the soil solution. Especially as they don't take into  
20 account the buffer capacity of soil, the water management involving dilutions and variations  
21 in  $pCO_2$ , and the effect of the plants uptake. We demonstrated that despite the high risk of  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

alkalinisation and sodication according to these indicators, in three west African situations of irrigation, soil degradation is not a fate. Considering the buffer effect of the soil (very high in Senegal), the possibilities of removing alkalinity from the plots by straw management, and finally the use of superficial flushing whether naturally (Foum Gleita), artificially with an effective drainage network (Niono) or accidentally with leaks in the bunds (Donaye), alkalination or sodication is not jeopardizing soil properties in well-conducted rice cropping system. These results using geochemical numerical modelling, attest that in Sahel, although the main rivers (river Senegal, Niger, Gorgol Noir) show an alkaline composition with positive  $RA_{calcite}$  and RSC values, their use for irrigation in paddy fields cannot be considered as directly responsible for soil alkalination. Barbiero et al. (1998), in the Republic of Niger, or van Asten et al. (2004) in Foum Gleita, found similar conclusions based on experimental

1 results. The presence of alkaline soils is more probably due to the alteration of an alkaline  
2 bed rock (in Foun Gleita) or to the presence of groundwater leaching through a deeper  
3 alkaline bed rock (in Niono). Unlike the usual assumptions, where rice cropping is held  
4 responsible for soil salinization or alkalinisation, this study show that in these cases, rice  
5 cropping can improve the soil quality when irrigation and residue management are  
6 performed properly. Namely a superficial lateral drainage performed once, at the beginning of  
7 the cropping season, although being sometimes seen as a waste of water by the local farmers,  
8 improves noticeably the sustainability of the system despite adverse composition of the  
9 irrigation water. More over, in the situation of alkalinisation risk, the exportation of the straw  
10 out of the plots contributes significantly to the control of the average alkalinity of soil,  
11 especially in the cases of Niono and Donaye.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table 1: Composition of irrigation water (I) and groundwater (GW) in the different locations

	Ca	Mg	Na	K	Alk.	Cl	SO <sub>4</sub>	RSC	pH	SAR	EC
	$\times 10^{-3} \text{ mol.l}^{-1}$										$\text{dS.m}^{-1}$
Foum <sup>I</sup>	0.78	0.38	0.32	0.12	1.4	0.11	0.09	0.24	7.54	0.42	0.17
Niono <sup>I</sup>	0.11	0.1	0.17	0.05	0.38	0.08	0	0.17	6.52	0.62	0.04
Niono <sup>GW</sup>	0.62	0.36	6.05	0.09	6.91	0.13	0.08	5.93	7.69	8.62	0.58
Donaye <sup>I</sup>	0.23	0.12	0.1	0.06	0.61	0.06	0.07	0.26	7.28	0.25	0.07

Table 2: Adsorbed cations of the different soils. ESP : exchangeable sodium percentage.

	XCa	XMg	XNa	XK	$\Sigma$ cat	ESP
	$\text{cmol}_c \text{kg}^{-1}$					%
Foum Gleita	11.8	4.8	0.3	0.6	17,5	1.71
Niono	5.5	2	1.5	0.1	9.1	16.48
Donaye	13	9	0.4	0.3	22.7	1.76

Table 3 : Minerals dissociation equations and solubility product values at 10<sup>5</sup> Pa and 298.15K

(1) : PHREEQC data base, (2) : estimated as a solid solution with data from PHREEQC data base, (3) : Kindis data base (Made, 1991) , (4) : fixed (Brooks et al. 1983)

Minerals	equations	log K (25 °C)
CO <sub>2</sub>		-1.8 <sup>(4)</sup>
Mg-Calcite	$\text{Ca}_{.98}\text{Mg}_{.02}\text{CO}_3 = .98 \text{Ca}^{2+} + .02\text{Mg}^{2+} + \text{CO}_3^{-2}$	-8.46608 <sup>(2)</sup>
Sepiolite	$\text{Mg}_2\text{Si}_3\text{O}_7.5\text{OH}:3\text{H}_2\text{O} + 4 \text{H}^+ + 0.5\text{H}_2\text{O} = 2 \text{Mg}^{2+} + 3 \text{H}_4\text{SiO}_4$	15.76 <sup>(1)</sup>
Kaolinite	$\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4 + 6 \text{H}^+ = \text{H}_2\text{O} + 2 \text{H}_4\text{SiO}_4 + 2 \text{Al}^{3+}$	7.44 <sup>(1)</sup>
Gypsum	$\text{CaSO}_4:2\text{H}_2\text{O} = \text{Ca}^{2+} + \text{SO}_4^{-2} + 2 \text{H}_2\text{O}$	-4.58 <sup>(1)</sup>
quartz	$\text{SiO}_2 + 2 \text{H}_2\text{O} = \text{H}_4\text{SiO}_4$	-3.98 <sup>(1)</sup>
beidellite	$\text{Si}_{3.8}\text{Al}_{1.7}\text{Fe}_{.3}\text{Mg}_{.2}\text{O}_{10}(\text{OH})_2\text{Ca}_{.2} + 6.8\text{H}^+ + 3.2 \text{H}_2\text{O} =$	-33.98 <sup>(3)</sup>
	$3.8\text{H}_4\text{SiO}_4 + 1.7\text{Al}^{3+} + .3\text{Fe}^{3+} + .2\text{Mg}^{2+} + .2\text{Ca}^{2+}$	



Table 4 : Average paddy uptake (grain + straw) and equivalent uptake in soil solution  
 (average yield of 5t/ha, and average bulk density of 1750kg.m<sup>-3</sup>)

	Dobermann & Fairhurst, 2000	equivalent ionic	
	%	species	mmol/l
N	0.65	NH <sub>4</sub> <sup>+</sup>	6.62
K	1.4	K <sup>+</sup>	5.1
Mg	0.2	Mg <sup>2+</sup>	2.35
Ca	0.3	Ca <sup>2+</sup>	2.13
Na	0.2	Na <sup>+</sup>	1.24
Σc			18.01
P	0.1	H <sub>2</sub> PO <sub>4</sub> <sup>-</sup>	0.46
Cl	0.5	Cl <sup>-</sup>	2.01
S	0.075	SO <sub>4</sub> <sup>2-</sup>	0.66
Σa			3.24
Σc - Σa		Alkalinity	14.3

## References

- 1  
2  
3  
4 Al Droubi, A., Fritz, B., Gac, J-Y. and Tardy, Y., 1980. Generalized residual alkalinity concept;  
5  
6 application to prediction of the chemical evolution of natural waters by evaporation.  
7  
8 Am. J. Sci., 280: 560-572  
9
- 10  
11 Arbol, I.P, Yadav, J.S.P and Massoud, FI., 1988. Salt-affected soils and their management.  
12  
13 Food and Agricultural Organization of the United Nations, Soils Bulletin 39, Rome 131  
14  
15 pp.  
16  
17
- 18  
19 Audry, P, 1961. Etude pédologique du cercle de Guidimaka (République Islamique de  
20  
21 Mauritanie). Etude du milieu naturel et étude pédologique. - Rapp. Off. Rech. sci. tech.  
22  
23 O-mer, Dakar, 173 p. multigr.,  
24  
25
- 26  
27 Barbiero, L. and Van-Vliet-Lanoë, B., 1998. The alkali soils of the middle Niger Valley; origins,  
28  
29 formation and present evolution. Geoderma, 84: 323-343.  
30  
31
- 32  
33 Barbiero, L.; Valles, V; Regard, A., Cheverry, C., 2001 Residual alkalinity as tracer to  
34  
35 estimate the changes induced by forage cultivation in a non-saline irrigated sodic soil  
36  
37 Agricultural Water Management. 50 : 229-241  
38
- 39  
40 Barbiero L., Ould Mohamedou A., Laperrousaz C., Furian S., Cunnac S., 2004. Polyphasic  
41  
42 origin of salinity in the Senegal delta and middle valley. Catena 58, 101-124.  
43
- 44  
45 Barbiero L., Ould Mohamedou A., Roger L., Furian S., Aventurier A., Rémy J.C., Marlet S.,  
46  
47 2005. The origin of vertisols and their relationship to Acid Sulfate Soils in the Senegal  
48  
49 Valley. Catena 59, 93-116.  
50
- 51  
52 Bertrand, R., Keita, B., Ndiaye, K.M., 1993. La dégradation des sols des périmètres irrigués  
53  
54 des grandes vallées sud-sahariennes: Cas de l'Office du Niger au Mali). Cahiers  
55  
56 Agricultures 2, 318 329.  
57  
58  
59  
60  
61  
62  
63  
64  
65

Bethke, C.M. 2007. *Geochemical and Biogeochemical Reaction Modeling*. Cambridge University Press. 564 pp.

Boivin, P, Favre, F and Maeght, J-L., 1998. Les sols de la moyenne vallée du fleuve Sénégal : caractéristiques et évolution sous irrigation. *Étude et Gestion des Sols*, 5: 235- 246

Boivin, P, Favre, F, Hammecker, C., Maeght, J-L., Delarivière, J., Poussin, J-C. and Wopereis, M.C.S., 2002. Processes driving soil solution chemistry in a flooded rice-cropped vertisol: analysis of long-time monitoring data. *Geoderma*, 110: 87-107

Bolan N.S. and Hedley M.J., 2003. Role of carbon, nitrogen, and sulfur cycles in soil acidification. In: Z. Rengel (Editor), *Handbook of soil acidity*. Marcel Dekker, New York, pp. 29-56.

Brinkman, R. 1979. *Ferrollysis, a soil-forming process in hydromorphic conditions*. Doctoral thesis. Agricultural Research Reports 887, Centre for Agricultural Publishing and Documentation, Wageningen.

Brooks, G.A., Folkoff, M.E., Box, E.O. 1983. A global model of soil carbon dioxide. *Earth Surface Processes and Land/onus* 8:79-88.

Carité, D. 1989. *Géologie en Mauritanie*. Edisud. 284 pp.

Charolais, M., Weber, V, 1994. Evolution des sols irrigués dans la moyenne vallée du fleuve Sénégal. M.Sc. Thesis, EPFL, Lausanne, Switzerland.

Condom, N., 2000. Analyse et modélisation des processus hydro-géochimiques de la salinisation des sols. Application aux sols rizicoles de l'Office du Niger (Mali). PhD Thesis. Ecole Nationale Supérieure d'Agronomie de Montpellier. 183 pp.

Dobermann, A. and Fairhurst, T.H., 2000. Rice: nutritional disorders and nutrient management. International Rice Research Institute & Potash and Phosphate Institute,

Singapore, 191 pp.

1 Eaton, F.M., 1950. Significance of carbonates in irrigation waters. *Soil Science*, 69:123-133

2  
3 Favre, F., Tessier, D., Abdelmoula M., Génin, J. M. ,Gates, W. P. , Boivin, P. , 2002. Iron  
4 reduction and changes in cation exchange capacity in intermittently waterlogged soil.  
5  
6 *European Journal of Soil Science*, 53: 175-183  
7  
8  
9

10  
11 Gaines, G.L., and Thomas, H.C., 1953. Adsorption studies on clay minerals. II. A formulation  
12 of the thermodynamics of exchange adsorption. *Journal of Chemical Physics*, 21 :714-  
13  
14 718.  
15  
16

17  
18 Haefele, S.; Wopereis, M. C. S.; Boivin, P and N'Diaye, A. M. 1999. Effect of puddling on soil  
19 desalinization and rice seedling survival in the Senegal River Delta. *Soil and Tillage*  
20  
21 *Research*, 51, 35-46  
22  
23  
24

25  
26 Hammecker, C. and Maeght, J-L., 1999. Synthèse sur l'évolution des sols sous culture  
27 (Sénégal). Lutte contre la dégradation des sols dans la vallée du fleuve Sénégal.  
28  
29 Document PSI, technical report.  
30  
31

32  
33 Hammecker, C., Maeght, J-L. And Wade M. 1999. Essai de drainage sur le périmètre de  
34 Donaye 8. Lutte contre la dégradation des sols dans la vallée du fleuve Sénégal.  
35  
36 Document PSI, technical report.  
37  
38  
39

40  
41 Hammecker, C., Antonino, A.C.D., Maeght, J-L. and Boivin, P. 2003. Experimental study and  
42 numerical modelling of the water transfers in an irrigated plot in Northern Senegal:  
43 evidences of air entrapment. *European Journal of Soil Science*. 54: 491-503  
44  
45  
46  
47

48  
49 Hardie, L. A. and Eugster, H. 1970. The evolution of closed basin brines. *Mineral. Soc. Amer.*  
50  
51 *Spec. Pap.*, 3, 273-290.  
52  
53

54 Helyar, K.R. and Porter, W.M., 1989. Soil acidification, its measurement and the processes  
55 involved. In: A.D. Robson (Editor), *Soil acidity and plant growth*. Academic press,  
56  
57  
58  
59

Sydney, pp. 61-101.

1  
2 House, W.A, Howson, M. R., Pethybridge, A. D., 1988. Crystallisation kinetics of calcite in the  
3  
4 presence of magnesium ions. *J. Chem. Soc., Faraday Trans. 1*, 84 : 2723 - 2734,  
5

6  
7 Madé, B., 1991. Modélisation thermodynamique et cinétique des réaction géochimiques dans  
8  
9 les interactions eau-roche. PhD thesis, Université Louis Pasteur, Strasbourg.  
10

11  
12 Maignien R., 1961. Contribution à l'étude des sols à pâturage du cercle du Gorgol. *Rapp. Off.*  
13  
14 *Rech. sci. tech. O-mer, Dakar*, 125 p. multigr.  
15

16  
17  
18 Parkhurst, D.L., Appelo C.A.J., 1999. User's guide to PHREEQC (version2)-a computer  
19  
20 program for speciation, batch-reaction, one-dimensional transport, and inverse  
21  
22 geochemical calculations. Water-Resources investigations report 99-4256 USGS, Denver,  
23  
24 Colorado  
25  
26

27  
28  
29 Pinton, R., Varanini, Z., Nannipieri, P. 2001. *The Rhizosphere: Biochemistry and Organic*  
30  
31 *Substances at the Soil-plant Interface*. CRC Press. 424 pp.  
32

33  
34  
35 Valles, V., Pachepsky, Y.A., Ponizovsky, A.A. 1991. Invariant criteria for irrigation water quality  
36  
37 assessment in arid and semi arid regions. In: *Genesis and Control of Fertility of Salt*  
38  
39 *Affected Soils*. ISSS Sub-Commission on Salt Affected Soils Science, USSR, V.V  
40  
41 *Dokuchaev Soil Institute, Moscow*, pp. 330-333.  
42  
43  
44

45  
46 van Asten, P.J.A. 2003. *Soil Quality and Rice Productivity Problems in Sahelian Irrigation*  
47  
48 *Schemes*, PhD thesis, Wageningen University, the Netherlands, 143 pp.  
49

50  
51 van Asten, P.J.A., Barbiéro L., Wopereis M.C.S. and van der Zee, S.E.A.T.M. 2003. Actual and  
52  
53 potential salt-related soil degradation in an irrigated rice scheme in the Sahelian zone of  
54  
55 Mauritania. *Agricultural Water Management*. 60:13-32  
56  
57  
58  
59  
60

van Asten, P. J. A., van 't Zelfde, J. A., van der Zee, S. E. A. T. M. and Hammecker, C. 2004.

1 The effect of irrigated rice cropping on the alkalinity of two alkaline rice soils in the  
2  
3 Sahel, Geoderma, 119: 233-247  
4  
5

6 van Beek, C.G.E. and Van Breemen, N. 1973. The alkalinity of alcali soils. J. Soil Sci. 24:129-  
7  
8  
9 136  
10

11 van Ranst, E. and de Coninck, F. 2002. Evaluation of ferrollysis in soil formation. European  
12  
13 Journal of Soil Science. 53:513-519  
14  
15

16 Rengel, Z. 2003. Handbook of soil acidity , CRC Press, 496 pp  
17  
18

19 Ribolzi, O., Valles, V., Bariac, T., 1996. Comparison of hydrograph deconvolutions using  
20  
21 residual alkalinity, chloride and oxygen 18 as hydrochemical tracers. Water Res. Resour.  
22  
23 32 (4), 1051-1059.  
24  
25  
26

27 Riverside Salinity Laboratory Staff, 1954. Diagnosis and Improvement of Saline and Alkali  
28  
29 Soils. Agriculture Handbook n°60, United States Department of Agriculture  
30  
31  
32

33 Schiess, S. 1995. Alcalinisation des sols irrigués dans la moyenne vallée du fleuve Sénégal :  
34  
35 diagnostique (sic) par évaluation du stock de carbonates. M.Sc. Thesis, EPFL, Lausanne,  
36  
37  
38 Switzerland.  
39  
40  
41

42 Sharma, P., Pantuwan, G., Ingram, K., DeDatta, S. 1994. Rainfed lowland rice roots: soil and  
43  
44 hydrological effects. In: GJD, K. (Ed.), Rice Roots Nutrient and Water Use. IRRI, Manila,  
45  
46 pp. 55 66.  
47  
48  
49

50 Slaton, N.A., Beyrouy, C.A., Wells, B.R., Norman, R.J., Gbur, E.E. 1990. Root growth and  
51  
52 distribution of two short-season rice genotypes. Plant and Soils. 121: 269-278  
53  
54  
55

56 Stumm, W., Morgan, J.J., 1970. Aquatic Chemistry : An Introduction Emphasizing Chemical  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 **Figure 1** : Location of the studied sites

4 **Figure 2** : Schematic representation water use dynamics in simulation : **a.** after irrigation, 2  
5 liters of superficial ponding water + 1 liter soil solution in the root zone, **b.** : after  
6 evaporation and just before the next irrigation, 1 liter of soil solution in the root zone.  
7  
8  
9

10  
11  
12 **Figure 3**: Schematic representation of the procedure adopted to simulate the effects of the  
13 leaching fraction (**L.F.**) and of the superficial flushing (**S.F.**). **a.** -1: the soil is saturated, 2 : at  
14 the end of the concentration period, part of the remaining solution is evacuated from the  
15 system (LF). **b.** -1 : at the end of an inter-cropping period precipitated minerals and  
16 concentrated solutions are close to the soil surface, 2: addition of a superficial layer of  
17 irrigation water, dissolving the minerals and equilibrating with soil solution and and  
18 exchangeable cations, 3: part of the superficial layer is evacuated.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 **Figure 4** : Evolution of irrigation water composition during concentration for  $p\text{CO}_2 = 10^{1.5} \text{ Pa}$   
34 and possibility for common minerals (calcite, quartz, sepiolite, kaolinite, gypsum) to  
35 precipitate.  
36  
37  
38  
39  
40  
41  
42

43 **Figure 5** : Evolution of irrigation water composition during concentration in the Riverside  
44 diagram.  
45  
46  
47  
48  
49  
50

51 **Figure 6** : Evolution of soil solution and adsorption complex for Foug Gleita.  $\text{RAC} : \text{RA}_{\text{calcite}}$ ,  
52 the straight horizontal solid line states for the precipitation of calcite.  
53  
54  
55  
56  
57

58 **Figure 7** : Evolution of soil solution and adsorption complex for Niono.  $\text{RAC} : \text{RA}_{\text{calcite}}$ , the  
59  
60  
61  
62  
63  
64  
65



1 straight horizontal solid line states for the precipitation of calcite and the dotted line for the  
2 precipitation of sepiolite.  
3  
4

5 **Figure 8** : Evolution of soil solution and adsorption complex for Donaye, RAC :  $RA_{\text{calcite}}$  , the  
6 straight horizontal solid line states for the precipitation of calcite and the dotted line for the  
7 precipitation of gypsum.  
8  
9

10  
11  
12  
13  
14  
15  
16 **Figure 9** : Evolution of pH, EC of soil solution, ESP and  $RA_{\text{calcite}}$  of the soil solution for Fourn  
17 Gleïta considering presence (solid line) and absence (dotted line) of plant uptake.  
18  
19

20  
21  
22  
23 **Figure 10** : Evolution of pH, EC of soil solution, ESP and  $RA_{\text{calcite}}$  of the soil solution for Niono  
24 considering presence (solid line) and absence (dotted line) of plant uptake.  
25  
26

27  
28 **Figure 11** : Evolution of pH, EC of soil solution, ESP and  $RA_{\text{calcite}}$  of the soil solution for  
29 Donaye considering presence (solid line) and absence (dotted line) of plant uptake.  
30  
31

32  
33  
34  
35  
36 **Figure 12** : Evolution of pH, EC of soil solution, ESP and  $RA_{\text{calcite}}$  of the soil solution for Fourn  
37 Gleïta, considering different superficial flushing rates, vertical drainage and plant uptake.  
38  
39

40  
41 **Figure 13** : Evolution of pH, EC of soil solution, ESP and  $RA_{\text{calcite}}$  of the soil solution for  
42 Niono, considering different superficial flushing rates, vertical drainage and plant uptake.  
43  
44

45  
46  
47  
48  
49 **Figure 14** : Evolution of pH, EC of soil solution, ESP and  $RA_{\text{calcite}}$  of the soil solution for  
50 Donaye, considering different superficial flushing rates and plant uptake.  
51  
52

53  
54  
55  
56 **Figure 15** : Comparison of soil quality between irrigated rice cropped plots and uncultivated  
57  
58  
59  
60

soil in Donaye. The diagrams were obtained with superficial (0-10cm) soil samples, n=130

in irrigated plots and n=95 in uncultivated soils.

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

Figure1

[Click here to download high resolution image](#)




Figure2

[Click here to download high resolution image](#)





Figure 3  
[Click here to download high resolution image](#)




Figure 4  
[Click here to download high resolution image](#)




Figure 5  
[Click here to download high resolution image](#)




Figure 6  
[Click here to download high resolution image](#)






Figure 7  
[Click here to download high resolution image](#)




Figure 8

[Click here to download high resolution image](#)




Figure 9  
[Click here to download high resolution image](#)




Figure 10  
[Click here to download high resolution image](#)




Figure 11

[Click here to download high resolution image](#)




Figure 12  
[Click here to download high resolution image](#)




Figure 13

[Click here to download high resolution image](#)




Figure 14

[Click here to download high resolution image](#)






Figure 15  
[Click here to download high resolution image](#)

