



Fishing activity, health characteristics and mercury exposure of Amerindian women living alongside the Beni River (Amazonian Bolivia).

Eric Bénéfice, Selma Luna Monrroy, Ronald Lopez-Rodriguez

► To cite this version:

Eric Bénéfice, Selma Luna Monrroy, Ronald Lopez-Rodriguez. Fishing activity, health characteristics and mercury exposure of Amerindian women living alongside the Beni River (Amazonian Bolivia).. International Journal of Hygiene and Environmental Health, 2010, 213 (6), pp.458-64. 10.1016/j.ijheh.2010.08.010 . ird-00547706

HAL Id: ird-00547706

<https://ird.hal.science/ird-00547706>

Submitted on 17 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Table 1 Fishing activity and fish consumption of the women in relation to hair mercury concentrations (H-Hg, µg/g)

| Variable | Number | Median H-Hg | 95% CI | H value ¹ p |
|-------------------------------|--------|----------------|---------|---------------------------|
| Livelihoods | | | | |
| Wage earners and craftsmen | 18 | 2.3 | 1.7-3.9 | 38.1 |
| Farming and logging | 88 | 3.3 | 2.7-3.9 | 0.000 |
| Fishing as principal activity | 57 | 8.2 | 6.4-9.1 | |
| Fish consumption | | | | |
| Once a day | 32 | 8.0 | 5.6-9.1 | 15.7 |
| At least once a week | 99 | 3.7 | 2.9-5.0 | 0.0003 |
| Less than once a week | 32 | 3.5 | 2.3-5.1 | |

¹ Z: Mann-Whitney U or Wilcoxon rank-sum test for difference in medians (two means comparisons).

Table 2 Characteristics of women examined during the study

| Variable | Mean/number | SD | median | Percent abnormal (95% CI) |
|---|--------------------------------|------|--------|--|
| Ethnic group | Esse Ejjas: 25 Tacanas: 148 | | | |
| Age (years) | 34.9 | 13.2 | 33.0 | |
| Height (cm) | 149.2 | 5.3 | 148.6 | % < 145 cm 21.3 (15.5 ~ 28.2) |
| BMI (kg/m^2) | 24.1 | 3.3 | 24.0 | % > 24.5 42.7 (35.3 ~ 50.5) |
| Hemoglobin ($\text{g}/100 \text{ ml}$) | 12.2 | 1.7 | 12.4 | % < 11 $\text{g}/100 \text{ ml}$ 22.3 (16.3 ~ 29.3) |
| H-Hg ($\mu\text{g}/\text{g}$) | 5.5 | 4.1 | 4.4 | % > 10 $\mu\text{g}/\text{g}$ 14.7 (9.6 ~ 21.1) |
| Systolic blood pressure (mm Hg) | 106.9 | 14.8 | 108 | % > 130 mmHg 4.1 (1.6 ~ 8.2) |
| Diastolic blood pressure (mm Hg) | 66.2 | 9.3 | 65 | % > 90 mmHg 0.0 (0.0 ~ 2.1) |

Table 3 Responses to neurological examination of contaminated and non-contaminated women

| Item | Response | H-Hg $\leq 5\mu\text{g/g}$ | H-Hg $> 5\mu\text{g/g}$ | χ^2 | p ^a |
|--------------------|-------------|----------------------------|-------------------------|----------|----------------|
| Visual field | normal | 78 | 57 | 0.31 | ns |
| | reduced | 7 | 7 | | |
| Paresthesia | no | 82 | 53 | 7.0 | 0.007 |
| | yes | 5 | 13 | | |
| Static balance | normal | 86 | 59 | 5.5 | 0.01 |
| | instability | 1 | 6 | | |
| Dynamic balance | Normal | 61 | 45 | 0.01 | ns |
| | Instability | 26 | 20 | | |
| Motor coordination | normal | 78 | 53 | 1.5 | ns |
| | weak | 9 | 11 | | |

^a statistical power (ns: not significant)

^b power value of the Fisher exact test

Table 4 Reproductive health parameters of the women in relation to hair mercury concentrations (H-Hg, µg/g)

| Variable | Number | Median H-Hg | 95% CI | Z values or H value ¹ p |
|---------------------------|--------|----------------|---------|---------------------------------------|
| Post-menopausal | 132 | 4.2 | 2.5-6.1 | Z = 0.7 |
| Pre-menopausal | 132 | 4.5 | 3.5-5.5 | ns |
| Pregnant ² | 17 | 3.2 | 1.3-3.8 | Z=2.4 |
| Not pregnant | 115 | 5.2 | 3.9-6.2 | 0.001 |
| Lactating | 59 | 5.5 | 4.5-6.4 | Z=2.2 |
| Not lactating | 72 | 3.5 | 2.8-5.2 | 0.02 |
| Child deaths | | | | |
| 0 child deaths | 79 | 3.5 | 2.6-4.4 | H= 9.0 |
| 1 child death | 33 | 5.2 | 3.5-7.8 | 0.01 |
| More than one child death | 50 | 5.6 | 3.7-7.9 | |
| Spontaneous abortion | | | | |
| 0 abortions | 114 | 4.0 | 3.3-5.6 | H=4.2 |
| 1 abortion | 30 | 4.4 | 2.7-6.1 | ns |
| More than one abortion | 19 | 8.0 | 2.9-9.6 | |

¹ Z: Mann-Whitney U or Wilcoxon rank-sum test for difference in medians (two means comparisons).

H: Kruskall Wallis one way analysis of variance on ranks (several means comparisons)

Ns= not significant

² Comparisons made only in pre-menopausal women

Table 5. Result of logistic regression between risk of mercury contamination, socio-cultural and physical characteristics of women

| Parameter | Wald Z-Value (Beta=0) | Wald Prob Level | Odds Ratio Exp(B) | Lower 95% CI | Upper 95% CI | Regression Coefficient (B or Beta) |
|------------------------------|--------------------------|-----------------|----------------------|--------------|--------------|---------------------------------------|
| Intercept | -1.55 | 0.12 | 0.03 | 0.00 | 2.64 | -3.63 |
| Age | -2.41 | 0.02 | 0.95 | 0.92 | 0.99 | -0.05 |
| BMI | 2.32 | 0.02 | 1.18 | 1.03 | 1.35 | 0.16 |
| Static balance abnormalities | 2.30 | 0.02 | 22.46 | 1.59 | 316.80 | 3.11 |
| Esse Ejjas ethnic group | 1.98 | 0.05 | 6.31 | 1.01 | 39.23 | 1.84 |
| Paresthesia | 2.12 | 0.03 | 4.48 | 1.12 | 17.94 | 1.50 |
| Fishing activity | -3.19 | 0.00 | 0.18 | 0.06 | 0.51 | -1.73 |
| Child death | 2.81 | 0.01 | 4.17 | 1.54 | 11.31 | 1.43 |
| Daily fish consumption | 2.73 | 0.01 | 4.80 | 1.56 | 14.79 | 1.57 |
| Blood hemoglobin | 0.49 | 0.63 | 1.07 | 0.83 | 1.37 | 0.06 |