



**HAL**  
open science

# The role of predation by harp seals (*Pagophilus groenlandicus*) in the collapse and non-recovery of northern Gulf of St. Lawrence cod (*Gadus morhua*)

Emmanuel Chassot, Daniel E. Duplisea, Mike O. Hammill, Amanda Caskenette, Nicolas Bousquet, Yvan Lambert, Garry H. Stenson

## ► To cite this version:

Emmanuel Chassot, Daniel E. Duplisea, Mike O. Hammill, Amanda Caskenette, Nicolas Bousquet, et al.. The role of predation by harp seals (*Pagophilus groenlandicus*) in the collapse and non-recovery of northern Gulf of St. Lawrence cod (*Gadus morhua*). *Marine Ecology Progress Series*, 2009, 379, pp.279-297. 10.3354/meps07897 . ird-00549074

**HAL Id: ird-00549074**

**<https://ird.hal.science/ird-00549074v1>**

Submitted on 21 Dec 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The role of predation by harp seals (*Pagophilus groenlandicus*) in the collapse and non-recovery of northern Gulf of St. Lawrence cod (*Gadus morhua*)

Emmanuel Chassot<sup>1,2,\*</sup>, Daniel Duplisea<sup>1</sup>, Mike Hammill<sup>1</sup>, Amanda Caskenette<sup>1</sup>, Nicolas Bousquet<sup>3</sup>, Yvan Lambert<sup>1</sup>, and Garry Stenson<sup>4</sup>

<sup>1</sup>*Fisheries and Oceans Canada, Institut Maurice-Lamontagne, 850 route de la Mer, Mont-Joli, QC, G5H 3Z4, CANADA*

<sup>2</sup>*IRD, CRH, Avenue Jean Monnet, BP 171, 34 203 Sète Cedex, FRANCE (present address)*

<sup>3</sup>*Département de Mathématiques et Statistiques, Université Laval, Pavillon Alexandre-Vachon, QC, G1K 7P4, CANADA*

<sup>4</sup>*Fisheries and Oceans Canada, Northwest Atlantic Fisheries Centre, PO Box 5667, St. John's, NF, A1C 5X1, CANADA*

Suggested running title: Harp seal predation on cod

Correspondence: Emmanuel Chassot

Telephone: +33(0)4 99 57 32 24  
Fax: +33(0)4 99 57 32 95  
Email: Emmanuel.Chassot@ird.fr

1 ABSTRACT: A statistical catch-at-age model was developed to assess the effects of predation  
2 by the northwest Atlantic harp seal population on northern Gulf of St. Lawrence cod by  
3 estimating the relative importance of different sources of mortality that affected the stock  
4 during a period of collapse and non-recovery. Cod recruitment at age 1 is modeled via a non-  
5 linear stock-recruitment relationship based on total egg production and accounts for changes in  
6 female length-at-maturity and cod condition. Natural mortality other than seal predation also  
7 depends on cod condition used as an integrative index of changes in environmental conditions.  
8 The linkage between seals and cod is modeled through a multi-age functional response that was  
9 derived from the reconstruction of the seal diet using morphometric relationships and stomach  
10 contents of more than 200 seals collected between 1998 and 2001. The model was fitted  
11 following a maximum likelihood estimation approach to a scientific survey abundance index  
12 (1984-2006). Model results show that the collapse of the northern Gulf of St. Lawrence cod  
13 stock was mainly due to the combination of high fishing mortality rates and poor environmental  
14 conditions in the early to mid-1990s contributing to the current state of recruitment overfishing.  
15 The increase in harp seal abundance during 1984-2006 was reflected by an increase in predation  
16 mortality for the young cod age-groups targeted by seals. Although current levels of predation  
17 mortality affect cod spawning biomass, the lack of recovery of the NGSL cod stock seems  
18 mainly due to the situation of very poor recruitment.

19 *Keywords : cod, harp seal, functional response, model, predation, recovery*

## INTRODUCTION

21 During the early 1990s, there was an almost simultaneous collapse among most of the Atlantic  
22 cod (*Gadus morhua* L.) fisheries in eastern Canada, leading to severe restrictions or moratoria  
23 on commercial fishing (Myers et al. 1997, Rice et al. 2003). The northern Gulf of St. Lawrence  
24 (NGSL) cod stock (NAFO divisions 3Pn4RS; Fig. 1) was historically the second largest cod  
25 population in the Western Atlantic with a spawning stock biomass (SSB) of 380,000 t in the  
26 early 1980s, supporting a fishery of more than 100,000 t in 1983 (MPO 2007). By the late  
27 1980s, the population and fishery declined to such an extent that a moratorium was imposed  
28 in 1994, leaving a SSB of only about 10,000 t. Subsequently, the population increased quite  
29 rapidly and in 1997, a small fishery reopened, although the stock has remained at about 50%  
30 of its minimum conservation biomass limit ( $B_{lim}$ ) of about 80-90,000 t.

31 There has been considerable debate concerning the causes for the decline in cod abundance  
32 and subsequent lack of recovery (Hutchings 1996, Shelton & Healey 1999, Shelton et al. 2006).  
33 Overfishing has been identified as the main factor that contributed to the collapse due to several  
34 interrelated common factors including underestimated discarding and misreporting (Myers et  
35 al. 1996, Myers et al. 1997, Savenkoff et al. 2004). For the NGSL cod stock, it has been  
36 suggested that changes in environmental conditions in the early 1990s also played a role in the  
37 fishery collapse through a decline in condition, which consequently resulted in reduced growth  
38 and recruitment (Lambert & Dutil 1997a, Dutil et al. 1999, Dutil & Lambert 2000). Poor  
39 condition is also suspected to have led to an increase in natural mortality through starvation  
40 and enhanced vulnerability to predation (Dutil et al. 1999). The current poor productivity  
41 of the stock associated with ongoing fishing activity has been proposed as a major factor  
42 explaining the lack of recovery for NGSL cod (Dutil et al. 2003, Shelton et al. 2006).

43 During the last 3 decades, there has been a marked increase in abundance of many pinniped  
44 populations in the northwest Atlantic (Hammill et al. 1998, Hammill & Stenson 2005). Higher  
45 natural mortality resulting from increased predation has also been put forward as a plausible  
46 hypothesis explaining the collapse and failure of northwest Atlantic groundfish populations

47 to recover (Bundy 2001, DFO 2003, Rice et al. 2003). In particular, harp seals (*Pagophilus*  
48 *groenlandicus*), the most abundant pinniped in the northwest Atlantic, have increased from  
49 less than 2 million in the early 1970s to almost 6 million today (Hammill & Stenson 2005).  
50 Harp seals summer in the Arctic, but migrate south along the Labrador coast in late autumn  
51 to overwinter off northeastern Newfoundland and in the NGSL (Fig. 2). The majority of  
52 the seals overwinter off the southeastern Labrador-northeastern Newfoundland coast (NAFO  
53 Divisions 2J3KL), while approximately 25-33% of the harp seals move into the NGSL (Stenson  
54 et al. 2002, 2003). Harp seals are generalist predators characterized by a diverse diet of fish  
55 and invertebrates (e.g. Lawson et al. 1998, Hammill & Stenson 2000). Stomach samples  
56 collected during the winter and spring in the NGSL indicate that their diet is highly variable  
57 both seasonally and geographically with average of approximately 5% made up by cod, mainly  
58 in the size range of 15-25 cm, although in some samples this proportion may be as high as 45%  
59 (Hammill & Stenson 2004, Stenson & Hammill 2004).

60 The correspondence between the decline of NGSL cod and the large increase in harp seal  
61 abundance (Fig. 3) has led to a hypothesis that harp seal predation may play a significant role  
62 in the cod lack of recovery (McLaren et al. 2001, DFO 2003). Seal predation on cod has been  
63 examined using bioenergetic models that derive cod annual consumption estimates by taking  
64 into account seasonal changes in feeding and variability in seal abundance, distribution and  
65 diet composition (Stenson et al 1997, Hammill & Stenson 2000, Stenson & Hammill 2004).  
66 However, because of the variable nature of the diets, these models have assumed an average  
67 proportion of cod in the diet, in spite of the marked changes in cod biomass that have been  
68 documented over the last 3 decades (MPO 2007). Assumptions about how predation changes in  
69 response to variations in prey abundance have been shown to be critical in predicting impacts  
70 of predators on commercially important prey (Yodzis 1994, Mackinson et al. 2003). However,  
71 individual and population level phenomena among predators and prey alike, such as search  
72 image, difficulty in finding prey, aggregation, and dispersion of predator and prey, can lead  
73 to significant departures from the linear assumptions about the relationship between prey  
74 abundance and the proportion of prey in the diet (Koen-Alonso 2006). In addition, harp seals  
75 can display strong preference for specific prey, although they have been shown to be neutrally

76 selective toward Atlantic cod (Lawson et al. 1998, Lindstrøm et al. 1998, Wathne et al. 2000).  
77 Modeling non-linear processes of the functional response of harp seals to cod appears therefore  
78 to be a key to modeling predation over a wide range of prey abundance such as observed in  
79 NGSL cod.

80 As described above, hypotheses related to fishing, seal predation, and environment have been  
81 proposed for the decline of NGSL cod but few studies have attempted to simultaneously address  
82 these hypotheses to determine which processes were at play at different times and how syner-  
83 gistic forcing could have led to the cod collapse. Here, a predator-prey model was constructed  
84 for NGSL cod, with mortality partitioned into components caused by harp seal predation and  
85 by other processes including an explicit influence of an integrated environmental forcer. The  
86 impact of seals on cod was modelled through a non-linear multi-age functional response similar  
87 to a multi-species functional response (Smout & Lindstrøm 2007). Cod condition was used as  
88 an index for the cumulative and lagged effects of environmental conditions related to factors  
89 such as food availability and temperature experienced by cod (Lambert & Dutil 1997a-b, Rätz  
90 & Lloret 2003). Changes in cod condition were assumed to affect natural mortality (Dutil  
91 & Lambert 2000) and fecundity, i.e. potential recruitment strength (Marteinsdottir & Begg,  
92 2002). The main objective of the analysis was to determine the relative impact of fishing, seal  
93 predation, and environment on the decline of NGSL cod by integrating these processes into a  
94 single consistent modeling framework. Such a framework is essential to both understand what  
95 happened to the stock and serve as a springboard for determining future states in the context  
96 of cod recovery and providing advice on exploitation of NGSL harp seal and cod populations.

## 97 **COD POPULATION MODEL**

98 Cod population dynamics were modelled through a separable statistical catch-at-age analysis  
99 (Fournier & Archibald 1982) without process error, where mortality was derived from three  
100 sources; the fishery, harp seal predation, and natural mortality due to causes other than harp  
101 seal predation (other predation, disease, virus, etc.). The Seal IMPact on Cod ABundance  
102 (SIMCAB) model considers 13 age classes of cod from 1 to 13 and 100% mortality occurs at

103 end of the 13th age class for all cohorts. Following Pope’s approximation (1972), mortality  
104 processes were modelled as successive steps, i.e. removals of cod by harp seal predation were  
105 taken instantaneously at the beginning of the year, fishing removals were taken instantaneously  
106 in the middle of the year, and residual mortality occurred between the 2 portions of the  
107 year (Table 2, D9-D11). The NGSL cod fishery changed considerably in the last decades,  
108 particularly after the 1994 moratorium when the winter mobile fishery dominated by otter  
109 trawlers was closed (Fréchet et al. 2003a). To account for changes in fishing pattern, 2  
110 periods of distinct fishery selectivity (partial recruitment) were considered: 1984-1993 and  
111 1994-2006. For consistency with knowledge available on NGSL cod dynamics and because  
112 of the convergence property of the virtual population analysis (VPA), the initial population  
113 age-structure was based on cod numbers estimated in 1984 from VPA (MPO 2007) and the  
114 model was run for the period 1984-2006. The parameters and variables used and the process  
115 and observation equations are given in Tables 1 and 2 respectively.

## 116 **A condition-dependent reproduction function**

117 Recruitment, i.e. fish numbers at age 1, was modelled through a Beverton and Holt (1957)  
118 stock-recruitment relationship based on an index of the reproductive potential of cod that  
119 was developed from an estimate of total egg production of the stock (TEP) (Table 2, D3).  
120 TEP accounted for yearly changes in cod numbers-at-age, sex ratio-at-age, the proportion of  
121 mature females-at-age, fecundity-at-age through length-at-age and condition factor of the fish.  
122 Previous studies have shown that length and condition factor have significant effects on the  
123 fecundity of cod in the NGSL (Lambert & Dutil 2000, Lambert et al. 2000). This dependence  
124 of fecundity on length and condition factor was used to develop a generalized linear model to  
125 determine the variation in the fecundity between 1984 and 2006 (Lambert 2008).

126 Sex ratio and maturity ogive data were derived from winter surveys conducted with the MV  
127 “Gadus Atlantic” (1984-1994) and spring samples available from the Groundfish Sentinel Fish-  
128 eries Program (<http://www.osl.gc.ca/pse/en/>). Fulton’s condition factor (Fulton 1902) based  
129 on length and weight data was also obtained from the winter surveys (1984-1994), the Sen-

130 tinal Fisheries Program (1995-2007), and supplementary research surveys conducted during  
131 the pre-spawning period in 1994, 1995, 1997, and 1998 (Dutil et al. 2003, Fréchet et al. 2007).

## 132 **A multi-age functional response of harp seals to cod**

133 To account for effects of prey abundance on predation, assuming no predator interference in  
134 the predation process, a multi-age functional response (FR), i.e. the number of cod consumed  
135 per seal per year, was considered (Table 2, D4). The FR of seal to cod is comprised of a  
136 maximum consumption rate for cod ( $J_{max}$ ), attack rates-at-age ( $\zeta_a$ ), and age-structured cod  
137 population. The shape parameter  $m$  of the FR determines the FR type, i.e.  $m = 1$  corresponds  
138 to a hyperbolic type 2 relationship and  $m > 1$  corresponds to a sigmoid shaped type 3 response  
139 (Koen-Alonso 2006).

140 The attack rates ( $\zeta_a$ ) were expressed relative to baseline attack rates ( $\zeta_a^0$ ) through a scaling  
141 factor ( $\rho$ ) that was estimated in the fitting process (see section 'Fitting the model'). Baseline  
142 attack rates were derived from cod biomass requirements for the NGSL harp seal population,  
143 cod abundance based on the matrix of numbers-at-age derived from virtual population analysis  
144 (MPO 2007), that was extended to ages 1-2 based on preliminary runs of the model without  
145 predation, and proportion and age-structure of cod in the seal diet in 1998-2001.

146 First, harp seal energetic requirements to maintain the seal population were estimated based  
147 on the daily gross energy intake of an individual harp seal derived from the allometric Kleiber  
148 equation that scales animal's metabolic rate to the 3/4 power of the animal's mass, and the  
149 age-structure of the seal population (Hammill & Stenson 2004, Stenson & Hammill 2004). 25%  
150 of the northwest harp seal population was assumed to move into the Gulf of St. Lawrence each  
151 year to spend on average 150 days per year in its northern part (Stenson et al. 2002, 2003).  
152 The biomass of cod consumed in 1998-2001 was then estimated based on the average energy  
153 of cod in winter, set equal to  $4.96 \text{ kJ g}^{-1}$  (Lambert & Dutil 1997a), and on the annual average  
154 proportion of cod in seals diet during November-March 1998-2001 (Table 3). The relative  
155 energetic contribution of cod to seal's diet was based on recovered otoliths and accounted for



156 harp seal population residency in the areas 4Ra-c, 4Rd-3Pn, and 4S of the NGSL (Fig. 2;  
157 Hammill & Stenson 2004). The annual number of cod consumed was then estimated based on  
158 the mean mass of cod consumed by harp seals and reallocated between cod age-groups based on  
159 information available from the analysis of seal stomach contents (Lawson et al 1998; Hammill  
160 & Stenson 2004). Lengths of cod ingested were estimated from otoliths found in seal stomachs  
161 and published fish length - otolith length regression equations (Hammill & Stenson 2004). Fish  
162 lengths were based on 671 otoliths found in 140 seal stomachs collected during November to  
163 March, 1989-2001 in the Gulf of St. Lawrence (Chassot et al. 2007). Harp seal size ranged  
164 from 102-166 cm. Otoliths collected from stomach contents were assumed to represent the size  
165 distribution of cod eaten. Proportions were estimated by fitting a mixture model of normal  
166 distributions to the prey size-frequency histogram by applying a constraint on the variance of  
167 cod size derived from DFO summer survey (Chassot et al. 2007).

168 The maximum consumption rate  $J_{max}$  was calculated based on the energy requirements of  
169 the NGSL harp seal population in 1998-2001 and a maximum value of cod proportion in seal  
170 stomachs derived from diet information available. Following the same approach as described  
171 above, fish biomass required to maintain the NGSL seal population was first calculated. Based  
172 on a maximum proportion of cod of 18.3% derived from the maximum proportions of cod in  
173 seals diet observed since 1980 and the relative residency time in each area of the NGSL, the  
174 potential maximum number of cod eaten per seal per year  $J_{max}$  was calculated. The baseline  
175 attack rates-at-age were finally calculated from the number of cod-at-age consumed per seal  
176 and cod numbers-at-age derived from the VPA in 1998-2001 (Table 2, D4). Since the baseline  
177 attack rates were based on a number of cod eaten calculated from the mean weight of cod in  
178 the reference period 1998-2001, the mass ratio  $\left(\frac{\overline{W}_t}{W_0}\right)$ , an index of cod weight, was included  
179 to account for yearly fluctuations in the mean mass of cod when calculating the number of cod  
180 predated by seals (Table 2, D5).

181 In order to illustrate the multi-age functional feeding response to changes in cod availability,  
182 a large range of levels of cod abundance were simulated for 2 distinct age-structures of the  
183 cod population. For each age-structure, cod-at-age consumed per seal per year were calculated

184 based on the model fitting results.

## 185 **Removals by natural mortality other than harp seal predation**

186 The rate of natural mortality excluding harp seal predation, i.e. residual mortality was mod-  
187 elled as an age-dependent function, in the form of a decreasing exponential as a function of  
188 age with a condition-dependent asymptote (Table 2, D6). The asymptote was modelled as a  
189 decreasing linear function of cod condition based on results of laboratory experiments relating  
190 natural mortality to cod condition (Dutil & Lambert 2000, Chassot et al. 2007). The condition  
191 factor which is an indicator of the energy content of cod was used here as an integrative index  
192 of environmental conditions (Lambert & Dutil 1997b). Alternative values for the asymptote  
193 of the residual mortality curve, i.e. a constant value of  $0.4 \text{ y}^{-1}$  and a time-dependent natural  
194 mortality derived from stock assessment models (MPO 2007), were considered to assess the  
195 impact on the results through the Akaike Information Criterion (AIC; Akaike 1974).

## 196 **Effects of seal predation vs. fishing on the reproductive potential**

197 Because seal predation and fishing are mortality sources for different cod age groups, a unit of  
198 mortality from each does not have the same impact on the population. Similarly to a multi-  
199 fleet biomass-per-recruit analysis, effects of changes in harp seal predation vs. fishing mortality  
200 were investigated by calculating the spawning stock biomass per recruit (SSB/R) for different  
201 levels of mortality. Such an analysis focuses on the long-term survival of a recruit and does  
202 not consider the linkage between the reproductive potential and recruitment at age 1. Based  
203 on the mortality rates estimated and averaged over the period 2002-2006 and assuming all else  
204 constant, we examined the relative importance of seal predation mortality and fishing mor-  
205 tality on the long-term stock reproductive potential under two scenarios: a high productivity  
206 environment and a low productivity environment. Considering a range of fishing mortality  
207 multipliers from 0 to 2, cod SSB/R was calculated at equilibrium for distinct levels condition  
208 because condition was assumed to affect the asymptote  $\vartheta$  of the residual natural mortality in

209 the model. 3 levels of seal predation mortality were considered corresponding to 0.5, 1, and  
210 1.5 times the average harp seal predation mortality estimated in 2002-2006. 2 scenarios of  
211 conditions represented different environmental conditions for the cod population, with values  
212 of  $\vartheta = 0.26$  and  $\vartheta = 0.35$  corresponding to a high and a low productivity regime scenario,  
213 respectively.

214

## FITTING THE MODEL

### 215 **Abundance and catch data**

216 Stratified random bottom-trawl surveys have been conducted in the NGSL annually, in summer  
217 months, since 1984. A consistent time-series of numbers-at-age, accounting for the changes in  
218 research vessels, was used as abundance indices for cod for the period 1984-2006 (Bourdages  
219 et al. 2003, 2007, Chassot et al. 2007).

220 The catch-at-age matrix of cod (in numbers) was obtained from the last updated stock as-  
221 sessment carried out in February 2007 (MPO 2007). All quantities caught and landed for  
222 both commercial and recreational fisheries, excluding discards, were included in the data. In  
223 the present analysis, reported landings were considered as the “true” catch despite potential  
224 discarding and misreporting in the fishery (Fréchet et al. 2003b, Savenkoff et al. 2004).

### 225 **Statistical estimation approach**

226 Maximum likelihood methods were used to estimate SIMCAB parameters. The likelihood  
227 function of the model was derived from assumptions on the statistical distribution of the 4  
228 datasets acquired from independent sampling processes: total abundance index in number,  
229 proportion-at-age for the abundance indices, total catch in number, and proportion of catch-  
230 at-age (Fournier & Archibald 1982, Quinn & Deriso 1999) (Table 5).

231 A stochastic observation equation accounting for correlated errors among ages was considered  
232 to link the abundance indices to the population model (Myers & Cadigan 1995) (Table 5, S1).  
233 For some years, some abundance indices-at-age for very young (age 1) or old cod (ages 11-13)  
234 were not observed during the surveys. For these years, total survey indices were considered as  
235 right-censored data, providing less information to the likelihood component than real observa-  
236 tions (Lawless 2003). This is true for the sum over ages of the logarithms of the survey indices  
237 since these indices are extrapolated to the scale of the NGSL and are positive numbers greater  
238 than 1. The observed likelihood of total surveys indices was then composed of a product of  
239 densities and survival functions, where a survival function is defined as 1 minus the distribu-  
240 tion function (Table 6, L1). The survival factor in the likelihood represents the integration,  
241 over all its possible values, of the probability that a random total survey index is higher than  
242 an underreported observation. It is a common way of including partial data information in a  
243 statistical analysis, but rarely used in fisheries analysis (for a notable exception see Hammond  
244 and Trenkel 2005).

245 Proportion-at-age for the abundance indices and catch data were assumed to follow dirichlet  
246 distributions to account for correlations among age groups (Fournier & Archibald 1982) (Table  
247 5, S2 and S3). Total catch observed were considered mainly dependent on the fishing year  
248 and assumed measured with lognormal observation errors (Table 5, S4). The assumption of  
249 log-normal error distributions for survey and catch data was checked through the residuals.

250 The maximum likelihood estimates were found by minimizing the negative sum of the log of  
251 the 4 likelihood components (i.e. objective function) given in Table 6. Minimizations were  
252 performed with the Nelder-Mead simplex non-linear optimization algorithm implemented in  
253 the R package (R Development Core Team 2008). The parameter space  $\theta$  to estimate included  
254 37 parameters from 644 observations, an observation being a survey index or catch number  
255 for a given age group in a given year: 2 for the TEP-recruitment relationship ( $R_{max}$  and  
256  $r$ ), 2 for the FR ( $m$  and  $\varrho$ ), 2 for the residual mortality curve ( $\alpha$  and  $\beta$ ), 1 for the survey  
257 catchability ( $q$ ), 6 for the selectivity of survey and fishing gears ( $\gamma_s, \delta_s, \gamma_c^1, \delta_c^1, \gamma_c^2, \delta_c^2$ ), 23 for  
258 the fishing mortality rates ( $F_t$ ), and 1 for the standard deviation of the observation error in

259 the total survey data ( $\psi$ ) (see Table 1 for definition). The maximum likelihood estimate of the  
260 observation error variance in the total catch data ( $\sigma_c^2$ ) was:

$$\sigma_c^2 = 2 \left( \sqrt{1 + \frac{1}{T} \sum_{t=1}^T \left( \log \frac{C_t^*}{C_t} \right)^2} - 1 \right) \quad (1)$$

261 Where the notations used are given in table 4.

262 Confidence statements about parameters were inferred using parametric bootstrap methods  
263 accounting for bias (Efron & Tibshirani 1998). 100 bootstrap replicate samples with the same  
264 dimensions as the datasets described above were generated by drawing from the probability  
265 distributions characterised by the parameters estimated at the maximum likelihood. Replicate  
266 samples were then submitted to the estimation procedure to obtain bootstrap replicates of the  
267 estimator and calculate percentiles of the distribution.

## 268 Sensitivity analysis

269 To complete the bootstrap analysis that only considered uncertainty in the catch and survey  
270 data, a sensitivity analysis was performed to account for uncertainty in some input parameters  
271 of the predation model and harp seal population in the NGSL (Hammill & Stenson 2004,  
272 Stenson & Hammill 2004). In a first step, the sensitivity of the results was examined by  
273 independently increasing parameter values by 10% from a baseline value (values from the  
274 standard run), re-fitting the model to observations for each sensitivity run, and assessing the  
275 change between the new and baseline results through a measure of relative sensitivity (Stenson  
276 & Hammill 2004, Mohn & Bowen 1996). The sensitivity to 5 parameters was investigated:  
277 the proportion of the northwest Atlantic population of harp seals moving into the NGSL, the  
278 average proportion of cod in seal's diet in 1998-2001, the maximum proportion of cod in seal's  
279 diet, the average energy of cod, and the Kleiber multiplier used as a constant multiplicative  
280 parameter in the Kleiber equation (Table 7). In order to summarize the model results and  
281 because predation by harp seals was the main focus of the analysis, only results for mean cod

282 biomass removed by seals and mean predation mortality rate for ages 1-4 during 1984-2006  
283 were presented. In a second step, the impacts of setting the value of the functional response  
284 type  $m$  to 1, i.e. a hyperbolic type 2 relationship, were examined.

## 285 **RESULTS**

### 286 **Age-structure of cod in seal's diet**

287 The size frequency histogram showed that 95% of cod found in harp seal stomachs were less than  
288 38 cm (Fig. 4). The mixture model fitted well the cod size-frequency data as indicated by the  
289 significance level for the goodness-of-fit test ( $p > 0.05$ ), based on the chi-square approximation  
290 to the likelihood ratio statistic (Du 2002). The model showed that cod consumed were young  
291 cod, with ages 1-4 representing 26.5%, 48.5%, 18%, and 7% of the cod in the diet, respectively  
292 (Fig. 4).

### 293 **Cod population dynamics**

294 The model fitted the survey and catch data well and the minimum value for the objective  
295 function was 6,149.6, corresponding to an AIC value of 12,373 (Figs. 5-6). The variances of  
296 the observation errors in the survey and catch data were low, the values of  $\psi$  and  $\sigma_c$  used as  
297 proxies of coefficients of variation in the case of lognormal likelihoods being 5.0% and 2.7%,  
298 respectively. Consequently, confidence intervals around parameters estimated by bootstrap  
299 analysis were small (Table 1). Considering a constant asymptote of  $0.4 \text{ y}^{-1}$  and a variable  
300 asymptote for the residual mortality curve led to higher values of AIC (12,583 and 12,696  
301 respectively) than for a condition-dependent asymptote, indicating a better fit of the model  
302 when residual mortality was related to cod condition.

303 SIMCAB estimates indicated a strong decline in cod abundance for all age-groups from 1984  
304 to 1993, consistent with the collapse of the northern Gulf cod stock in the mid-1990s (Fig. 5).

305 Abundance indices for ages 4-8 then remained quite stable from the mid-1990s to the early  
306 2000s, the stock showing no clear sign of recovery despite the moratoria on fishing implemented  
307 in 1994-1996 and 2003 (Fig. 5). By contrast, abundance indices for cod aged 10-13 displayed  
308 increasing trends from the mid-1990s to the 2000s; these age-groups representing a very small  
309 proportion of the stock abundance but about 10% of the SSB on average for the period 1984-  
310 2006 (Fig. 5). The declines in abundance were accompanied by a decrease in SSB from more  
311 than 330,000 t in the early 1980s to a minimum of less than 30,000 t in 1994 and showed a  
312 small increase thereafter. SSB estimates were thus below the conservation biomass limit of  
313 80-90,000 t and indicated a clear case of recruitment overfishing for the stock.

314 The number of cod caught showed a strong decrease from the 1980s to the mid-1990s for ages  
315 5-10 that represent the large majority of the biomass harvested (Fig. 6). SIMCAB predictions  
316 appeared bumpy in the 1980s with catches underestimated in 1986 and 1988 for all age-groups  
317 (Fig. 6). This mismatch between catch observed and predicted seems mostly due to the  
318 exceptional and somewhat surprising abundance indices for ages 4-10 in 1987 (Fig. 5).

### 319 **Mortality components in different age-groups**

320 Total mortality divided into age-groups showed that sources and magnitude of mortality varied  
321 in time and age (Fig. 7). Young cod aged 1-2 experienced high natural mortality, varying  
322 between about  $1.9 \text{ y}^{-1}$  in the mid-1980s to more than  $2.4 \text{ y}^{-1}$  in the mid-1990s (Fig. 7a).  
323 Predation mortality by harp seals only represented a low proportion of mortality for ages 1-2  
324 throughout the period 1984-2006 but increased from about 6% in the 1980s to more than 12%  
325 in the mid-2000s.

326 In addition to seal predation and residual natural mortality, cod aged 3-6 were subject to  
327 fishing mortality, mainly in the 1980s (Fig. 7b). The decrease in fishing mortality from the  
328 mid-1990s coincided with the change in fishing pattern, i.e. the commercial fishery after 1994  
329 was only conducted by fixed gears (longlines, gill nets, and hand lines) targeting larger cod.  
330 Residual natural mortality showed an increase from the 1980s to the mid-1990s associated with

331 lower condition during this period; the rates in the most recent years were similar to the levels  
332 estimated in the mid-1980s. Seal predation mortality for ages 3-6 remained quite stable during  
333 1984-2006 with its proportion in total mortality increasing to more than 20% in the 2000s due  
334 to the decrease in fishing and residual natural mortality.

335 Cod aged 7-10 not preyed upon by harp seals showed high total mortality, particularly in the  
336 mid-1980s and early 1990s with rates higher than  $1.0 \text{ y}^{-1}$  (Fig. 7c). The high interannual  
337 variability in mortality rates was mainly due to the strong variations in fishing mortality that  
338 represented more than 65% of the total mortality in some years (Fig. 7c). The low fishing  
339 mortality values estimated in 1986 and 1988 could be due to the high abundance indices  
340 observed in 1987 (Fig. 5). As for ages 3-6, residual natural mortality was the highest in the  
341 mid-1990s, corresponding to a low-condition period for cod. Although fishing mortality rates  
342 showed a decreasing trend in the 2000s, the values remained high for some years relative to  
343 the low abundance of the stock, particularly in 2005.

#### 344 **Harp seal predation**

345 Predation mortality rates for cod aged 1-4 showed a steady increase alongside the increase  
346 in harp seal abundance from  $0.1 \text{ y}^{-1}$  in 1984 to more than  $0.3 \text{ y}^{-1}$  in 2000 (Fig. 8). The  
347 rates then decreased to remain quite stable around  $0.25 \text{ y}^{-1}$  in the most recent years. Cod  
348 biomass removed annually by seal predation was estimated to be around 10,000 t from 1985  
349 to 1995. Removals then showed an increasing trend with a peak above 30,000 t in 2000 and  
350 a temporally averaged median of 16,000 t in the 2000s. The high interannual variability in  
351 biomass removed was mainly due to the variations in abundance of cod aged 1-2 that represent  
352 the major age-groups preyed upon by harp seals.

353 The exponent of the multi-age functional response fitted to the data was estimated to be 2.13  
354 ( $\pm 0.015$ ), indicating a sigmoid shaped type 3 response characteristic of sharp changes in the  
355 relative importance of the cod age-groups consumed as the result of a relatively small change in  
356 their availability in the environment. The changes in biomass removed by seal predation were



357 then related to both changes in seal population energy requirements and changes in relative  
358 cod abundance between age-groups. For instance, the predicted multi-age functional response  
359 showed that cod age 2 would be the main target of harp seals when considering different levels  
360 of cod abundance described by the 1984 population age-structure (Fig. 9). Considering an  
361 age-structure similar to low abundance year of 1995, characterized by a high proportion of  
362 cod aged 1 in the population, would lead to a very different response of seals with age-1 cod  
363 becoming the major prey and the other age-groups quickly reaching their respective maximum  
364 in seal consumption (Fig. 9).

### 365 **Sensitivity analysis**

366 The sensitivity analysis showed that the results of the model concerning harp seal predation  
367 were robust to the assumptions made about the average energy of cod and the values of  
368 contribution of cod to seal's diet (Table 7). Increasing the contribution of cod to seal's diet  
369 in 1998-2001 did not affect the total biomass removed but led a relatively small increase in  
370 predation mortality rates by decreasing the cod population. Although the increase in maximum  
371 cod consumption led to an increase in the maximum rate of cod consumption  $J_{max}$ , it poorly  
372 affected harp seal predation. Predation appeared quite sensitive to a change in the size of  
373 the seal population and the Kleiber multiplier of the bioenergetic model (Table 7). Changes  
374 in input parameters could be counter-intuitive due to the complexity and non-linearity of  
375 the model; hence, the biomass removed by predation during 1984-2006 was decreased by an  
376 increase in average energy of cod while it led to an increase in the predation mortality rates.  
377 The relative sensitivity of predation rates could have a different sign than the sensitivity of  
378 biomass removed since predation rates depend both on the cod biomass removed and cod  
379 numbers-at-age.

380 Constraining the predation model to follow a hyperbolic type 2 relationship strongly affected  
381 the results of the model and resulted in a lower quality of fit than for the standard run (AIC  
382 = 12,697). Setting the shape parameter of the functional feeding response to 1 led to reduced  
383 harp seal predation and increased cod numbers-at-age and spawning stock biomass.

## 384 **Effects of seal predation vs. fishing on the reproductive potential**

385 The SSB/R curves showed that the reproductive potential of the NGS� cod was affected dif-  
386 ferently by changes in seal predation and fishing mortality (Fig. 10). The distinct productivity  
387 regimes represented by 2 levels of residual mortality modified the values of SSB/R, with low  
388 productivity leading to lower values of SSB. In both cases of productivity regimes, the nearly  
389 horizontal curves indicated that changes in SSB/R would be poorly modified by changes in fish-  
390 ing mortality while variations in seal predation mortality would have a higher effect on SSB/R  
391 (Fig. 10). This result was mainly due to the low fishing mortality rate of  $0.15 \text{ y}^{-1}$  estimated  
392 during 2002-2006. Considering the harp seal predation mortality estimated in 2002-2006 and a  
393 low residual mortality (i.e. a high productivity regime), the SSB/R was estimated to be equal  
394 to 129 g (Fig. 10 - intermediate dashed line curve). In this case and for a fishing mortality  
395 fixed at the level of 2000-2006, a decrease in the predation mortality rate from the mean value  
396 of  $0.26 \text{ y}^{-1}$  in 2000-2006 to a level around  $0.13 \text{ y}^{-1}$  observed in the mid-1980s would allow  
397 almost doubling the SSB/R (Fig. 10 - upper dashed line curve). Changes in fishing mortality  
398 assuming constant predation mortality would be less beneficial to the SSB, a decrease from  
399 the fishing mortality rate of  $0.15 \text{ y}^{-1}$  in 2002-2006 to a low rate of  $0.03 \text{ y}^{-1}$  increasing the  
400 SSB from 129 g to 171 g per recruit. Considering a low productivity regime strongly decreased  
401 the SSB/R that was estimated to be 73 g in 2000-2006, due to the higher residual mortality  
402 decreasing the probability of young cod to reach maturity (Fig. 10 - intermediate solid line  
403 curve). In this case, the SSB/R for a predation mortality of  $0.13 \text{ y}^{-1}$  and constant fishing  
404 mortality would only be 131 g (Fig. 10 - upper solid line curve). For the 2002-2006 predation  
405 mortality, a decrease in fishing mortality would slowly increase the SSB/R to a maximum of  
406 97 g in the situation of a moratorium.

407

## **DISCUSSION**

408 The SIMCAB statistical catch-at-age model was developed to assess the effects of predation  
409 by the northwest Atlantic harp seal population on NGS� cod by estimating the relative im-

410 portance of different sources of mortality that affected the stock during a period of collapse  
411 and non-recovery. SIMCAB is a consistent modeling framework which combines the effects of  
412 fishing, predation mortality and environment (through the condition index) on cod population  
413 dynamics. SIMCAB showed that the increase in harp seal abundance during 1984-2006 re-  
414 sulted in an increase in predation mortality for cod age-groups targeted by seals. Despite the  
415 increasing importance of harp seal predation in cod mortality, the collapse of the NGS� cod  
416 stock appears mainly due to the combination of high fishing mortality rates and poor environ-  
417 mental conditions that affected the productivity of the population in the early to mid-1990s  
418 and led to the current state of recruitment overfishing. SIMCAB results also show, however,  
419 that current levels of predation mortality could increase stock rebuilding time by decreasing the  
420 probability of fish to reach maturity. Although cod condition has improved in recent years, the  
421 current lack of recovery of the NGS� cod seems due mainly to the very low spawner biomass,  
422 driven both by the fishery inherently targeting larger fish and increased harp seal predation.  
423 Within an ecosystem perspective, management scenarios based on a decline in seal population  
424 to promote stock rebuilding should however consider environmental conditions, prey availabil-  
425 ity for seals and other cod predators of the NGS� that all affect cod productivity and might  
426 give rise to unexpected outcomes with a decline in seal abundance.

## 427 **Multispecies modeling and functional response**

428 Separating the sources of mortality affecting animal populations is a major issue in ecology  
429 and has a long history in fisheries science (Andersen & Ursin 1977). This is particularly  
430 important in the context of multispecies management as predator control has been proposed  
431 for some time as a beneficial tool for fisheries (Flaaten 1988, FRCC, 1999). Multispecies  
432 models are useful tools to address the issues raised about the effects of marine mammals on  
433 fisheries (Yodzis 1998). In particular, predator-prey models have been used to evaluate the  
434 effects of fur seal (*Arctocephalus pusillus pusillus*) predation on hakes (*Merluccius capensis* and  
435 *Merluccius paradoxus*) in the Benguela system (Punt & Butterworth 1995) and more recently  
436 the effects of northwest Atlantic grey seal (*Halichoerus grypus*) predation on cod population  
437 dynamics and causes of collapse and non recovery on the eastern Scotian shelf (ESS) have been

438 explored (Mohn & Bowen 1996, Fu et al. 2001, Trzcinski et al. 2006). In the Benguela system,  
439 modeling the biological interaction between cape fur seals and the cape hakes suggested that  
440 an initial reduction in seal numbers would be beneficial, but the resulting decrease in seal  
441 consumption would also result in increased predation by a less commercially important hake  
442 species *M. capensis* on the more commercially important hake *M. paradoxus*, resulting in little  
443 net benefit (Punt and Butterworth 1995). Among northwest Atlantic grey seals, the first  
444 modeling approach included predation mortality explicitly in cohort analysis and showed that  
445 grey seals had little effect on the collapse of cod on the ESS (Mohn & Bowen 1996). By  
446 contrast, more recent analyses suggested that increasing predation mortality since the 1990s  
447 could affect the survival of immature cod and contribute to the failure of the ESS and southern  
448 Gulf of St. Lawrence cod stocks to recover (Fu et al. 2001, Chouinard et al. 2005, Trzcinski  
449 et al. 2006).

450 The SIMCAB model is an age-structured population dynamics model structurally similar to  
451 the models used for grey seals on the ESS but it differs in several important points. A major  
452 difference with the models from Mohn & Bowen (1996) and Fu et al. (2001) but addressed  
453 by Trzcinski et al. (2006) concerns the linkage between cod abundance and seal consumption  
454 through the functional feeding response. The form of the FR is a key issue in multispecies  
455 modeling, especially at low abundance, because it defines how predators impact their prey as  
456 a function of prey abundance (Yodzis 1994, Mackinson et al. 2003). Most multispecies models  
457 such as the Multi-Species Virtual Population Analysis (MSVPA; Magnússon 1995) are based  
458 on hyperbolic type 2 functional relationships between prey abundance and predation rates  
459 (for a review see Pláganyi 2007). These models assume constant ration formulations where  
460 per-capita consumption is set equal to the predator's required daily ration, consistent with  
461 the interpretation that feeding selectivities are independent of prey abundance. A sigmoidal  
462 functional response is however likely more appropriate when modeling generalist predators  
463 (Magnússon & Pálsson, 1991) and recent analyses based on field data tend to support a type  
464 3 relationship (Middlemas et al. 2006, Kempf et al. in press). In Ecopath with Ecosim  
465 models (Walters et al. 1997), the functional form of interactions is based on the 'foraging  
466 arena' concept that allows exploring alternative forms of functional response but biological

467 and ecological justifications about this concept remain controversial (Pláganyi 2007).

468 In the present analysis, cod age-groups were considered distinct prey, consistently with the  
469 opportunistic behaviour of harp seals (Hammill & Stenson 2000) and the size-based structure  
470 of marine food webs that regulates predation processes (e.g. Shin & Cury 2004). The multi-age  
471 FR fitted to the data was a sigmoid shaped type 3 response, suggesting that the response of  
472 seals to changes in cod abundance might be complex and highly non-linear, due to predator  
473 preferences and potential for prey switching as a function of the relative abundance of prey  
474 types in the environment. A sigmoidal shape for the FR would be consistent with the available  
475 knowledge on the spatio-temporal and vertical distribution of cod that differs between cod  
476 age-groups, juvenile cod being generally found in shallower waters and closer to the shore  
477 than adult cod (e.g. Castonguay et al. 1999). Based on survey conducted on board the MV  
478 “Gadus Atlantica” in winter from 1978 to 1994, young cod age-groups have been shown to  
479 occupy different depths and areas in the NGS (DFO unpublished data), which could favor  
480 the emergence of a type 3 FR. Cod migration and distribution patterns might also change in  
481 time in relation to changes in environmental conditions affecting their habitat (Castonguay et  
482 al. 1999).

483 The sensitivity analysis showed that a type 2 FR would be less consistent with both the data  
484 and outputs from other predation models (Hammill & Stenson 2004, Duplisea & Hammill  
485 2006). Although cod predation removals were of the same order of magnitude as those found  
486 with models based on linear assumptions about seal consumption (Hammill & Stenson 2004,  
487 Duplisea & Hammill 2006), the use of a FR led to different conclusions by accounting for  
488 changes in cod age-groups abundance in time. Our results are consistent with Middlemas et  
489 al. (2006) who provided empirical support for a type 3 FR of seals to Atlantic salmon in an  
490 estuarine system in Scotland; however, in a multispecies or multi-age context, the interpretation  
491 of the FR type is not straightforward as the changes in alternative prey abundance influence  
492 the response of the predator to the availability of any prey (Smout & Lindstrøm 2007). This  
493 will be important for harp seals who have been shown to show strong preference for pelagic  
494 species such as capelin (*Mallotus villosus*) but are neutrally selective towards cod (Lawson et

495 al. 1998). Attempts to estimate statistical parameters of a FR have revealed the difficulty  
496 of relating microscale observations of stomach contents to the effective feeding responses of  
497 marine species at the macroscale (Pláganyi 2007). This is further complicated in the case of  
498 harp seal because biomass estimates of important alternative prey such as capelin, sandlance  
499 (*Ammodytes* sp), Arctic cod (*Boreogadus saida*) and krill (*Euphausidae* sp) are not available  
500 for the NGSL. New data collection, experiments, and analyses such as conducted by Kempf  
501 et al (in press) may help clarify harp seal sigmoidal feeding behaviour since modeling such  
502 non-linear processes is key to explore the potential impact of management scenarios based on  
503 a decline in seal population.

#### 504 **Cod population dynamics**

505 Despite major differences with the current method used to assess the cod stock status (e.g.  
506 no commercial data included in the analysis), SIMCAB estimates of cod abundance, fishing  
507 mortality, and SSB accorded with the last assessment of the northern Gulf cod (MPO 2007).  
508 Results were consistent with the collapse of the stock in the late 1980s and the current lack of  
509 recovery (MPO 2007). In the present analysis, cod abundance in the initial year 1984 was based  
510 on the numbers-at-age provided by VPA results, considering that the convergence property of  
511 this method would give reliable estimates of abundance in the past (Jones 1961). Setting  
512 the initial cod numbers helped in the estimation of the parameters linking cod abundance to  
513 survey abundance indices, i.e. catchability and selectivity. Although a statistically 'optimal'  
514 solution was obtained through the fitting process, the information provided to the model was  
515 insufficient to distinguish between the level of residual mortality and magnitude of recruitment  
516 at age 1, i.e. there could be a correlation between the maximum recruitment ( $R_{max}$ ) and the  
517 residual mortality parameters ( $\alpha$  and  $\beta$ ). Such a correlation could affect the absolute values of  
518 residual mortality at age 1 but would not modify the model results for the other cod age-groups  
519 for which information is provided through survey data.

520 A novel aspect of our cod population model is a stock-recruitment relationship based on egg  
521 production, which for cod is considered a better measure of the true reproductive potential

522 of the stock than spawning biomass (Marshall et al. 2006). The high interannual variations  
523 in abundance indices for cod aged 1-2 included both natural survival variability of cod larvae  
524 and juveniles, and sampling noise associated with the research vessel gear that does not well  
525 select very small cod ( $< 15$  cm) predated by seals. In the context of trophic interactions  
526 where small fish form the bulk of the diet of many predators, collecting data on fish larvae and  
527 juveniles abundance appears as a key issue to explain the factors driving prerecruit survival  
528 and recruitment.

529 In addition, total egg production in our model was derived from a statistical model that related  
530 egg production to length-at-maturity and condition of mature females (Lambert 2008). Hence,  
531 recruitment in the model accounted for changes in growth of cod that could be density and/or  
532 temperature dependent (Swain et al. 2003), as well as changes in environment that could affect  
533 egg productivity through cod condition. Natural mortality other than harp seal predation was  
534 assumed age- and condition-dependent based on a comparative analysis between laboratory  
535 feeding experiments and wild cod collected in the Gulf of St. Lawrence in the 1990s (Lambert  
536 & Dutil 1997a). The residual natural mortality term for old ages was consistent with temporal  
537 patterns estimated with virtual population analysis (Grégoire & Fréchet 2005). Accounting  
538 for effects of condition on residual mortality led to a better fit of the model and suggested that  
539 the decline in the condition and energy reserves of cod during the 1990s may have lowered the  
540 productivity of the stock and contributed to its collapse (Lambert & Dutil 1997a).

541 In contrast to the VPA, SIMCAB allows the separation of seal predation mortality from fishing  
542 mortality and other mortality through time and accounts for environmental influences on  
543 mortality and cod fecundity. Modeling the mechanistic processes involved in predation is a  
544 valuable alternative to the classic assumption of constant natural mortality rates to include  
545 some ecosystem components in stock assessment methods (e.g. Lindstrøm et al. 2002) and  
546 progressively move toward ecosystem-based fishery management (Pikitch et al. 2004). This  
547 seems particularly critical for conducting stock projections to evaluate the expected effects of  
548 fishery management rules (e.g. Hollowed et al. 2000) in a context where predation is suspected  
549 to delay the NGSL cod stock recovery.

## 550 Cod recovery within an ecosystem perspective

551 The impact of changes in predation and fishing mortality on the SSB/R was assessed for  
552 different values of residual natural mortality assumed here to represent different environmental  
553 conditions. Diagnostics about a decline in seal predation were then shown to be dependent  
554 on the productivity of the stock related to the environment (Dutil et al. 2003). Residual  
555 mortality affecting young cod was shown to be more important than harp seal predation but  
556 not explicitly modelled in the present analysis. Predation mortality has been shown to be the  
557 dominant source of mortality for small cod ( $\leq 35$  cm) in the 1990s and 2000s, representing  
558 about 95% of total mortality (Savenkoff et al. 2006). In the NGSL, small cod have been shown  
559 to be preyed upon by large cod through cannibalism, other large demersals (e.g. white hake  
560 *Urophycis tenuis*), grey seals and boreal cetaceans including mysticetes such as minke whale  
561 (*Balaenoptera acutorostrata*) and odontocetes such as white-beaked dolphins (*Lagenorhynchus*  
562 *albirostris*) (Morissette et al. 2006, Savenkoff et al. 2004, 2006, 2007). While cannibalism  
563 decreased in the last decades due to the decline in adult stock biomass, it still represented an  
564 important component of mortality for small cod in the early 2000s (Savenkoff et al. 2007). A  
565 high proportion of the total mortality of cod aged 3-10 was also not explained by harp seal  
566 predation or fishing and attributed to residual mortality. This residual mortality includes both  
567 a predation mortality component due to large cod predators such as grey seals and another  
568 mortality component that could involve starvation, disease, and parasites (Savenkoff et al.  
569 2004, 2006). In SIMCAB, mortality linked to these latter factors was related to cod condition  
570 through the asymptote of residual mortality, assuming that fish in bad condition would have  
571 little energy reserves to survive over the winter months or critical stages of their life cycle  
572 (Lambert and Dutil 1997, Dutil et al. 1999).

573 Based on the large database of cod stomachs collected by Fisheries and Oceans Canada dur-  
574 ing numerous research and commercial fishing-vessel surveys since the mid-1980s, modelling  
575 explicitly cannibalism within SIMCAB could help separating the components of residual mor-  
576 tality for small cod. Following the one-way interaction modelled between harp seal and cod,  
577 predation by other species than harp seals such as grey seals could also be included in the



578 model if suitable data are available. The use of ecosystem models such as mass-balanced and  
579 end-to-end models (Travers et al. 2007) in complement to SIMCAB would allow to represent  
580 the full complexity of the food web, to identify the major factors affecting cod mortality, to  
581 include bottom-up effects of prey availability on production and mortality, and eventually to  
582 compare simulation outputs for assessing the robustness of the results.

583 Long-term projections based on the SSB/R analysis did not consider issues of recruitment in  
584 a context where the low reproductive potential of the stock associated with a low productivity  
585 are major factors explaining its lack of recovery (Dutil et al. 2003, Shelton et al. 2006).  
586 Projections performed with SIMCAB to estimate recovery time should also account for the  
587 effects of condition on recruitment (Lambert et al. 2000, Marteinsdottir & Begg 2002) and  
588 growth (Dutil et al. 1999) that would also modify the response of the cod population to  
589 the implementation of management rules based on a decline in seal population. Considering  
590 alternative projection scenarios based on different harvest control rules for the commercial cod  
591 fishery and different environmental regimes is important to determine the utility of managing  
592 seals to increase the recovery rate of the NGSL cod population.

## 593 CONCLUSION

594 The main objectives of this study were to examine the relative impacts of fishing, environmental  
595 conditions and seal predation on the decline and recovery of the NGSL cod population within  
596 a consistent modeling framework. Within this framework harp seals could play an important  
597 role in the recovery of NGSL cod. This framework lends itself to adding additional components  
598 which might result in different conclusions. For example, harp seals are generalist predators  
599 and their impact will be affected by the availability of alternative prey. Elsewhere it has been  
600 shown that incorporating even only a few other prey alternatives may result in different and  
601 unexpected outcomes (Punt & Butterworth 1995, Morissette et al. 2006, Matthiopoulos et al.  
602 2007). Therefore, although harp seals may have an important impact on recovery of NGSL cod  
603 population, efforts to manage seals with this objective in mind may not achieve the expected

604 outcomes because of other components within the NGSL marine ecosystem.

605 *Acknowledgements.* We are grateful to P Carter, S Turgeon, D McKinnon, W Penney, B  
606 Stockwood, D Wakeham, and H Bourdages for providing data on seal diet and cod abundance  
607 indices. K Trzcinski and B Mohn made useful suggestions on the modeling approach, M-N  
608 Bourassa and JF Gosselin helped in map drawing. The work benefited from discussions with  
609 A Fréchet, P Schwab, J Gautier, and M Koen-Alonso. Comments by 3 anonymous referees  
610 greatly helped to improve and to clarify the manuscript. This work was financed by DFO's  
611 species at risk funding program (SARCEP).

## 612 **LITERATURE CITED**

613 Akaike H (1974) A new look at the statistical model identification. *IEEE Trans Auto Contr*  
614 19:716–723

615 Andersen KP, Ursin E (1977) A multispecies extension to the Beverton and Holt theory of  
616 fishing, with accounts of phosphorus circulation and primary production. *Meddelelser Fra*  
617 *Danmarks Fiskeri- og Havundersøgelse*, NS Vol 7:319–435

618 Beverton RJH, Holt SJ (1957) On the dynamics of exploited fish populations. *Fisheries inves-*  
619 *tigations series 2.* Ministry of Agriculture, Fisheries and Food. London, UK

620 Bourdages H, Archambault D, Morin B, Fréchet A, Savard L, Grégoire F, Bérubé M (2003)  
621 Preliminary results from the groundfish and shrimp multidisciplinary survey from August 2003  
622 in the northern Gulf of St. Lawrence. *DFO Can Sci Advis Sec Res Doc 2003/078*

623 Bourdages H, Savard L, Archambault D, Valois S (2007) Résultats de l'expérience de la pêche  
624 comparative d'août 2004 et 2005 dans le nord du golfe Saint-Laurent, entre le NGCC Alfred  
625 Needler et le NGCC Teleost. *Rapp tech can sci halieut aquat 26xx : ix + 57p*

626 Bundy A (2001) Fishing on ecosystems: the interplay of fishing and predation in Newfoundland-  
627 Labrador. *Can J Fish Aquat Sci* 58:1153–1157

628 Castonguay M, Rollet C, Fréchet A, Gagnon P, Gilbert D, Brêthes JC (1999) Distribution  
629 changes of Atlantic cod (*Gadus morhua* L.) in the northern Gulf of St Lawrence in relation to  
630 an oceanic cooling. ICES J Mar Sci 56: 333–344

631 Chassot E, Caskenette A, Duplisea DE, Hammill MO, Bourdages H, Lambert Y, Stenson GB  
632 (2007) A model of predation by harp seals (*Phoca groenlandica*) on the northern Gulf of St.  
633 Lawrence stock of Atlantic cod (*Gadus morhua*). DFO Can Sci Advis Sec Res Doc 2007/066

634 Chouinard GA, Swain DP, Hammill MO, Poirier GA (2005) Covariation between grey seal  
635 (*Halichoerus grypus*) abundance and natural mortality of cod (*Gadus morhua*) in the southern  
636 Gulf of St. Lawrence. Can J Fish Aquat Sci 62:1991–2000

637 DFO (2003) Proceedings of the zonal assessment meeting - Atlantic cod. DFO Can Sci Advis  
638 Sec Proceed Ser 2003/021

639 Duplisea DE, Hammill MO (2006) An approach to modelling the influence of harp seal (*Phoca*  
640 *groenlandica*) predation on decline and recovery of the Northern Gulf of St. Lawrence cod  
641 (*Gadus morhua*). ICES CM 2006/L: 02

642 Dutil J-D, Castonguay M, Gilbert D, Gascon D (1999) Growth, condition, and environmental  
643 relationships in Atlantic cod (*Gadus morhua*) in the northern Gulf of St. Lawrence and  
644 implications for management strategies in the Northwest Atlantic. Can J Fish Aquat Sci  
645 56:1818–1831

646 Dutil J-D, Y Lambert (2000) Natural mortality from poor condition in Atlantic cod (*Gadus*  
647 *morhua*). Can J Fish Aquat Sci 57:826–836

648 Dutil J-D, Gauthier J, Lambert Y, Fréchet A, Chabot D (2003) Cod stocks rebuilding and fish  
649 bioenergetics: low productivity hypothesis. DFO Can Sci Advis Sec Res Doc 2003/060

650 Efron B, Tibshirani RJ (1998). An introduction to the Bootstrap Chapman & Hall

651 Flaaten O (1988) The economics of multispecies harvesting. Springer-Verlag, Berlin, Germany

652 Fournier D, Archibald CP (1982) A general theory for analyzing catch at age data. Can J Fish  
653 Aquat Sci 39:1195–1207

654 FRCC (1999) Conservation requirements for the Gulf of St. Lawrence groundfish stocks and  
655 cod stocks in divisions 2GH and 3Ps and science priorities. Letter to Minister, April 1999

656 Fréchet A, Butler R, Hussey E, Kennedy J, Rose G (2003b) Quality at landing in relation to  
657 gears used in the 3Pn, 4R cod fishery, 2000-2002. DFO Can Sci Advis Sec Res Doc 2003/066

658 Fréchet A, Gauthier J, Schwab P, Bourdages H, Chabot D, Collier F, Grégoire F, Lambert  
659 Y, Moreault G, Pageau L, Spingle J (2003a) The status of cod in the Northern Gulf of St.  
660 Lawrence (3Pn, 4RS) in 2002. DFO Can Sci Advis Sec Res Doc 2003/065

661 Fréchet A, Gauthier J, Schwab P, Bourdages H, Tournois, C, Spingle J, Way M, Collier F  
662 (2007) The status of cod in the Northern Gulf of St. Lawrence (3Pn, 4RS) in 2006. DFO Can  
663 Sci Advis Sec Res Doc 2007/068

664 Fu C, Mohn R, Fanning LP (2001) Why the Atlantic cod (*Gadus morhua*) stock off eastern  
665 Nova Scotia has not recovered. Can J Fish Aquat Sci 58:1613–1623

666 Fulton T (1902) Rate of growth of seas fishes. Sci Invest Fish Div Scot Rept 20

667 Grégoire F, Fréchet A (2005) Calculation of northern Gulf of St. Lawrence cod (*Gadus morhua*)  
668 natural mortality for the 1990 to 2004 period. DFO Can Sci Advis Sec Res Doc 2005/019

669 Hammill MO, Stenson GB, Myers RA, Stobo WT (1998) Pup production and population  
670 trends of the grey seal (*Halichoerus grypus*) in the Gulf of St. Lawrence. Can J Fish Aquat  
671 Sci 55:423–430

672 Hammill MO, Stenson GB (2000) Estimated prey consumption by harp seals (*Phoca groen-*  
673 *landica*), hooded seals (*Cystophora cristata*), grey seals (*Halichoerus grypus*) and harbour seals  
674 (*Phoca vitulina*) in Atlantic Canada. J Northw Atl Fish Sci 26:1–23

675 Hammill MO, Stenson GB (2004) Estimated consumption of Atlantic cod (*Gadus morhua*)  
676 by harp seals (*Phoca groenlandica*), in NAFO Zone 4RS. DFO Can Sci Advis Sec Res Doc  
677 2004/093

678 Hammill MO, Lesage V, Carter P (2005) What do harp seals eat? Comparing diet composition  
679 from different compartments of the digestive tract with diets estimated from stable isotope  
680 ratios. Can J Zool 83:1365–1372

681 Hammill MO, Stenson GB (2005) Abundance of Northwest Atlantic harp seals (1960-2005).  
682 DFO Can Sci Advis Sec Res Doc 2005/090

683 Hammond TR, Trenkel VM (2005) Censored catch data in fisheries stock assessment, ICES J  
684 Mar Sci, 62:1118–1130

685 Hollowed AB, Ianelli JN, Livingston PA (2000) Including predation mortality in stock assess-  
686 ments: a case study for Gulf of Alaska walleye Pollock. ICES J. Mar. Sci 57:279–293

687 Hutchings JA (1996) Spatial and temporal variation in the density of northern cod and a  
688 review of hypotheses for the stock's collapse. Can J Fish Aquat Sci 53:943–962

689 Jones R (1961) The assessment of the long-term effects of changes in gear selectivity and fishing  
690 effort. Mar Res Scot 2: 1–19

691 Kempf A, Floeter J, Temming A (in press) Predator-prey overlap induced Holling type III  
692 functional response: an example from the North Sea fish assemblage. Mar Ecol Prog Ser

693 Koen-Alonso M (2006) A process-oriented approach to the multispecies functional response.  
694 In Rooney N, McCann KS, Noaks DLG (eds): From energetics to ecosystems: the dynamics  
695 and structure of ecological systems, Chapter 1, p 1-36

696 Lambert, Y. 2008. Why should we closely monitor fecundity in marine fish populations? J  
697 Northw Atl Fish Sci 41:93–106

- 698 Lambert Y, Dutil J-D (1997a) Condition and energy reserves of Atlantic cod (*Gadus morhua*)  
699 during the collapse of the northern Gulf of St. Lawrence stock. *Can J Fish Aquat Sci* 54:2388–  
700 2400
- 701 Lambert Y, Dutil J-D (1997b) Can simple condition indices be used to monitor and quantify  
702 seasonal changes in the energy reserves of Atlantic cod (*Gadus morhua*)? *Can J Fish Aquat*  
703 *Sci* 54:104–112
- 704 Lambert Y, Dutil J-D (2000) Energetic consequences of reproduction in Atlantic cod (*Gadus*  
705 *morhua*) in relation to spawning level of somatic energy reserves. *Can J Fish Aquat Sci*  
706 57:815–825
- 707 Lambert Y, Dutil J-D, Ouellet P (2000) Nutritional condition and reproductive success in wild  
708 fish populations. In: Norberg B, Kjesbu OS, Taranger GL, Andersson E, Stefansson SO (eds)  
709 *Proc 6th Int Symp Rep Phys Fish. Inst Mar Res & Univ Bergen, Bergen, Norway*, p 77-84
- 710 Lawless JF (2003) *Statistical Models and Methods for Lifetime Data*, 2nd edition. John Wiley  
711 and Sons, Hoboken
- 712 Lawson JW, Anderson JT, Dalley EL, Stenson GB (1998) Selective foraging by harp seals  
713 *Phoca groenlandica* in nearshore and offshore waters of Newfoundland, 1993 and 1994. *Mar*  
714 *Ecol Prog Ser* 163:1–10
- 715 Lindstrøm U, Harbitz A, Haug T, Nilssen KT (1998) Do harp seals *Phoca groenlandica* exhibit  
716 particular prey preferences? *ICES J Mar Sci* 55:941–953
- 717 Lindstrøm U, Haug T, Røttingen I (2002) Predation on herring, *Clupea harengus*, by minke  
718 whales, *Balaenoptera acutorostrata*, in the Barents Sea *ICES J Mar Sci* 59:58–70
- 719 Mackinson S, Blanchard JL, Pinnegar JK, Scott R (2003) Consequences of alternative func-  
720 tional responses in models exploring whale-fishery interactions. *Mar Mam Sci* 19:661–681
- 721 Magnússon KG, Pálsson OK (1991) Predator-prey interactions of cod and capelin in Icelandic

722 waters. ICES Mar Sci Symp 193:153–170

723 Marshall CT, Needle CL, Thorsen A, Kjesbu OS, Yaragina NA (2006) Systematic bias in  
724 estimates of reproductive potential of an Atlantic cod (*Gadus morhua*) stock: implications for  
725 stock-recruit theory and management. Can J Fish Aquat Sci 63:980–994

726 Marteinsdottir G, Begg GA (2002) Essential relationships incorporating the influence of age,  
727 size and condition on variables required for estimation of reproductive potential in Atlantic  
728 cod *Gadus morhua*. Mar Ecol Prog Ser 235:235–256

729 Matthiopoulos J, Graham K, Smout S, Asseburg C, Redpath S, Thirgood S, Hudson P, Har-  
730 wood J (2007) Sensitivity to assumptions in models of generalist predation on a cyclic prey.  
731 Ecology 88:2576–2586

732 McLaren I, Brault S, Harwood J, Vardy D (2001) Report of the eminent panel on seal man-  
733 agement. Department of Fisheries and Oceans, Ottawa, Canada, 143pp

734 Middlemas SJ, Barton TR, Armstrong JD, Thompson PM (2006) Function and aggregative  
735 response of harbour seals to changes in salmonid abundance. Proc R Soc London Ser B  
736 273:193–198

737 Mohn R, Bowen WD (1996) Grey seal predation on the eastern Scotian Shelf: modelling the  
738 impact on Atlantic cod. Can J Fish Aquat Sci 53:2722–2738

739 Morissette L, Hammill MO, Savenkoff C (2006) The trophic role of marine mammals in the  
740 northern Gulf of St. Lawrence Mar Mam Sci 22:74–103

741 MPO (2007) Évaluation du stock de morue du nord du golfe du Saint-Laurent (3Pn, 4RS) en  
742 2006. Secr can de consult sci du ministère des Pêches et des Océans (MPO), Avis sci 2007/003

743 Myers RA, Barrowman NJ, Hoenig JM, Qu Z (1996) The collapse of cod in Eastern Canada:  
744 the evidence from tagging data. ICES J Mar Sci 53:629–640

745 Myers RA, Hutchings JA, Barrowman NJ (1997) Why do fish stocks collapse? The example  
746 of cod in Atlantic Canada. *Ecol Applic* 7:91–106

747 Myers RA, Cadigan NG (1995) Statistical analysis of catch-at-age data with correlated errors.  
748 *Can J Fish Aquat Sci* 52:1265–1273

749 Pikitch E, Santora C, Babcock E, Bakun A, Bonfil R, Conover D, Dayton P, Doukakis P,  
750 Fluharty D, Heneman B, Houde E, Link J, Livingston P, Mangel M, McAllister M, Pope J,  
751 Sainsbury K (2004) Ecosystem-based fishery management. *Science* 305:346–347

752 Pope JG (1972) An investigation of the accuracy of Virtual Population Analysis using cohort  
753 analysis. *ICNAF Res Bull* 9:65–74

754 Punt AE, Butterworth DS (1995) The effects of future consumption by the Cape fur seal on  
755 catches and catch rates of the Cape hakes. 4. Modelling the biological interaction between  
756 Cape fur seals *Arctocephalus pusillus pusillus* and the Cape hakes *Merluccius capensis* and *M.*  
757 *paradoxus* *S Afr J Mar Sci* 16:255–285

758 Quinn TJ II, Deriso RB (1999) *Quantitative Fishery Dynamics*. Oxford, New York

759 Rätz H-J, Lloret J (2003) Variation in fish condition between Atlantic cod (*Gadus morhua*)  
760 stocks, the effect on their productivity and management implications. *Fish Res* 60:369–380

761 R Development Core Team (2008) *R: A Language and Environment for Statistical Computing*  
762 R Foundation for Statistical Computing Vienna, Austria ISBN 3-900051-07-0

763 Rice JC, Shelton PA, Rivard D, Chouinard GA, Fréchet A (2003) Recovering Canadian Atlantic  
764 cod stocks: the shape of things to come? *ICES CM* 2003/U: 06

765 Savenkoff C, Castonguay M, Chabot D, Fréchet A, Hammill MO, Morissette L (2006) Main  
766 prey and predators and estimates of mortality of Atlantic cod (*Gadus morhua*) in the northern  
767 Gulf of St. Lawrence during the mid-1980s, mid-1990s, and early 2000s. *Can Tech Rep Fish*  
768 *Aquat Sci* 2666



769 Savenkoff C, Castonguay M, Chabot D, Hammill MO, Bourdages H, Morissette L (2007)  
770 Changes in the northern Gulf of St. Lawrence ecosystem estimated by inverse modelling:  
771 evidence of a fishery-induced regime shift? *Estuar Coast Shelf Sci* 73:711–724

772 Savenkoff C, Castonguay M, Vézina AF, Despatie S-P, Chabot D, Morissette L, Hammill MO  
773 (2004) Inverse modelling of trophic flows through an entire ecosystem: the northern Gulf of  
774 St. Lawrence in the mid-1980s. *Can J Fish Aquat Sci* 61:2194–2214

775 Shelton PA, Healey, BP (1999) Should depensation be dismissed as a possible explanation  
776 for the lack of recovery of the northern cod (*Gadus morhua*) stock? *Can J Fish Aquat Sci*  
777 56:1521–1524

778 Shelton PA, Sinclair AF, Chouinard GA, Mohn R, Duplisea DE (2006) Fishing under low  
779 productivity conditions is further delaying recovery of Northwest Atlantic cod (*Gadus morhua*).  
780 *Can J Fish Aquat Sci* 63:235–238

781 Shin Y-J, Cury P (2004) Using an individual-based model of fish assemblages to study the  
782 response of size spectra to changes in fishing. *Can J Fish Aquat Sci* 61:414–431

783 Smout S, Lindstrøm U (2007) Multispecies functional response of the minke whale *Balaenoptera*  
784 *acutorostrata* based on small-scale foraging studies. *Mar Ecol Prog Ser* 341:277–291

785 Stenson GB, Hammill MO, Kingsley MCS, Sjare B, Warren WG, Myers RA (2002) Is there ev-  
786 idence of increased pup production in northwest Atlantic harp seals, *Pagophilus groenlandicus*?  
787 *ICES J Mar Sci* 59:81–92

788 Stenson GB, Hammill MO (2004) Quantifying uncertainty in estimates of Atlantic cod (*Gadus*  
789 *morhua*) consumption by harp seals (*Phoca groenlandica*). *DFO Can Sci Advis Sec Res Doc*  
790 2004/089

791 Stenson GB, Rivest L-P, Hammill MO, Gosselin J-F, Sjare B (2003) Estimating pup production  
792 of harp seals, *Pagophilus groenlandicus*, in the Northwest Atlantic. *Mar Mam Sci* 19:141–160

- 793 Swain DP, Sinclair AF, Castonguay M, Chouinard GA, Drinkwater KF, Fanning LP, Clark  
794 DS (2003) Density-versus temperature-dependent growth of Atlantic cod (*Gadus morhua*) in  
795 the Gulf of St. Lawrence and on the Scotian Shelf. *Fish Res* 59:327–341
- 796 Travers M, Shin Y, Jennings S, Cury P (2007) Towards end-to-end models for investigating  
797 the effects of climate and fishing in marine ecosystems. *Prog Oceanogr* 75:751–770
- 798 Trzcinski MK, Mohn R, Bowen WD (2006) Continued decline of an Atlantic cod population:  
799 how important is gray seal predation? *Ecol Applic* 16:2276–2292
- 800 Walters CJ, Christensen V, Pauly D (1997) Structuring dynamic models of exploited ecosys-  
801 tems from trophic mass-balance assessments. *Rev Fish Biol Fish* 7:139-172
- 802 Wathne JA, Haug T, Lydersen C (2000) Prey preference and niche overlap of ringed seals *Phoca*  
803 *hispida* and harp seals *P. groenlandica* in the Barents Sea. *Mar Ecol Prog Ser* 194:233–239
- 804 Yodzis P (1994) Predator-prey theory and management of multispecies fisheries. *Ecol Applic*  
805 4:51–58
- 806 Yodzis P (1998) Local trophodynamics and interaction of marine mammals and fisheries in the  
807 Benguela ecosystem. *J Anim Ecol* 67:635–658

808 **List of Figures**

809     1     Location of the northern Gulf of St. Lawrence (NAFO divisions 4RS) . . . . . 35

810     2     Distribution area of northwest Atlantic harp seals (*Pagophilus groenlandicus*) . . . . . 36

811     3     Harp seal abundance (black dots) ( $\pm$  standard deviation) derived from the model of Hammill  
812             and Stenson (2004) and cod spawning stock biomass (white diamonds) derived from virtual  
813             population analysis (MPO 2007) in the northern Gulf of St. Lawrence during 1984-2006 . . . . . 37

814     4     Size-frequency histogram of cod ingested derived from otoliths collected in seal stomachs with  
815             fitted mixture distribution representing cod of age 1-4 . . . . . 38

816     5     Survey numbers-at-age observed (cross) and predicted by the SIMCAB model (solid line).  
817             Scales were selected for readability along *y*-axes . . . . . 39

818     6     Catch-at-age observed (cross) and predicted by the SIMCAB model (solid line). Scales were  
819             selected for readability along *y*-axes . . . . . 40

820     7     Cumulative annual mortality rates for cod aged (a) 1-2, (b) 3-6, and (c) 7-10. Dark grey: seal  
821             predation mortality; light grey: natural mortality other than seal predation; white: fishing  
822             mortality. Scales were selected for readability along *y*-axes . . . . . 41

823     8     Annual predation mortality rate for cod aged 1-4 with 90% bootstrap confidence intervals . . . . . 42

824     9     Predicted multi-age functional response of harp seal to cod for 2 distinct age-structures of  
825             the population. Solid line corresponds to the age-structure of the year 1984; dashed line  
826             corresponds to the age-structure of the year 1995 characterized by a high abundance of cod  
827             aged 1. Grey area represents 90% bootstrap confidence intervals. Scales were selected for  
828             readability along *y*-axes . . . . . 43

829     10    Spawning stock biomass per recruit (SSB/R) as a function of fishing mortality for 3 different  
830             levels of harp seal predation and 2 distinct productivity regimes (see text for details) . . . . . 44




Figure 1: Location of the northern Gulf of St. Lawrence (NAFO divisions 4RS)




Figure 2: Distribution area of northwest Atlantic harp seals (*Pagophilus groenlandicus*)




Figure 3: Harp seal abundance (black dots) ( $\pm$  standard deviation) derived from the model of Hammill and Stenson (2004) and cod spawning stock biomass (white diamonds) derived from virtual population analysis (MPO 2007) in the northern Gulf of St. Lawrence during 1984-2006




Figure 4: Size-frequency histogram of cod ingested derived from otoliths collected in seal stomachs with fitted mixture distribution representing cod of age 1-4

# Cod abundance indices




Figure 5: Survey numbers-at-age observed (cross) and predicted by the SIMCAB model (solid line). Scales were selected for readability along y-axes






Figure 6: Catch-at-age observed (cross) and predicted by the SIMCAB model (solid line). Scales were selected for readability along  $y$ -axes




Figure 7: Cumulative annual mortality rates for cod aged (a) 1-2, (b) 3-6, and (c) 7-10. Dark grey: seal predation mortality; light grey: natural mortality other than seal predation; white: fishing mortality. Scales were selected for readability along  $y$ -axes




Figure 8: Annual predation mortality rate for cod aged 1-4 with 90% bootstrap confidence intervals




Figure 9: Predicted multi-age functional response of harp seal to cod for 2 distinct age-structures of the population. Solid line corresponds to the age-structure of the year 1984; dashed line corresponds to the age-structure of the year 1995 characterized by a high abundance of cod aged 1. Grey area represents 90% bootstrap confidence intervals. Scales were selected for readability along  $y$ -axes




Figure 10: Spawning stock biomass per recruit (SSB/R) as a function of fishing mortality for 3 different levels of harp seal predation and 2 distinct productivity regimes (see text for details)

Table 1: Parameters and variables used in SIMCAB. NS: not shown

Notation	Definition	Origin	Value	Equation
$B$	Cod biomass (t)	Calculated	NS	D1
$N$	Cod numbers (number)	Calculated	NS	D1-D5,D9,D11-D12
$w_a$	Cod weight-at-age (t)	Fixed	NS	D1-D2
$SSB$	Spawning stock biomass (t)	Calculated	NS	D2
$TEP$	Total egg production (number)	Calculated	NS	D3,D8
$\xi$	Sex ratio	Fixed	NS	D3
$\phi$	Proportion of maturing females	Fixed	NS	D1,D3
$f$	Fecundity (number of eggs cod <sup>-1</sup> )	Fixed	NS	D3
$A$	Last age-group	Fixed	13	D3
$\lambda$	Functional response of harp seals to cod (number seal <sup>-1</sup> year <sup>-1</sup> )	Calculated	Fig. 9	D4,D5
$\zeta$	Attack rate (number seal <sup>-1</sup> year <sup>-1</sup> )	Calculated	NS	D4
$\zeta^{ref}$	Reference attack rate (number seal <sup>-1</sup> year <sup>-1</sup> )	Calculated	NS	D4
$\rho$	Scaling factor of the attack rates	Estimated	0.021 (0.003)	D4
$J_{max}$	Maximum consumption rate (number seal <sup>-1</sup> year <sup>-1</sup> )	Calculated	996	D4
$m$	Shape parameter defining the functional response type	Estimated	2.13 (0.015)	D4
$\rho$	Number of feeding days spent by seals in Gulf each year	Fixed	150	D5

Table 1: Parameters and variables used in SIMCAB (continued)

Notation	Definition	Origin	Value	Equation
$\overline{W}_t$	Cod mean weight for ages targeted by seals (t)	Calculated	NS	D5
$\overline{W}_0$	Cod mean weight for ages targeted by seals in 1998-2001 (t)	Calculated	288e-6	D5
$S$	Seal numbers (number)	Fixed	Fig. 3	D5
$M$	Residual natural mortality rate ( $y^{-1}$ )	Calculated	Fig. 7	D6
$\alpha$	Intercept of the $M$ curve ( $y^{-1}$ )	Estimated	5.03 (0.013)	D6
$\beta$	Slope of the $M$ curve ( $y^{-1} \text{ a}^{-1}$ )	Estimated	0.90 (0.002)	D6
$\vartheta$	Asymptote of the $M$ curve ( $y^{-1}$ )	Calculated	NS	D6
$F$	Fishing mortality rate ( $y^{-1}$ )	Estimated	Fig. 7	D7
$s_{a,c}$	Partial recruitment	Calculated	NS	D7
$\gamma_c^1$	Shape parameter of the partial recruitment (1984-1993)	Estimated	1.95 (0.007)	D7
$\delta_c^1$	Age at which 50% of the individuals are vulnerable to fishing gear (1984-1993)	Estimated	5.39 (0.006)	D7
$\gamma_c^2$	Shape parameter of the partial recruitment (1994-2006)	Estimated	1.64 (0.017)	D7
$\delta_c^2$	Age at which 50% of the individuals are vulnerable to fishing gear (1994-2006)	Estimated	6.46 (0.016)	D7
$R$	Recruitment (number)	Calculated	NS	D8
$R_{max}$	Maximum number of recruits produced (number)	Estimated	1.9e9 (6.9e6)	D8
$r$	TEP needed to produce recruitment equal to $R_{max}/2$ (number of eggs)	Estimated	5.3e11 (1.3e11)	D8
$N'$	Cod numbers in the middle of the year (number)	Calculated	NS	D9-D10
$P$	Cod numbers predated by the seals (number)	Calculated	NS	D9

Table 1: Parameters and variables used in SIMCAB (continued)

Notation	Definition	Origin	Value	Equation
$C$	Fishery catch (number)	Calculated	Fig. 6	D10-D11,D14-D15
$I$	Abundance index (number)	Calculated	Fig. 5	D12
$\varsigma_{a,s}$	Survey selectivity-at-age	Calculated	NS	D12
$q$	Survey catchability	Estimated	1.38 (0.076)	D12
$\gamma_s$	Shape parameter of the survey selectivity	Estimated	2.99 (0.028)	D12
$\delta_s$	Age at which 50% of the individuals are vulnerable to the survey gear	Estimated	2.72 (0.007)	D12
$p_{a,t,s}$	Proportion of number-at-age in the survey	Calculated	NS	D13
$p_{a,t,c}$	Proportion of catch-at-age	Calculated	NS	D15



Table 2: Deterministic process and observation equations in the SIMCAB model.  $a$  and  $t$  index age and time respectively

---

<b>State moments</b>	
(D1)	$B_t = \sum_{a=1}^A w_a N_{a,t}$
(D2)	$SSB_t = \sum_{a=1}^A w_a N_{a,t} \phi_{a,t}$
(D3)	$TEP_t = \sum_{a=1}^A N_{a,t} \xi_{a,t} \phi_{a,t} f_{a,t}$
<b>Mortality components</b>	
(D4)	$\lambda_{a,t} = \frac{J_{max} \zeta_a N_{a,t}^m}{J_{max} + \sum_i \zeta_i N_{i,t}^m}$ with $\zeta_a = \varrho \zeta_a^0$
(D5)	$P_{a,t} = \frac{\lambda_{a,t} N_{a,t}}{\left(\frac{\bar{W}_t}{W_0}\right) S_t \frac{\rho}{365}}$
(D6)	$M_{a,t} = \vartheta_t + \alpha \exp\left(\frac{-a}{\beta}\right)$
(D7)	$F_{a,t} = \zeta_{a,c}^k F_t$ with $\zeta_{a,c}^k = \frac{1}{1 + \exp(-\gamma_c^k (a - \delta_c^k))}$
<b>Process functions</b>	
(D8)	$R_t = \frac{R_{max} TEP_{t-1}}{r + TEP_{t-1}}$
(D9)	$N'_{a,t} = (N_{a,t} - P_{a,t}) \exp(-M_{a,t}/2)$
(D10)	$C_{a,t} = N'_{a,t} (1 - \exp(-F_{a,t}))$
(D11)	$N_{a+1,t+1} = (N'_{a,t} - C_{a,t}) \exp(-M_{a,t}/2)$
<b>Observation functions</b>	
(D12)	$I_t = \sum_{a=1}^A q \zeta_{a,s} N_{a,t}$ with $\zeta_{a,s} = \frac{1}{1 + \exp(-\gamma_s (a - \delta_s))}$
(D13)	$p_{a,t,s} = \frac{I_{a,t}}{\sum_{a=1}^A I_{a,t}}$
(D14)	$C_t = \sum_{a=1}^A C_{a,t}$
(D15)	$p_{a,t,c} = \frac{C_{a,t}}{\sum_{a=1}^A C_{a,t}}$

---

Table 3: Proportion of Atlantic cod in harp seal diet weighted by their relative residency in each area during the period November-March. The year 1998 refers to the period November 1997-March 1998 and similarly for the other years

Year	Sample size			Proportion
	4Ra-c	4Rd-3Pn	4S	
1998	27	21	-	0.0506
1999	28	22	24	0.0538
2000	25	39	29	0.0382
2001	30	16	-	0.0377

Table 4: Parameters and variables used in the stochastic equations and likelihood components.  $a$ ,  $t$ ,  $s$ , and  $c$  index age, year, survey, and catch respectively

Notation	Definition	Equation
$I_{a,t}^*$	Observed abundance index	L1
$J_{a,t}^*$	log-observed abundance index	S1
$T_1$	Number of uncensored data, with $T_1 = \text{Card}(\bar{I})$	L1
$\Phi$	Cumulative standard normal distribution	L1
$\psi$	Standard deviation of the observation error in the survey data	L1
$n_{t,s}^*$	Observed total number of fish taken by the survey vessels	L2
$p_{a,t,s}^*$	Observed proportion of number-at-age, with $p_{a,t,s}^* = \frac{I_{a,t}^*}{\sum_{a=1}^A I_{a,t}^*}$	L2
$p_{a,t,s}$	Predicted proportion of number-at-age	L2
$T$	Number of years	L2-L4
$\Gamma$	Gamma distribution	L2-L3
$p_{a,t,c}^*$	Observed proportion of catch-at-age	L3
$p_{a,t,c}$	Predicted proportion of catch-at-age	L3
$n_{t,c}^*$	Total catch sampled to establish age-proportions	L3
$C_t^*$	Observed total catch	L4
$C_t$	Predicted total catch	L4
$\sigma_c$	Standard deviation of the observation error in the catch data	L4

Table 5: Definitions used to extend the deterministic model in Table 2 to a stochastic model. iid: independent and identically distributed;  $\sim$ : distributed as;  $\mathcal{N}$ : normal distribution; *Dir*: Dirichlet distribution

---

(S1)	$J_t^* = \sum_{a=1}^A J_{a,t}^* \stackrel{iid}{\sim} \mathcal{N}\left(\sum_{a=1}^A \log I_{a,t}(\theta), \psi^2\right)$	$\psi^2 = A\sigma^2 + A^2\tau^2$
	where $J_{a,t}^* = \log(I_{a,t}^*) = \log(I_{a,t}) + \epsilon_{a,t} + \eta_t$	$\epsilon_{a,t} \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2)$ $\eta_t \stackrel{iid}{\sim} \mathcal{N}(0, \tau^2)$
(S2)	$p_{a,t,s}^* \stackrel{iid}{\sim} \text{Dir}(p_{1,t,s}(\theta), \dots, p_{A,t,s}(\theta), n_{t,s}^*)$	
(S3)	$p_{a,t,c}^* \stackrel{iid}{\sim} \text{Dir}(p_{1,t,c}(\theta), \dots, p_{A,t,c}(\theta), n_{t,c}^*)$	
(S4)	$\log C_t^* \stackrel{iid}{\sim} \mathcal{N}\left(\log C_t(\theta) - \frac{\sigma_c^2}{2}, \sigma_c^2\right)$	

---

Table 6: Likelihood components of the SIMCAB model. The generic notation  $\theta$  represents the set of parameters to estimate

Equation	Likelihood
(L1)	$L(\{I_{a,t}^*\} \theta) = \underbrace{\frac{1}{(2\pi)^{\frac{T_1}{2}} \psi^{T_1}} \exp\left\{-\sum_{t \in \bar{I}} \frac{\left(\sum_{a=1}^A \log \frac{I_{a,t}^*}{I_{a,t}(\theta)}\right)^2}{2\psi^2}\right\}}_{\text{real observations}} \underbrace{\prod_{t \in I} \left\{1 - \Phi\left(\frac{\sum_{a=1}^A \log \frac{I_{a,t}^*}{I_{a,t}(\theta)}}{\psi}\right)\right\}}_{\text{censored observations}}$
(L2)	$L(\{p_{a,t,s}^*\} \theta) = \prod_{t=1}^T \Gamma(n_{t,s}^* + 1) \frac{\prod_{a=1}^A (p_{a,t,s}^*)^{n_{t,s}^* p_{a,t,s}(\theta)}}{\prod_{a=1}^A \Gamma(n_{t,s}^* p_{a,t,s}(\theta) + 1)}$
(L3)	$L(\{p_{a,t,c}^*\} \theta) = \prod_{t=1}^T \Gamma(n_{t,c}^* + 1) \frac{\prod_{a=1}^A (p_{a,t,c}^*)^{n_{t,c}^* p_{a,t,c}(\theta)}}{\prod_{a=1}^A \Gamma(n_{t,c}^* p_{a,t,c}(\theta) + 1)}$
(L4)	$L(\{C_t^*\} \theta) = \frac{1}{(2\pi)^{\frac{T}{2}} \sigma_c^T} \exp\left\{-\frac{1}{2\sigma_c^2} \sum_{t=1}^T \left(\log \frac{C_t^*}{C_t(\theta)} - \frac{\sigma_c^2}{2}\right)^2\right\}$

---

Table 7: Relative sensitivity (%) of biomass removed ( $BR$ ) by seals and predation mortality rate for cod aged 1-4 ( $M_p(1-4)$ ) to a 10% increase in input parameters

Parameter	Base value	$BR$	$M_p(1 - 4)$
% in the NGSL	25	67	129
% cod contribution to diet 1998-2001	4.53	1	25
% cod maximum contribution to diet	18.3	-6	-2
Average energy of cod ( $\text{kJ g}^{-1}$ )	4.96	-13	40
Kleiber multiplier (kJ)	293	67	129