



HAL
open science

Rodent borne diseases in Thailand: targeting rodent carriers and risky habitats

Vincent Herbreteau, Frédéric Bordes, Sathaporn Jittapalapong, Yupin Supputamongkol, Serge Morand

► To cite this version:

Vincent Herbreteau, Frédéric Bordes, Sathaporn Jittapalapong, Yupin Supputamongkol, Serge Morand. Rodent borne diseases in Thailand: targeting rodent carriers and risky habitats. *Infection Ecology & Epidemiology*, 2012, 2, 5 p. 10.3402/iee.v2iO.18637 . ird-00714369v1

HAL Id: ird-00714369

<https://ird.hal.science/ird-00714369v1>

Submitted on 4 Jul 2012 (v1), last revised 4 Jul 2012 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 *Infection Ecology & Epidemiology (in press)*

2

3 **Rodent-borne diseases in Thailand: targeting rodent carriers and risky habitats**

4

5 Vincent Herbreteau¹, Frédéric Bordes², Sathaporn Jittapalapong³, Yupin Supputamongkol⁴,

6 Serge Morand^{2,5}

7

8

9 ¹ Espace-Dev, IRD, Maison de la télédétection, Montpellier, France

10 ² Institut des Sciences de l'Evolution, CNRS-UM2, CC65, Université de Montpellier2,

11 34095 Montpellier, France

12 ³ Faculty of Veterinary Sciences, Kasetsart University, Bangkok, Thailand.

13 ⁴ Siriraj Hospital, Mahidol University, Bangkok, Thailand

14 ⁵ UR22 AGIRs, CIRAD, Campus International de Baillarguet

15 34398 Montpellier, France

16

17

18 **Corresponding author**

19 Serge Morand,

20 Institut des Sciences de l'Evolution, UMR CNRS-IRD-UM2, CC65,

21 Université de Montpellier 2, F-34095 Montpellier, France

22 Tel +33 (0)6 88 50 57 13

23 serge.morand@univ-montp2.fr

24

25

1 **Abstract**

2 We developed a simple method to target rodent reservoirs in Thailand by using published
3 studies screening microparasites in rodents from Thailand. Microparasite richness was
4 correlated with both rodent sample size and pathogens' screening effort. The residual
5 variation of this correlation helps at identifying major rodent reservoirs and potential risky
6 habitats.

7

8 **Key words:** microparasite richness, rodents, Southeast Asia, zoonosis, habitat,
9 prioritization.

10 **Running Title:** Targeting rodent-borne disease carriers in SEA

11

12

13 **Background**

14 Targeting reservoirs of zoonotic diseases is a challenge particularly in countries presenting
15 high biodiversity. Comparative analysis, which aims at investigating ecological and
16 evolutionary patterns among species, may help at indentifying wildlife reservoirs using
17 published studies.

18

19 **Introduction**

20 We aim at presenting a simple way to prioritize reservoir species in relation to their capacity
21 to carry multiple agents of potential diseases. We illustrate this simple method with rodent-
22 borne diseases in Thailand, a country that showed major outbreaks of several rodent-borne
23 diseases in the past years (1). We focused on rodents as carriers, vectors or reservoirs of
24 numerous zoonotic diseases, notably microparasites (2). Thailand presents the advantage to
25 harbour a rich biodiversity, although at threat (3), and to have been quite intensively surveyed

1 for rodent-borne diseases. Using published surveys investigating rodent-borne diseases in
2 rodents in this country we aim at identifying the major rodent species reservoirs, at least in
3 term of their capacity to host multiple pathogens. For this, we first investigate the relationship
4 between the pathogen species richness observed among rodent species and the screening
5 effort and, second, we use the residual variations of this relationship to prioritize rodents and
6 evaluate the potential risky habitats.

7

8 **Materials and methods**

9 We compiled surveys of microparasites investigated in rodents trapped in Thailand (from 27
10 references given in supplementary appendix). The data comprise a total of 17,358 rodents
11 from 18 species of murine rodents that have been investigated for a total of 10 microparasites
12 (Table 1). The microparasites were viruses, bacteria and protozoans. Viruses were:
13 Hantaviruses, Lymphocytic choriomeningitis virus (family *Arenaviridae*, genus *Arenavirus*),
14 Rabies virus and Hepatitis E virus. These viruses are directly transmitted between rodent
15 hosts and are all major pathogens of humans. Three bacteria were investigated: *Leptospira*
16 *spp.* agents of leptospirosis, *Bartonella spp.* agents of bartonellosis and *Orientia*
17 *tsutsugamushi*, the agent of scrub typhus. *Bartonella sp.* and *Orientia tsutsugamushi* are
18 arthropod- borne agents, whereas *Leptospira spp.* are indirectly transmitted via contact with
19 water or soils contaminated by urine of infected rodents. From the three protozoans, only
20 *Toxoplasma gondii* can infect humans whereas *Trypanosoma spp.* and *Babesia spp.* infect
21 livestock and more rarely humans. Altogether 1,716 rodents (10%) have been found positive
22 for at least one pathogen. Microparasite richness was defined as the number of pathogen
23 species for which each rodent species was found positive.

1 We used information on main habitats of rodents following Jittapalapong *et al.* (4,5), Ivanova
2 *et al.* (6) and Suntsov *et al.* (7): forests, dry lands near forests, non-flooded lands, paddy fields
3 and irrigated/flooded agricultural lands, houses.

4

5 **Results**

6 Total host sample size varies widely among rodent species, from 5683 *Rattus tanezumi* to 9
7 *Mus musculus*. This last species was removed from the analysis due to its low sample size.
8 This great variability can be explained by the variability in abundance of each species but also
9 by unequal sampling among habitats. All rodent species have not been screened for all
10 selected microparasites, with number of pathogens investigated varying according among
11 rodent species (Table 1). The detection of microparasites then depends on both the pathogens'
12 screening effort (number of pathogens tested) and the host sampling size (number of
13 individual hosts trapped and tested for a given microparasite).

14 Using multiple regression analysis between microparasite richness and host sample size and
15 pathogens' screening effort as independent variables, we found that microparasite species
16 richness was positively related to both independent variables ($P < 0.0001$, host sample size
17 being log transformed). By investigating the residual variations among rodent hosts (Fig.1B),
18 we show that several rodent species harboured more pathogens than that was expected by the
19 regression model (i.e. positive residual values), particularly *Rattus adamanensis*, *Bandicota*
20 *savilei*, *R. argentiventer* and *R. norvegicus*. Two species appeared to harbour less pathogen
21 species than expected by the regression model (i.e. negative residuals values): *Mus cervicolor*
22 and *M. caroli*.

23 Similar reasoning on habitats suggests that higher pathogen richness than expected from
24 correlation with sampling effort (i.e. positive residual values) are found in non-flooded lands,
25 forests and paddy fields.

1

2 **Conclusions**

3 We have developed here a simple method for prioritizing/targeting rodents that are best
4 carriers of rodent-borne diseases, in a sense that they harbour more pathogen species than
5 expected on the basis of the relationship between pathogen richness and sampling effort.

6 Controlling sampling effort in comparative analysis is usual as pathogen/parasite richness is
7 highly correlated with the efforts done to sample their hosts. However, host sample size is
8 also often positively correlated with some host features such as host density or host
9 geographical range (8). A host living in high density and a wide geographical range is more
10 sampled than a host living in low density and on restricted range. As parasite/pathogen
11 richness is found correlated with host density and/or host geographical range (9-10), there are
12 potential confounding effects between these host features and the sampling effort to detect
13 these parasites/pathogens. Here, rather to determine the potential determinants of pathogen
14 richness in rodents, which is a difficult task due to the lack of knowledge on many life traits
15 of the species living in Southeast Asia, we used the residual variations of the pathogen
16 richness / sampling effort relationship. Then, high residuals values mean high pathogen
17 richness whatever the explaining factor. Interestingly, our results suggest some rodent species
18 that are not commonly investigated to target for pathogen screening or surveillance such as *R.*
19 *adamanensis* or *B. savilei*.

20 The second result suggest that non-flooded lands and forests should be more taken into
21 caution, whereas much surveys focused on paddy rice fields and households, although for
22 obvious reasons. There is growing empirical evidence that some ecosystems are prone to alter
23 or improve parasite transmissions (11-13). The recent study of Duffy et al. (14) emphasizes
24 that ecosystems with high productivity (and then high host densities) could select hosts for
25 being more resistant to infections to limit epidemics and parasite transmission but also less

1 fecund (due to trade –offs between reproduction and immunity) compare to host living in
2 ecosystems with low productivity. Assuming the reality of differences in pathogen richness
3 between rodent species and the various habitats, our results call for future studies in Asian
4 ecosystems to improve the processes prone to explain such patterns.
5 Finally the simple method developed here based on known research effort in pathogens’
6 screening of wildlife can present some interest in surveillance prioritization (15) by allocating
7 wildlife surveillance effort to specific rodent species in specific habitats.

8

9 **Acknowledgements**

10 This study is supported by the French ANR Biodiversity ANR 07 BDIV 012, project
11 CERoPath, “Community Ecology of Rodents and their Pathogens in a changing
12 environment”.

13

14 **References**

- 15 1. Sejvar J, Tangkanakul W, Ratanasang P, Dowell SF, Sangjun N, Bragg S, et al. An
16 outbreak of leptospirosis, Thailand--the importance of the laboratory. *Southeast Asian J*
17 *Trop Med Public Health* 2005;36:289-95.
- 18 2. Meerburg BG, Singleton GR, Kijlstra A. Rodent-borne diseases and their risks for public
19 health. *Critic Rev Microb* 2009:1–50.
- 20 3. Sodhi NS, Koh LP, Brook BW, Ng PKL. Southeast Asian biodiversity: an impending
21 disorder. *TREE* 2004;9:654-60.
- 22 4. Jittapalapong S, Inpankaew T, Sarataphan N, Herbreteau V, Hugot, JP, Morand S et al.
23 Molecular detection of divergent trypanosomes among rodents of Thailand. *Infection*
24 *Gen Evol* 2008;8: 445-49.

- 1 5. Jittapalapong S, Sarataphan N, Maruyama S, Hugot JP, Morand S, Herbreteau V.
2 Seroprevalence of *Toxoplasma gondii* infections of rodents in Thailand. Vector-Borne
3 Zoonot Dis 2011;11:231-37.
- 4 6. Ivanova S, Herbreteau V, Blasdell K, Chaval Y, Buchy P, Guillard B et al. *Leptospira* and
5 rodents in Cambodia: environmental determinants of infection. Am J Trop Med Hyg
6 2012; (in press).
- 7 7. Suntsov VV, Ly TVH, Adler GH. Distribution of rodents along a gradient of disturbance
8 on the Tay Nguyen Plateau of southern Vietnam. Mammalia 2003 67:379-83.
- 9 8. Guégan JF, Kennedy CR. Parasite richness/ sampling effort/host range: the fancy three-
10 piece jigsaw puzzle. Parasitol Today 1996;12:367-70.
- 11 9. Poulin R, Morand S. The diversity of parasites. Quart Rev Biol 2000;75: 277-93.
- 12 10. Bordes S, Morand S. The impact of multiple infections on wild animal hosts: a review.
13 Infect Ecol Epidemiol 2011;1:7346.
- 14 11. Cottontail VM, Wellinghausen N, Kalko EKV. Habitat fragmentation and haemoparasites
15 in the common fruit bat, *Artibeus jamaicensis* (Phyllostomidae) in a tropical lowland
16 forest in Panamá. Parasitology 2009;136:1133-45.
- 17 12. Gillespie TR, Chapman, CA. Forest fragmentation, the decline of an endangered primate
18 and changes in host parasite interactions relative to an unfragmented forest. Am J Primatol
19 2008;70:222-30.
- 20 13. Rohr J, Schotthoefer, AM, Raffel, TR, Carrick, HJ, Halstead N, Hoverman J et al.
21 Agrochemicals increase trematode infections in a declining amphibian species. Nature.
22 2008 455:1235- 39.
- 23 14. Duffy MA, Ochs JH, Penczykowski RM, Civitello DJ, Klausmeir CA, Hall SR.
24 Ecological context influences epidemic size and parasite driven evolution. Science
25 2012;335:1636-38.

1 15. McIntyre KM, Hawkes I, Waret-Szkuta A, Morand S, Baylis M. The H-Index as a
2 quantitative indicator of the relative impact of human diseases. PLoS ONE. 2011;6:
3 e19558.

4

1

2 **Figure captions**

3

4 Figure 1. Residual values in microparasite richness, corrected for rodent sample size and
5 pathogens' screening effort, positively sorted according (A) to rodent species and (B) to
6 habitats. High positive values of residuals in microparasite richness indicate higher
7 species diversity of microparasites than expected by the regression modelling and can
8 help at identifying "good" rodent carriers of pathogens or risky habitats, at least in term
9 of high diversity of pathogens than can be encountered herein.

10

1 Table 1. Survey of infection by microparasites (viruses, bacteria, protozoans) of rodent
 2 species in Thailand, with number of positive individuals and number of investigated
 3 individuals between brackets (see references in supplementary materials)

Species	<i>Leptospira</i> <i>spp.</i>	<i>Orientia</i> <i>spp.</i>	<i>Bartonella</i> <i>Spp.</i>	Hanta virus	Herpes virus	LCM virus	Rabies virus	<i>Toxoplasma</i> <i>gondii</i>	<i>Trypanosoma</i> <i>spp.</i>	<i>Babesia</i> <i>spp.</i>
<i>Bandicota indica</i>	102 (1006)	101 (755)	12 (167)	60 (932)	3 (164)	20 (166)	0 (276)	1 (37)	11 (192)	17 (30)
<i>Bandicota savilei</i>	12 (464)	52 (189)	2 (33)	3 (197)	2 (12)	5 (14)	0 (17)	0 (11)	13 (64)	
<i>Berylmys</i> <i>berdmorei</i>	0 (6)	2 (9)	5 (20)	0 (12)			0 (3)	0 (4)	1 (23)	
<i>Berylmys bowersi</i>		0 (9)		0 (5)				0 (5)	0 (3)	
<i>Leopoldamys</i> <i>edwardsi</i>							0 (23)	2 (16)	4 (10)	
<i>Maxomys surifer</i>		0 (19)	8 (33)	0 (6)			1 (150)	2 (38)	6 (22)	
<i>Mus caroli</i>	0 (6)	0 (69)	0 (5)	0 (67)	0 (6)	0 (3)				
<i>Mus cervicolor</i>	0 (12)	0 (17)	1 (2)	0 (85)				0 (1)	0 (13)	
<i>Mus musculus</i>	0 (4)			0 (2)		0 (3)				
<i>Mus cookii</i>				0 (40)				0 (1)	0 (17)	
<i>Niviventer</i> <i>fulvescens</i>		1 (4)	0 (8)	0 (10)			0 (11)	0 (13)	0 (2)	
<i>Rattus</i> <i>andamanensis</i>		1 (12)					0 (1)	0 (7)	1 (3)	
<i>Rattus</i> <i>argentiventer</i>	6 (102)	5 (23)		0 (107)	15 (31)	1 (9)			4 (62)	
<i>Rattus exulans</i>	48 (1242)	20 (465)	1 (71)	24 (667)	0 (3)	3 (47)	0 (1)	1 (79)	22 (266)	0 (17)
<i>Rattus losea</i>	6 (86)	82 (638)	2 (24)	1 (119)	0 (10)	0 (2)	0 (4)	0 (1)	3 (12)	
<i>Rattus norvegicus</i>	179 (860)	11 (36)		26 (309)	22 (48)	17 (54)	0 (1)	0 (34)	1 (14)	
<i>Rattus tanezumi</i>	107 (1858)	542 (2284)	7 (73)	21 (900)	5 (74)	3 (69)	0 (139)	11 (256)	7 (143)	
<i>Rattus tiomanicus</i>				0 (97)	26 (105)			0 (2)		

4


5

1


3 Figure 1

5

A



B



1 **Supplemental file**
2 **References for the data on microparasites in rodents**

3
4
5

5 **Hantaviruses**

- 6 Blasdel K, Cosson JF, Chaval Y, Herbreteau V, Douangboupha B, Jittapalapong S,
7 Lundqvist A, Hugot JP, Morand S, Buchy, P Rodent-borne Hantaviruses in
8 Cambodia, Laos PDR and Thailand. *EcoHealth* 2012 [ahead of print].
- 9 Elwell MR, Ward GS, Tingpalapong M, et al. Serologic evidence of Hantaan-like virus in
10 rodents and man in Thailand. *Southeast Asian J Trop Med Pub Health* 1985;
11 6:349-354.
- 12 LeDuc JW, Smith GA, Childs JE, et al. Global survey of antibody to Hantaan-related
13 viruses among peridomestic rodents. *Bulletin of the World Health Organization*
14 1986;64:139-144.
- 15 Nitatpattana N, Chauvancy G, Dardaine J, et al. Serological study of hantavirus in the
16 rodent population of Nakhon Pathom and Nakhon Ratchasima provinces
17 Thailand. *Southeast Asian J Trop Med Pub Health* 2000a; 31:277-282.
- 18 Nitatpattana N, Chauvancy G, Jumronsawat K, et al. Preliminary study on potential
19 circulation of arenaviruses in the rodent population of Nakhon Pathom Province,
20 Thailand and their medical importance in an evolving environment. *Southeast*
21 *Asian J Trop Med Pub Health* 2000b 31:62-65.
- 22 Nitatpattana N, Henrich T, Palabodeewat S, et al. Hantaan virus antibody prevalence in
23 rodent populations of several provinces of north-eastern Thailand. *Trop Med Int*
24 *Health* 2002; 7:1-6
- 25 Pattamadilok S, Lee BH Kumperasart, K, .al. Geographical distribution of hantaviruses in
26 Thailand and potential human health significance of Thailand virus. *Am J Trop*
27 *Med Hyg* 2006; 75:994-1002.

1 Tantivanich S, Ayuthaya PI, Usawattanakul W, et al. Hantaanvirus among urban rats `

2 from a slum area in Bangkok. Southeast Asian J Trop Med Public Health 1992;

3 23:504- 509

4

5 **Arenaviruses**

6 Nitatpattana N, Chauvancy G, Jumronsawat K, et al. Preliminary study on potential

7 circulation of arenaviruses in the rodent population of Nakhon Pathom Province,

8 Thailand and their medical importance in an evolving environment. Southeast

9 Asian J Trop Med Pub Health 2000b 31:62-65.

10

11 **Rabies Virus**

12 Brown JL, Tingpalapong M, Andrews WK. Serological survey of feral rodents in Thailand

13 for evidence of rabies virus infection J Wild Dis 1979; 15:601-606.

14

15 **Hepatitis E Virus**

16 Schmidt-Chanasit J et al. Evidence for Hepatitis E Virus infection among rodents in

17 Thailand. In: MEEGID VIII, 2006; Bangkok

18

19 **Leptospirosis**

20 Bunnag T, Potha U, Thirachandra S, Impand P. Leptospirosis in man and rodents in

21 North and Northeast Thailand. South Asian J Trop Med Pub Health 1983; 14:481-

22 487.

23 Douchawee G, Phulsuksombat D, Naigowi P, et al. Survey of leptospirosis of small

24 mammals in Thailand. Southeast Asian J Trop Med Pub Health 2005; 36:1516-

25 1522.

- 1 Imvithaya A, Warachit P, Naigowit P, et al. Survey of Host Reservoir of Rodent-Borne
2 Kositanont U, Naigowit P, Imvithaya A, et al. Prevalence of antibodies to *Leptospira*
3 serovars in rodents and shrews trapped in low and high endemic areas in
4 Thailand. J Med Ass Thailand 2003; 86:136-142.
- 5 Phulsuksombati D, Sangjun N, Khoprasert Y, et al. Leptospire in rodent, northeastern
6 region 1999-2000. 2001; 10:516-525.
- 7 Phulsuksombati D, Tangkanakul W, Sangjun N, et al. Isolation of leptospire from
8 rodents in Nakhonratchasima province. J Health Sc 1999; 8:360-369.
- 9 Wangroongsarb P, Petkanchanapong W, Yasaeng S, et al. Survey of leptospirosis among
10 rodents in epidemic areas of Thailand. J Trop Med Parasitol 2002; 25:55-58.
- 11
- 12 ***Bartonella spp.***
- 13 Castle KT, Kosoy M, Lerdthusnee K, et al. Prevalence and diversity of *Bartonella* in
14 rodents of northern Thailand: a comparison with *Bartonella* in rodents from
15 southern China. Am J Trop Med Hyg 2004; 70: 429-433.
- 16 Saisongkorh W, Wootta W, Sawanpanyalert P, et al. "*Candidatus Bartonella*
17 *thailandensis*": A new genotype of *Bartonella* identified from rodents. Vet Microb
18 2009
- 19
- 20 ***Orientia tsutsugamushi***
- 21 Coleman RE, Taweesak M, Linthicum KT, et al. Occurrence of *Orientia tsutsugamushi* in
22 small mammals from Thailand. Am J Trop Med Hyg 2003; 69:519-524.
- 23 Strickman D, Smith CD, Kevin D, et al. Pathology of *Rickettsia tsutsugamushi* infection in
24 *Bandicota savilei*, a natural host in Thailand. Am J Trop Med Hyg 1994; 51:416-
25 423.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Trypanosoma spp.

Dantrakool A, Somboon P, Hashimoto T, et al. Identification of a New Type of *Babesia* Species in Wild Rats (*Bandicota indica*) in Chiang Mai Province, Thailand. J Clin Microb 2004; 42:850-854.

Jittapalapong S, Inpankaew T, Sarataphan N, et al. Molecular detection of divergent trypanosomes among rodents of Thailand. Infection, Genetics and Evolution 2008; 8:445-449

Milocco C, Kamyinkird K, Desquesnes M, Jittapalapong S, Herbreteau V, Chaval Y, Douangboupha B, Morand S (2012) Molecular demonstration of *Trypanosoma evansi* and *Trypanosoma lewisi* DNA in wild rodents from Cambodia, Lao PDR and Thailand. Transboundary and Emerging Diseases (in press)

Babesia spp.

Dantrakool A, Somboon P, Hashimoto T, et al. (2004) Identification of a New Type of *Babesia* Species in Wild Rats (*Bandicota indica*) in Chiang Mai Province, Thailand. J Clin Microb 2004; 42:850-854.

Toxoplasma gondii

Jittapalapong S, Sarataphan N, Maruyama S, Hugot JP, Morand S, Herbreteau V (2011) Seroprevalence of *Toxoplasma gondii* infections of rodents in Thailand. Vector-Borne Zoonot Dis 11:231-237.