



Statistics of the French purse seine fishing fleet targeting tropical tunas in the Indian Ocean (1981-2011)

Emmanuel Chassot, Laurent Floch, Patrice Dewals, Isabelle Terrier, Pierre Chavance

► To cite this version:

Emmanuel Chassot, Laurent Floch, Patrice Dewals, Isabelle Terrier, Pierre Chavance. Statistics of the French purse seine fishing fleet targeting tropical tunas in the Indian Ocean (1981-2011). IOTC Working Party on Tropical Tunas, Oct 2012, Grand-Baie, Mauritius. 29 p. multigr. ird-00780451

HAL Id: ird-00780451

<https://ird.hal.science/ird-00780451>

Submitted on 24 Jan 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Statistics of the French purse seine fishing fleet targeting tropical tunas in the Indian Ocean (1981-2011)

Chassot E.^a, L. Floch^{a,*}, P. Dewals^b, I. Terrier^a, P. Chavance^a

^a*Institut de Recherche pour le Développement, UMR 212 EME (IRD/Ifremer/Université de Montpellier 2),
Avenue Jean Monnet, BP 171, 34 203 Sète Cedex, FRANCE*

^b*Institut de Recherche pour le Développement, Seychelles Fishing Authority, BP 570, Victoria,
SEYCHELLES*

Abstract

In 2011, the French purse seine fishing fleet of the Indian Ocean was composed of 8 large size vessels that represented a total carrying capacity of about 8,000 t. Catches reached a total of 43,000 t and were composed of 50%, 42%, and 8% of yellowfin, skipjack, and bigeye, respectively. After a period of increase during 2006-2008, the fishing effort of the fleet has been decreasing to reach a minimum of 1,800 searching days in 2011. The decrease in effort was associated with a contraction of the fleet fishing grounds in the recent years and mainly characterized by a strong decrease in the number of sets made on free-swimming schools; a total of 1,800 fishing sets being made in 2011 compared to about 4,200 y⁻¹ in the mid-2000s. Hence, the percentage of sets made on FAD-associated schools steadily increased since 2004 to reach more than 65% in 2010-2011, FAD-fishing resulting in 75% of the total catch of the French purse seine fishing fleet in 2011. Species-specific catch rates (in t per searching day) on FAD-associated schools reveal increasing trends over 1981-2011 with high interannual variability while no clear trend is apparent in the time series of catch rates on free-swimming schools. The year 2011 is characterized by a large proportion of yellowfin in the catch on FAD-associated schools that might indicate a good recruitment. For both fishing modes, the mean weight in the catch of the 3 tropical tunas has decreased since the mid-2000s with values in 2011 equal to 7.1 kg, 2.4 kg, and 5.4 kg for yellowfin, skipjack, and bigeye, respectively.

Keywords: fish aggregating device, free-swimming school, *Katsuwonus pelamis*, purse seine fishing, *Thunnus albacares*, *Thunnus obesus*

*Corresponding author

Email address: Laurent.Floch@ird.fr (L. Floch)

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 2 |
| 2 | Fishing capacity and effort | 3 |
| 2.1 | Fishing fleet | 3 |
| 2.2 | Carrying capacity | 3 |
| 2.3 | Fishing and searching days | 4 |
| 2.4 | Fishing grounds | 5 |
| 2.5 | Fishing activities | 6 |
| 3 | Fisheries production | 9 |
| 3.1 | Catch levels | 9 |
| 3.2 | Spatial distribution of the catch | 9 |
| 3.3 | Catch rates | 11 |
| 3.4 | Size structure of the catch | 16 |
| 3.5 | Mean weight in the catch | 17 |
| 4 | References | 21 |
| 5 | Appendix tables | 22 |

1. Introduction

Statistical data for the French purse seine fishing fleet have been collected by the “Institut de Recherche pour le Développement” (IRD; ex-ORSTOM) since the arrival of the first purse seiners in the early 1980s. French purse seiners target yellowfin (*Thunnus albacares*), skipjack (*Katsuwonus pelamis*), and bigeye tuna (*Thunnus obesus*) through two major fishing modes that result in different species and size composition of the catch: log/FAD-associated (FAD) and free-swimming schools (FSC). The acronym “FAD”, which stands for drifting fish aggregating device, is used here to describe any type of floating object used for increasing tuna catchability. This includes natural objects (e.g. logs, palm branches) and anthropogenic floating objects, such as man-made bamboo rafts equipped with radio-range beacons, satellite transmitters or scanning sonars. Fishing sets made on whales were classified as free-swimming school sets whereas sets made on whale sharks (*Rhincodon typus*) were classified as FAD sets (Pallarés and Hallier 1997). The fleet activities are described through a suite of fisheries indicators that provide information on fishing effort, catch, catch rates, size structure and mean weights for the 3 principal market tropical tunas, with a particular focus on the year 2011.

2. Fishing capacity and effort

2.1. Fishing fleet

The number of vessels of the French purse seine fishing fleet in the Indian Ocean rapidly increased from 2 in 1981 to a maximum of 26 during 1984-1985 with the arrival of vessels from the Atlantic Ocean. Since then, the number of vessels steadily decreased to reach a minimum of 8 in 2010-2011 (Fig. 1 and Table 1). The size of the vessels progressively increased in the French purse seine fishery over the last 30 years. The number of small-size vessels (capacity < 600 t) decreased throughout the 1980s and 1990s to become 0 in the mid-2000s while medium-size vessels (capacity between 601-800 t) disappeared from the fishery in the late 2000s (Fig. 1).




Figure 1: Fishing capacity of the French purse seine fleet in the Indian Ocean. Annual changes in the number of purse seiners by size category (barplots) and total carrying capacity (solid line with circles) during 1981-2011. Capacity was weighted by the vessel-specific proportion of the year at sea (in months). The vessel size category (t) was computed as 0.7 times the capacity expressed in m^3

2.2. Carrying capacity

The total carrying capacity expressed in tonnage strongly increased from the arrival of the first purse seiner in 1981 to about 14,000 t in the mid-1980s. Despite the steady decrease in the total number of vessels, the capacity remained stable through the 1990s and 2000s




Figure 2: Changes in mean vessel carrying capacity for the French purse seine fishing fleet in the Indian Ocean, 1981-2011. The vessel-specific carrying capacity was weighted by the relative annual catch of each purse seiner. Solid line indicates mean linear regression

and increased to a maximum of 17,000 t in 2007 (Fig. 1). Indeed, the mean vessel-specific carrying capacity of the French purse seiners increased from about 600 t in the early 1980s to more than 1,000 t from the mid-2000s (Fig. 2; slope = $+13 \text{ t}^{-1}$, p -value < 0.001). In the recent years, the capacity of the French fleet decreased to reach about 8,000 t during 2010-2011. Meanwhile, the capacity of the French overseas territories increased from less than 500 t in 2006 to more than 5,000 t in 2011 due to the construction of new purse seiners (Chassot et al. 2012).

2.3. Fishing and searching days

Fishing effort expressed in searching time (days) was computed by subtracting the time spent setting the gear from the fishing time. The time spent setting the gear was estimated by regressions linking duration and size of sets, from at-sea measurements made by scientific observers. The total number of fishing and searching days showed similar patterns over 1981-2011 with a major increase in the mid-1980s and a steady decrease until 2005 thereafter (Fig. 3). An increase in effort to a value of almost 5,000 fishing days was then observed due to the increase in the number of purse seiners from 14 in 2003 to 17 during 2006-2008. Since then, the nominal fishing effort strongly decreased and

reached in 2011 values of about 2,110 and 1,800 fishing days and searching days, respectively.




Figure 3: Changes in nominal effort over time. Annual total number of fishing and searching days for the French purse seine fishing fleet in the Indian Ocean during 1981-2011

2.4. Fishing grounds

The major fishing grounds of the French purse seine fishing fleet in 2011 were located in the north of the Mozambique Channel, west of the Seychelles, and along the Somali EEZ boundary (Fig. 4). A few fishing activities took place south of the Seychelles at longitude between 55° and 60°E and in the east of the Chagos EEZ.

After a major expansion of the fishing grounds during the early 1980s, the spatial extent of the fishery steadily increased over time from a total of about 400 squares in the mid-1980s to a mean value of 490 squares in the late 2000s (Fig. 5). The peak observed in 1998 corresponded to a particular climatic event that has been described elsewhere (Murtugudde 1999, Murtugudde et al. 2000) and led the purse seiners to explore the eastern part of the Indian ocean for fishing (Ménard et al. 2006). The different selection criteria, i.e. minimum catch, effort, or sets, did not modify the temporal patterns in the spatial extent of the fishery (Fig. 5).




Figure 4: Fishing grounds. Spatial distribution of fishing effort (in searching days) of the French purse seine fishing fleet in 2011

2.5. Fishing activities

The total annual number of fishing sets made by the French purse seine fleet showed a general pattern consistent with the annual variations in carrying capacity and fishing effort of the fleet (Fig. 6 and Table 4). The fleet showed a strong decrease in the number of sets from more than 4,300 in the mid-2000s to less than 2,000 in 2010-2011. The total number of sets showed a pattern very similar to the annual changes in the number of sets made on free-swimming schools (Fig. 6). The annual number of sets made on FAD-associated schools was more stable over time (around 1,700 y^{-1}) and in opposite phase compared to sets made on free-swimming schools. The percentage of FAD-associated over free-swimming schools varied around a mean of 47% (SD = 10%) with FAD-associated fishing predominating from the mid-1990s to the early 2000s while sets made on free-swimming schools were more frequent during 2003-2008. The percentage of sets made on FAD-associated schools steadily increased from 37% in 2004 to reach more than 65% in 2010-2011.

The percentage of successful sets (i.e. positive tuna catch) made on FAD-associated schools has been high over the period 1981-2011, the mean annual value varying between a minimum of 85% in 1984 and a maximum of 95% in 1991. After a decrease during the development of the fishery, the percentage of successful sets on free-swimming schools steadily increased from a mean value of 37% in 1986 to an average of 54% during 2007-2011 (slope = +0.5% y^{-1} , p -value < 0.001). By contrast, the high success of sets on FAD-associated schools does




Figure 5: Changes in spatial extent of the fishery over time. Annual number of 1-degree squares explored by the French purse seine fishing fleet during 1981-2011




Figure 6: Fishing activities. Annual number of fishing sets in the French purse seine fishery on FAD-associated and free-swimming schools during 1981-2011. Line with solid circles indicates the percentage of sets made on FAD-associated schools over free-swimming schools. Grey solid line indicates the 50% value

not show any clear temporal pattern over the period. In the recent years, the success rate for FAD-associated schools seems to show an increasing trend that could be related to the improvement and increasing use of lateral echo sounders.

3. Fisheries production

3.1. Catch levels

The French purse seine fishery showed strong interannual variations in the catch over 1987-2008 with a minimum of 56,000 t in 1998 and a maximum of 108,000 t during 2003-2004 (Fig. 7). Since then, the total catch of the fleet strongly declined in relation with the decrease in the number of vessels and reached 43,000 t in 2011. Catches on FAD-associated schools represented on average 60% (SD = 8%) of the total fleet catch over 1981-2011. Following the large catch of tunas on free-swimming schools during 2003-2005, the percentage of FAD-associated catch steadily increased (slope = +4.3%, p -value < 0.001) to reach more than 70% during 2009-2011. Catches on FAD-associated schools appeared more stable than catches on free-swimming schools during 1981-2011 (Fig. 8). They were largely dominated by skipjack that represented an annual mean of 62% (SD = 6.5%) of all species caught, varying from 53% in 1997 to more than 75% in 1991 (Fig. 8a). Catches of bigeye, re-estimated from multispecies sampling operations conducting at unloading (Pallarés and Hallier 1997), varied between 3.7% and 13% of the total of catch on FAD-associated schools during 1981-2011. Catches of yellowfin on FAD-associated schools were stable over the period representing on average 27% (SD = 5.2%) of the catch.

Catches made on free-swimming schools showed strong interannual variations in the last two decades from a maximum of 50,000 t in 2004 to a minimum of 8,000 t in 2011 (Fig. 8b). The changes over time in free-swimming schools catches drove the trends in the total catch of the fishery. The catches were largely dominated by yellowfin that represented an annual average of 74% (SD = 9%) of the FSC catch during 1981-2010, and a similar value of 74% in 2011. However, while free-swimming schools represented more than 60% of the total catch of yellowfin during 1981-2010, it only represented 38% of the total yellowfin catch in 2011. The predominance of yellowfin in the French purse seine catch from FAD-associated fishing (> 50% of the catch) had only been observed during 1997-1999 during the strong El-Niño event. In the recent period, this pattern stems from the strong decrease in the number of sets made on free-swimming schools which might reveal a change in purse seine fishing strategy due to a decrease in tuna free-swimming schools in conjunction with the Somali piracy threat (Chassot et al. 2010), the implementation of the BIOT marine protected area that decreased purse seine fishing grounds as well as to the high sale price of skipjack (Table 4). In addition, the proportion of yellowfin in FAD-associated catch reached more than 40% in 2011 while it represented an average of 28% (SD = 5%) during 1982-2010 (see below).

3.2. Spatial distribution of the catch

The French purse seine fleet fishing grounds in 2011 were concentrated in the north of the Mozambique Channel and the Somali area where FAD-fishing predominated while




Figure 7: Total fishery production. Catch by species of the French purse seine fishing fleet during 1981-2011




Figure 8: Fishery production by major fishing mode. Catch by species of the French purse seine fishing fleet on (a) FAD-associated and (b) free-swimming schools during 1981-2011

fishing on free-swimming schools mainly occurred around and south-east of the Seychelles islands, and in the Mozambique to a lesser extent (Fig. 9). As compared to previous years (2006-2010), the composition of the catch on FAD-associated schools in the Somali area was described by an important part of yellowfin that represented more than 50% of the catch in most of the 1x1 degree squares west of 55°E and north of 10°S (Figs. 11-12). This might result from a good recruitment of yellowfin in 2011 as suggested by the size structure of the catch (see Section 3.4).




Figure 9: Spatial distribution of tuna catches of the French purse seine fishing fleet in 2011

Besides, catches on free-swimming schools in 2011 were predominated by skipjack in the Mozambique Channel (Fig. 13), as generally observed in this area, while the remaining of the catch was scattered along latitude 5°S and predominated by yellowfin, with a few squares characterized by 15-20% of bigeye (Fig. 13).

3.3. Catch rates

Raw catch rates expressed in tonnes per searching day ($t d^{-1}$) for the principal market tunas exhibit an increasing trend over 1981-2011, in particular from the early 1980s to the mid-1990s (Fig. 15). While skipjack and yellowfin are mainly caught in different fishing modes, their respective CPUEs were in the same order of magnitude, i.e. a mean value of 10.3 and 10 t per searching day for yellowfin and skipjack, respectively. Also, the CPUE time series showed very similar trends over time, with the notable exception of 2003-2005




Figure 10: Spatial distribution of tuna catches of the French purse seine fishing fleet in 2006-2010




Figure 11: Spatial distribution of tuna catches of the French purse seine fishing fleet made on FAD-associated schools in 2011




Figure 12: Spatial distribution of tuna catches of the French purse seine fishing fleet made on FAD-associated schools in 2006-2010




Figure 13: Spatial distribution of tuna catches of the French purse seine fishing fleet made on free-swimming schools in 2011




Figure 14: Spatial distribution of tuna catches of the French purse seine fishing fleet made on free-swimming schools in 2006-2010

described by abnormally high catch rates for yellowfin on free-swimming schools (Fig. 16b). Raw catch rates of bigeye also showed an increasing trend during 1981-2011 (slope = +4.4%, p -value < 0.001) with an average of 1.88 t d^{-1} (SD = 0.4 t d^{-1}).

Skipjack catch rates predominated on FAD-associated schools during 1981-2011 and increased quickly through the 1980s while stabilising around $10 \text{ t searching d}^{-1}$ (SD = 2.3 t d^{-1}) since the early 1990s (Fig. 16a). Yellowfin catch rates have shown an increasing trend since the early 1980s with some interannual variability. The maximum catch rates for yellowfin on FAD-associated schools have been observed in 2011, i.e. an annual value of 7.3 t d^{-1} . Bigeye raw catch rates on FAD-associated schools have remained stable around a mean value of 1.4 t d^{-1} since the late 1990s (Fig. 16a).

Yellowfin catch rates predominated for sets made on free-swimming schools and remained quite stable over time (mean of 7 t d^{-1} during 1984-2011), with abnormally high values during 2003-2005 > 13.5 t d^{-1} (Fig. 16b). In 2011, the catch rate of yellowfin in free-swimming schools was less than 4.5 t d^{-1} . Catch rates for skipjack have shown high interannual variability during 1986-2011 (mean = 2 t d^{-1} , SD = 1 t d^{-1}) and a decreasing trend in the recent years, with catch rates < 1 t d^{-1} during 2009-2011. Catch rates for bigeye in free-swimming schools have shown a shift from a mean value around 0.3 t d^{-1} dur-




Figure 15: Annual catch rates (in t per searching day) of the French purse seine fishing fleet in the Indian Ocean during 1981-2011

ing the mid-1980s to the mid-1990s to a mean annual value of 0.6 t d^{-1} throughout the 2000s.




Figure 16: Annual catch rates (in t per searching day) of the French purse seine fishing fleet on (a) FAD-associated and (b) free-swimming schools in the Indian Ocean during 1981-2011

3.4. Size structure of the catch

The size structure of the catch (expressed in number) in 2011 appeared similar for the 3 tuna species to the average year representing the period 2006-2010, with the exception of skipjack caught on free-swimming schools that were smaller (Fig. 17). The sizes of yellowfin caught on FAD-associated schools were described by 2 modes for small (46 cm) and intermediate sizes (56 cm) while larger individuals ($> 80 \text{ cm}$) were also caught but more seldomly. The year 2011 was characterized by a high peak of small yellowfin (size range of 42-48 cm fork length) which might indicate a good recruitment. Bigeye caught on FAD-associated schools in 2011 were only juvenile fishes of median size of 52 cm (sd = 9 cm). Skipjack showed a unique size mode described by a median of 46 cm (min = 31 cm and max = 69 cm) similar to 2006-2010 during which the median size of skipjack was 47 cm.

Similarly, the biomass of fish caught by size class showed overall similar patterns between 2011 and 2006-2010 except for skipjack caught smaller on free-swimming schools in 2011 (Fig. 18). For yellowfin caught on FAD-associated schools, the year 2011 was described by larger biomass of fishes comprised between 60-100 cm. In relation with the decrease in activities on free-swimming schools, the biomass of large yellowfin and bigeye ($> 100 \text{ cm}$) caught was lower in 2011 than for the average year 2006-2010. The biomass of skipjack 44-50 cm long caught on FAD-associated schools, which represent the bulk of skipjack catch, was very similar in 2011 than during 2006-2010 while there was a strong decrease in

skipjack catch on free-swimming schools (Table 5-7).

3.5. Mean weight in the catch

Time series of the mean weight in the catch for the principal market tropical tunas were computed as the species-specific annual biomass over the total number of fishes caught. The sampling design used to collect size-frequency data aboard French purse seiners throughout the 1980s varied over time (Pianet 1999) therefore the mean weight was computed from 1991 for data consistency. The mean weight highly differed between fishing modes and showed strong interannual variations during 1991-2011 (Fig. 19). The mean weight of yellowfin caught on free-swimming schools decreased from more than 37 kg in 1991 to a minimum of about 15 kg in 1998-1999, before progressively increasing thereafter to reach about 38 kg in 2004-2005. Since then, the mean weight decreased to about 28 kg during 2009-2011 (Fig. 19a). It is noteworthy that the yellowfin mean weight showed an increase from 25 kg in 2010 to 29 kg in 2011. After an initial decrease from about 10 kg in the early 1990s, the mean weight of yellowfin in the catch made on FAD-associated schools stabilised at around 5.7 kg ($SD = \pm 1.1$ kg) from the mid-1990s to the late 2000s. In 2011, the mean weight of yellowfin caught on FAD-associated schools was less than 5 kg. Overall, the mean weight of yellowfin caught in the French purse seine fleet has shown a strong decrease from about 18 kg during 2003-2005 to about 7 kg in 2011 (slope = -1.4 kg^{-1} ; p -value < 0.001) due to increasing proportion of FAD-fishing in the fishery over the last decade (Fig. 6 – dotted line).

The overall mean weight of skipjack was more stable than yellowfin during 1991-2011 and driven by the mean weight of skipjack caught on FAD-associated schools. It varied between a maximum of 3 in 1992 and a minimum of 2 kg in 2008 for fishes caught on FAD-associated schools (Fig. 19b). After a decrease during 2007-2008, the mean weight remained stable at about 2.3 kg during 2009-2011. The mean weight of skipjack caught on free-swimming schools has shown a strong and significant decrease from more than 4.2 kg in 2003-2004 to 2.4 kg in 2009-2011 (slope = -0.3 kg^{-1} , p -value < 0.001).

The mean weight of bigeye varied around 5.8 kg ($SD \pm 1$ kg) during 1991-2011 for the whole fishery (Fig. 19c). It showed strong interannual variations for individuals caught on free-swimming schools with a pattern very similar to yellowfin tuna (Pearson's $r = 0.77$; p -value < 0.001). The mean weight of bigeye caught on FAD-associated schools showed a significant decreasing trend during 1991-2011 (slope = -0.06 kg^{-1} , p -value < 0.01) with high interannual variations. In 2011, the mean weight of bigeye caught on free-swimming schools and FAD-associated schools was 18.3 kg and 3.8 kg, respectively.

Acknowledgments. We thank ORTHONGEL, Pascal Cauquil, Céline Rodriguez, Julien Barde and all past and current personnel for helpful assistance in data collection and management. We are particularly grateful to Renaud Pianet, Viveca Nordstrøm and Alain Fonteneau for their dedication to the monitoring of tropical tuna fisheries and their support to the Observatoire Thonier. The collaboration with Alicia Delgado de Molina (IEO) and




Figure 17: Size structure of the catch. Size distribution (in numbers) of the species-specific catch for the French purse seine fishing fleet in 2011 (red line) and for an average year representing the period 2006-2010 (black line)




Figure 18: Size structure of the catch. Size distribution (in weight) of the French purse seine fishing fleet in 2011 (red line) and for an average year representing the period 2006-2010 (black line)




Figure 19: Annual time series of mean weight (kg) for (a) yellowfin tuna (YFT), (b) skipjack tuna (SKJ), and (c) bigeye tuna (BET) for each fishing mode during 1991-2011

Cindy Assan (SFA) was instrumental in data processing. This work was financed by the European Data Collection Framework (DCF, Reg 199/2008 and 665/2008) and supported by the Direction des Pêches Maritimes et de l'Aquaculture (DPMA).

4. References

Chassot, E., P. Dewals, L. Floch, V. Lucas, M. Morales-Vargas, and D.M. Kaplan. 2010. Analysis of the effects of Somali piracy on the European tuna purse seine fisheries of the Indian Ocean. Proceedings of the 13th Session of the IOTC Scientific Committee. IOTC-2010-SC-09, 26p.

Chassot, E., Floch, L., P. Dewals, I. Terrier, and P. Chavance. 2012. Statistics of the purse seine fleet of the French overseas territories targeting tropical tunas in the Indian Ocean (2001-2011). Proceedings of the 14th Session of the IOTC Working Party on Tropical Tunas, IOTC-2012-WPTT14-21, 27p.

Ménard, F., F. Marsac, E. Bellier, and B. Cazelles. 2007. Climatic oscillations and tuna catch rates in the Indian Ocean: a wavelet approach to time series analysis. Fisheries Oceanography 16: 95–104.

Murtugude, R. G., S. R. Signorini, J. R. Christian, A. J. Busalacchi, C. R. McClain, and J. Picaut. 1999. Ocean color variability of the tropical Indo-Pacific basin observed by SeaWiFS during 1997–1998. Journal of Geophysical Research 104: 18351–18351.

Murtugude, R. G., J. P. McCreary, and A. J. Busalacchi. 2000. Oceanic processes associated with anomalous events in the Indian Ocean with relevance to 1997-1998. Journal of Geophysical Research 105: 3295–3306.

Pallarés, P & J.P. Hallier. 1997. Analyse du schéma d'échantillonnage multispécifique des thonidés tropicaux (Programme n° 95/37), Technical report, IEO/ORSTOM

Pianet, R. 1999. Évolution du système de collecte et de traitement des données de la pêche thonière des senneurs européens et assimilés de 1981 à 1998. IOTC Working Party on Data Collection and Statistics, September 1999. IOTC Proceedings 2: 74-96.

5. Appendix tables

Table 1: Annual number of purse seiners by size category and total carrying capacity of the French tropical tuna purse seine fishing fleet of the Indian Ocean during 1981-2011. Total carrying capacity (CC) was weighted by the proportion of the year at sea (in months)

| Year | 50-400 | 401-600 | 601-800 | 801-1200 | 1201-2000 | >2000 | Total | CC |
|------|--------|---------|---------|----------|-----------|-------|-------|-------|
| 1981 | 1 | 0 | 0 | 1 | 0 | 0 | 2 | 233 |
| 1982 | 1 | 1 | 0 | 2 | 0 | 0 | 4 | 945 |
| 1983 | 1 | 6 | 0 | 5 | 0 | 0 | 12 | 3907 |
| 1984 | 0 | 11 | 6 | 9 | 0 | 0 | 26 | 14566 |
| 1985 | 0 | 11 | 6 | 9 | 0 | 0 | 26 | 15945 |
| 1986 | 0 | 9 | 5 | 8 | 0 | 0 | 22 | 14526 |
| 1987 | 1 | 6 | 5 | 9 | 0 | 0 | 21 | 13983 |
| 1988 | 1 | 6 | 5 | 9 | 0 | 0 | 21 | 14699 |
| 1989 | 1 | 6 | 5 | 9 | 0 | 0 | 21 | 14285 |
| 1990 | 0 | 7 | 5 | 9 | 0 | 0 | 21 | 12939 |
| 1991 | 0 | 4 | 3 | 9 | 2 | 0 | 18 | 12943 |
| 1992 | 0 | 4 | 2 | 9 | 2 | 0 | 17 | 14220 |
| 1993 | 0 | 4 | 2 | 9 | 2 | 0 | 17 | 14180 |
| 1994 | 0 | 4 | 2 | 9 | 2 | 0 | 17 | 13743 |
| 1995 | 0 | 4 | 2 | 9 | 2 | 0 | 17 | 14199 |
| 1996 | 0 | 3 | 2 | 10 | 2 | 0 | 17 | 13341 |
| 1997 | 0 | 3 | 2 | 10 | 4 | 0 | 19 | 14013 |
| 1998 | 0 | 3 | 2 | 8 | 3 | 0 | 16 | 13074 |
| 1999 | 0 | 2 | 2 | 8 | 3 | 0 | 15 | 12523 |
| 2000 | 1 | 1 | 2 | 8 | 3 | 0 | 15 | 12736 |
| 2001 | 1 | 1 | 2 | 9 | 5 | 0 | 18 | 12261 |
| 2002 | 0 | 1 | 2 | 8 | 5 | 0 | 16 | 14011 |
| 2003 | 0 | 0 | 1 | 8 | 5 | 0 | 14 | 13676 |
| 2004 | 0 | 0 | 2 | 8 | 5 | 0 | 15 | 14090 |
| 2005 | 0 | 0 | 2 | 9 | 5 | 0 | 16 | 13818 |
| 2006 | 0 | 0 | 2 | 10 | 5 | 0 | 17 | 16805 |
| 2007 | 0 | 0 | 2 | 10 | 5 | 0 | 17 | 16949 |
| 2008 | 0 | 0 | 2 | 10 | 5 | 0 | 17 | 16035 |
| 2009 | 0 | 0 | 0 | 9 | 6 | 0 | 15 | 10878 |
| 2010 | 0 | 0 | 0 | 6 | 2 | 0 | 8 | 8275 |
| 2011 | 0 | 0 | 0 | 6 | 2 | 0 | 8 | 8093 |

Table 2: Annual nominal fishing effort of the French purse seine fleet expressed in fishing and searching days during 1981-2011. Searching days was derived from the total time spent at sea corrected for periods of damage, route towards port, and purse seine operation

| Year | Fishing days | Searching days |
|------|--------------|----------------|
| 1981 | 91 | 75 |
| 1982 | 277 | 235 |
| 1983 | 1582 | 1247 |
| 1984 | 5323 | 4310 |
| 1985 | 6308 | 5319 |
| 1986 | 5876 | 4732 |
| 1987 | 5300 | 4240 |
| 1988 | 5683 | 4606 |
| 1989 | 5492 | 4649 |
| 1990 | 5013 | 4202 |
| 1991 | 4146 | 3496 |
| 1992 | 4417 | 3656 |
| 1993 | 4528 | 3853 |
| 1994 | 4490 | 3766 |
| 1995 | 4633 | 3901 |
| 1996 | 4381 | 3734 |
| 1997 | 4427 | 3835 |
| 1998 | 4189 | 3651 |
| 1999 | 3690 | 3152 |
| 2000 | 3742 | 3187 |
| 2001 | 3703 | 3101 |
| 2002 | 3938 | 3242 |
| 2003 | 3488 | 2756 |
| 2004 | 3836 | 3039 |
| 2005 | 3845 | 3096 |
| 2006 | 4714 | 3940 |
| 2007 | 4921 | 4208 |
| 2008 | 4254 | 3589 |
| 2009 | 2692 | 2253 |
| 2010 | 2132 | 1797 |
| 2011 | 2109 | 1796 |

Table 3: Annual number of 1-degree squares explored by the French purse seine fishing fleet during 1981-2011

| Year | TOTAL | #sets>0 | Catch>0 | Effort>1 d | Effort>5 d |
|------|-------|---------|---------|------------|------------|
| 1981 | 73 | 26 | 25 | 18 | |
| 1982 | 133 | 47 | 40 | 53 | 10 |
| 1983 | 257 | 112 | 101 | 137 | 60 |
| 1984 | 574 | 274 | 261 | 342 | 182 |
| 1985 | 496 | 340 | 322 | 384 | 267 |
| 1986 | 406 | 310 | 297 | 333 | 223 |
| 1987 | 416 | 329 | 305 | 323 | 206 |
| 1988 | 393 | 282 | 267 | 300 | 210 |
| 1989 | 442 | 315 | 303 | 355 | 229 |
| 1990 | 444 | 336 | 320 | 353 | 215 |
| 1991 | 411 | 334 | 321 | 332 | 200 |
| 1992 | 404 | 345 | 333 | 330 | 196 |
| 1993 | 414 | 333 | 325 | 328 | 217 |
| 1994 | 438 | 356 | 348 | 363 | 228 |
| 1995 | 445 | 367 | 362 | 371 | 229 |
| 1996 | 522 | 405 | 392 | 409 | 243 |
| 1997 | 524 | 415 | 392 | 422 | 257 |
| 1998 | 755 | 551 | 528 | 556 | 243 |
| 1999 | 611 | 426 | 411 | 418 | 195 |
| 2000 | 498 | 359 | 343 | 361 | 196 |
| 2001 | 439 | 340 | 319 | 344 | 203 |
| 2002 | 546 | 402 | 381 | 405 | 233 |
| 2003 | 410 | 313 | 302 | 293 | 186 |
| 2004 | 470 | 345 | 323 | 330 | 171 |
| 2005 | 441 | 353 | 336 | 337 | 198 |
| 2006 | 512 | 396 | 379 | 371 | 218 |
| 2007 | 461 | 371 | 354 | 348 | 222 |
| 2008 | 503 | 409 | 397 | 391 | 230 |
| 2009 | 550 | 344 | 332 | 339 | 163 |
| 2010 | 421 | 312 | 301 | 295 | 136 |
| 2011 | 376 | 268 | 256 | 252 | 123 |

Table 4: Number of positive and null sets by fishing mode made by the French purse seine fishing fleet of the Indian ocean during 1981-2011. FAD = Fish Aggregating Device; FSC = Free-Swimming School

| | ALL | | | FAD | | | FSC | | |
|------|-------|----------|------|-------|----------|------|-------|----------|------|
| | Total | Positive | Null | Total | Positive | Null | Total | Positive | Null |
| 1981 | 56 | 44 | 12 | 32 | 29 | 3 | 24 | 15 | 9 |
| 1982 | 143 | 105 | 38 | 72 | 63 | 9 | 71 | 42 | 29 |
| 1983 | 1068 | 766 | 302 | 540 | 470 | 70 | 528 | 296 | 232 |
| 1984 | 3601 | 2211 | 1390 | 1143 | 971 | 172 | 2458 | 1240 | 1218 |
| 1985 | 3780 | 2274 | 1506 | 1353 | 1210 | 143 | 2427 | 1064 | 1363 |
| 1986 | 4446 | 2429 | 2017 | 1628 | 1393 | 235 | 2818 | 1036 | 1782 |
| 1987 | 4414 | 2813 | 1601 | 1908 | 1676 | 232 | 2506 | 1137 | 1369 |
| 1988 | 4824 | 2823 | 2001 | 1309 | 1177 | 132 | 3515 | 1646 | 1869 |
| 1989 | 3583 | 2243 | 1340 | 1436 | 1310 | 126 | 2147 | 933 | 1214 |
| 1990 | 4126 | 2527 | 1599 | 1189 | 1093 | 96 | 2937 | 1434 | 1503 |
| 1991 | 3630 | 2448 | 1182 | 1622 | 1538 | 84 | 2008 | 910 | 1098 |
| 1992 | 4602 | 2980 | 1622 | 1708 | 1569 | 139 | 2894 | 1411 | 1483 |
| 1993 | 4163 | 2763 | 1400 | 1810 | 1611 | 199 | 2353 | 1152 | 1201 |
| 1994 | 4332 | 3099 | 1233 | 2326 | 2068 | 258 | 2006 | 1031 | 975 |
| 1995 | 4486 | 3066 | 1420 | 2276 | 2052 | 224 | 2210 | 1014 | 1196 |
| 1996 | 3956 | 2883 | 1073 | 2221 | 1956 | 265 | 1735 | 927 | 808 |
| 1997 | 3607 | 2714 | 893 | 2301 | 2035 | 266 | 1306 | 679 | 627 |
| 1998 | 3328 | 2454 | 874 | 2117 | 1828 | 289 | 1211 | 626 | 585 |
| 1999 | 3238 | 2369 | 869 | 1750 | 1553 | 197 | 1488 | 816 | 672 |
| 2000 | 3333 | 2475 | 858 | 1818 | 1555 | 263 | 1515 | 920 | 595 |
| 2001 | 2984 | 2175 | 809 | 1327 | 1223 | 104 | 1657 | 952 | 705 |
| 2002 | 3325 | 2597 | 728 | 1924 | 1822 | 102 | 1401 | 775 | 626 |
| 2003 | 3651 | 2464 | 1187 | 1570 | 1405 | 165 | 2081 | 1059 | 1022 |
| 2004 | 4062 | 2580 | 1482 | 1511 | 1378 | 133 | 2551 | 1202 | 1349 |
| 2005 | 4442 | 3051 | 1391 | 1683 | 1532 | 151 | 2759 | 1519 | 1240 |
| 2006 | 4644 | 3148 | 1496 | 1893 | 1743 | 150 | 2751 | 1405 | 1346 |
| 2007 | 4381 | 2932 | 1449 | 1944 | 1747 | 197 | 2437 | 1185 | 1252 |
| 2008 | 3995 | 2874 | 1121 | 1936 | 1760 | 176 | 2059 | 1114 | 945 |
| 2009 | 2548 | 2060 | 488 | 1636 | 1502 | 134 | 912 | 558 | 354 |
| 2010 | 1927 | 1554 | 373 | 1304 | 1212 | 92 | 623 | 342 | 281 |
| 2011 | 1810 | 1446 | 364 | 1217 | 1132 | 85 | 593 | 314 | 279 |

Table 5: Catch by species for the French purse seine fishing fleet of the Indian ocean during 1981-2011

| Year | YFT | SKJ | BET | ALB | OTH | TOTAL |
|------|-------|-------|------|------|-----|--------|
| 1981 | 188 | 158 | 23 | 0 | 56 | 425 |
| 1982 | 1081 | 792 | 145 | 0 | 0 | 2018 |
| 1983 | 10400 | 8153 | 1536 | 0 | 136 | 20225 |
| 1984 | 39268 | 21979 | 5081 | 224 | 228 | 66781 |
| 1985 | 37706 | 29183 | 6477 | 445 | 483 | 74293 |
| 1986 | 40911 | 38786 | 6636 | 200 | 693 | 87227 |
| 1987 | 41012 | 41620 | 6701 | 217 | 43 | 89593 |
| 1988 | 56766 | 38094 | 7251 | 177 | 732 | 103020 |
| 1989 | 33548 | 45750 | 5764 | 6 | 0 | 85068 |
| 1990 | 45351 | 27873 | 5663 | 36 | 31 | 78954 |
| 1991 | 36597 | 38630 | 4217 | 848 | 0 | 80292 |
| 1992 | 43287 | 44081 | 2580 | 1344 | 0 | 91292 |
| 1993 | 37792 | 47155 | 3773 | 292 | 0 | 89012 |
| 1994 | 34025 | 57209 | 4101 | 282 | 0 | 95617 |
| 1995 | 37573 | 47365 | 5781 | 336 | 0 | 91055 |
| 1996 | 33861 | 38958 | 5500 | 377 | 0 | 78696 |
| 1997 | 29661 | 30222 | 6706 | 514 | 0 | 67103 |
| 1998 | 20892 | 29370 | 5305 | 446 | 0 | 56013 |
| 1999 | 29289 | 41619 | 7326 | 150 | 0 | 78384 |
| 2000 | 33696 | 40407 | 5464 | 309 | 122 | 79998 |
| 2001 | 31253 | 32074 | 5452 | 645 | 174 | 69599 |
| 2002 | 34568 | 54204 | 7802 | 194 | 195 | 96963 |
| 2003 | 63101 | 38258 | 6334 | 608 | 368 | 108670 |
| 2004 | 63174 | 37323 | 6798 | 77 | 649 | 108021 |
| 2005 | 57198 | 43220 | 6453 | 86 | 184 | 107140 |
| 2006 | 44495 | 47640 | 5573 | 850 | 233 | 98791 |
| 2007 | 32660 | 30438 | 6132 | 305 | 3 | 69539 |
| 2008 | 37642 | 29520 | 6794 | 952 | 10 | 74919 |
| 2009 | 22195 | 28690 | 5761 | 295 | 3 | 56944 |
| 2010 | 22490 | 20985 | 3584 | 29 | 16 | 47103 |
| 2011 | 21192 | 17871 | 3593 | 238 | 0 | 42894 |

Table 6: Catch by species made on FAD-associated schools for the French purse seine fishing fleet of the Indian ocean during 1981-2011

| Year | YFT | SKJ | BET | ALB | OTH | TOTAL |
|------|-------|-------|------|-----|-----|-------|
| 1981 | 37 | 128 | 20 | 0 | 56 | 240 |
| 1982 | 442 | 709 | 131 | 0 | 0 | 1282 |
| 1983 | 3959 | 6637 | 1381 | 0 | 136 | 12114 |
| 1984 | 10692 | 17600 | 3762 | 0 | 191 | 32244 |
| 1985 | 14623 | 26582 | 4993 | 14 | 459 | 46671 |
| 1986 | 15353 | 31040 | 4953 | 0 | 693 | 52038 |
| 1987 | 17926 | 30205 | 4937 | 0 | 20 | 53089 |
| 1988 | 12763 | 28633 | 4675 | 0 | 602 | 46673 |
| 1989 | 13769 | 26850 | 4499 | 0 | 0 | 45118 |
| 1990 | 10312 | 21046 | 3513 | 0 | 31 | 34902 |
| 1991 | 7895 | 36252 | 3048 | 0 | 0 | 47195 |
| 1992 | 11877 | 38559 | 2261 | 8 | 0 | 52705 |
| 1993 | 10982 | 39820 | 1958 | 5 | 0 | 52765 |
| 1994 | 12122 | 44944 | 3329 | 22 | 0 | 60417 |
| 1995 | 17561 | 38428 | 4720 | 16 | 0 | 60725 |
| 1996 | 15698 | 32902 | 4863 | 69 | 0 | 53532 |
| 1997 | 16963 | 25966 | 6406 | 65 | 0 | 49400 |
| 1998 | 11568 | 24810 | 4328 | 12 | 0 | 40718 |
| 1999 | 16449 | 31024 | 5831 | 103 | 0 | 53407 |
| 2000 | 16050 | 31939 | 4150 | 41 | 122 | 52302 |
| 2001 | 8844 | 26273 | 3884 | 108 | 174 | 39283 |
| 2002 | 13654 | 47378 | 6337 | 0 | 171 | 67540 |
| 2003 | 16810 | 33837 | 3429 | 0 | 134 | 54209 |
| 2004 | 13959 | 31473 | 4882 | 0 | 339 | 50653 |
| 2005 | 15399 | 31270 | 3667 | 0 | 184 | 50520 |
| 2006 | 14074 | 36207 | 4042 | 0 | 185 | 54507 |
| 2007 | 11986 | 23001 | 4127 | 3 | 3 | 39119 |
| 2008 | 11476 | 25446 | 3901 | 2 | 10 | 40835 |
| 2009 | 9936 | 26710 | 4226 | 10 | 3 | 40884 |
| 2010 | 10925 | 19827 | 2443 | 4 | 16 | 33216 |
| 2011 | 13153 | 16536 | 2277 | 30 | 0 | 31997 |

Table 7: Catch by species made on free-swimming schools for the French purse seine fishing fleet of the Indian ocean during 1981-2011

| Year | YFT | SKJ | BET | ALB | OTH | TOTAL |
|------|-------|-------|------|------|-----|-------|
| 1981 | 151 | 31 | 4 | 0 | 0 | 185 |
| 1982 | 638 | 83 | 14 | 0 | 0 | 736 |
| 1983 | 6441 | 1516 | 155 | 0 | 0 | 8111 |
| 1984 | 28576 | 4380 | 1319 | 224 | 37 | 34537 |
| 1985 | 23083 | 2601 | 1484 | 432 | 24 | 27623 |
| 1986 | 25558 | 7747 | 1683 | 200 | 0 | 35189 |
| 1987 | 23086 | 11415 | 1764 | 217 | 23 | 36505 |
| 1988 | 44003 | 9461 | 2575 | 177 | 130 | 56347 |
| 1989 | 19779 | 18900 | 1265 | 6 | 0 | 39951 |
| 1990 | 35039 | 6827 | 2150 | 36 | 0 | 44052 |
| 1991 | 28702 | 2378 | 1169 | 848 | 0 | 33097 |
| 1992 | 31410 | 5522 | 319 | 1336 | 0 | 38587 |
| 1993 | 26810 | 7335 | 1815 | 287 | 0 | 36247 |
| 1994 | 21903 | 12265 | 772 | 260 | 0 | 35200 |
| 1995 | 20012 | 8937 | 1061 | 320 | 0 | 30330 |
| 1996 | 18163 | 6056 | 637 | 308 | 0 | 25164 |
| 1997 | 12698 | 4256 | 300 | 449 | 0 | 17703 |
| 1998 | 9324 | 4560 | 977 | 434 | 0 | 15295 |
| 1999 | 12840 | 10595 | 1495 | 47 | 0 | 24977 |
| 2000 | 17646 | 8468 | 1314 | 268 | 0 | 27696 |
| 2001 | 22409 | 5801 | 1568 | 538 | 0 | 30316 |
| 2002 | 20913 | 6826 | 1465 | 194 | 24 | 29422 |
| 2003 | 46291 | 4422 | 2906 | 608 | 235 | 54461 |
| 2004 | 49215 | 5850 | 1916 | 77 | 310 | 57368 |
| 2005 | 41799 | 11950 | 2786 | 86 | 0 | 56620 |
| 2006 | 30421 | 11433 | 1531 | 850 | 48 | 44283 |
| 2007 | 20675 | 7438 | 2005 | 302 | 0 | 30420 |
| 2008 | 26166 | 4074 | 2893 | 950 | 0 | 34084 |
| 2009 | 12259 | 1980 | 1535 | 285 | 0 | 16059 |
| 2010 | 11565 | 1157 | 1140 | 25 | 0 | 13888 |
| 2011 | 8039 | 1335 | 1316 | 208 | 0 | 10897 |