

## Accounting for age uncertainty in growth modeling, the case study of yellowfin tuna (Thunnus albacares) of the Indian Ocean

Emmanuelle Dortel, F. Massiot-Granier, Etienne Rivot, J. Million, Jean-Pierre Hallier, Eric Morize, Jean-Marie Munaron, N. Bousquet, Emmanuel Chassot

## ▶ To cite this version:

Emmanuelle Dortel, F. Massiot-Granier, Etienne Rivot, J. Million, Jean-Pierre Hallier, et al.. Accounting for age uncertainty in growth modeling, the case study of yellowfin tuna (Thunnus albacares) of the Indian Ocean. PLoS ONE, 2013, 8 (4), pp.e60886. 10.1371/journal.pone.0060886 . ird-00833540

## HAL Id: ird-00833540 https://ird.hal.science/ird-00833540

Submitted on 13 Jun 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. APPENDIX A. Presentation of data used in the coupling of ageing error and growth models

TABLE A1. Summarize of data used in ageing error model; *It*: section between nucleus and OTC mark , *Im*: section between the OTC mark and edge, *Ir*: section between the nucleus and edge

	Fish from RTTP-IO program knowledge of time-at-liberty: with withou			without	Fish from WSTTP program
Otolith section	It	Ir	Im	It	It
Number of fishes	80	14	27	30	38
Fork length at tagging (cm)	43 to 85	50 to 72			
Fork length at recapture (cm)	49.7 to 135.4	59.5 to 114.1	49.7 to 131	47.9 to 146.5	19 to 46.6
Time-at-liberty (days)	43 to 969	33 to 414	43 to 969		



FIG. A1. Tagging area (gray-colored) and points of tag recovery (circles) of RTTP-IO program and sampling area (square) of WSTTP program

## APPENDIX B. Evaluation by simulation of ageing error model: general approach and results



FIG. B1. Simulation framework for testing the ageing error model; different sources of uncertainty are added to simulated ages to randomly generate noisy increments that were then used as inputs in the model



FIG. B2. Marginal posterior distributions of the ageing error model parameters (black) compared with the simulated distributions (grey) with individual variability (right) and without (left); a, b and c represents the first, second and third simulated data set respectively



FIG. B3. Boxplot of the relative mean square error (RMSE) of estimated age by ageing error model for 2,3,4 or 5 readings of the same otolith; a, b and c represents the first, second and third simulated data set respectively

TABLE B1. Comparison of the RMSE values of estimated age by ageing error model using a Wilcoxon test; 2L, 3L, 4L and 5L correspond to the number of readings of the same otolith; a, b and c represents the first, second and third simulated data set respectively

		4L	3L	2L
a	5L	V = 54527 p-value = 0.01224		
	4L		V = 48750 p-value = 1.767e-05	
	3L			V = 45892 p-value = 2.260e-07
b	5L	V = 54789 p-value = 0.01535		
	4L		V = 69539 p-value = 0.0324	
	3L			V = 45324 p-value = 8.683e-08
с	5L	V = 46959 p-value = 1.257e-06		
	4L		V = 69539 p-value = 0.0324	
	3L			V = 45324 p-value = 8.683e-08



FIG. B4. Boxplot of the relative mean square error (RMSE) of estimated age by the ageing error model, the conventional method and the intermediate method for 2, 3, 4 or 5 readings of the same otolith; a, b and c represents the first, second and third simulated data set respectively

TABLE B2. Comparison of the RMSE values of estimated age by ageing error model, the traditional method and the intermediate method using a Wilcoxon test; 2L, 3L, 4L and 5L correspond to the number of readings of the same otolith; a, b and c represents the first, second and third simulated data set respectively

			Traditional method	Intermediate method		
a	Ageing error model	2L	V = 28741, p-value < 2.2e-16	V = 55196, p-value = 0.02155		
		3L	V = 28165, p-value < 2.2e-16	V = 57314, p-value = 0.1004		
		4L	V = 28080, p-value < 2.2e-16	V = 59300, p-value = 0.3037		
		5L	V = 28796, p-value < 2.2e-16	V = 59429, p-value = 0.3229		
	Intermediate method	2L	V = 17140, p-value < 2.2e-16			
		3L	V = 16009, p-value < 2.2e-16			
		4L	V = 16139, p-value < 2.2e-16			
		5L	V = 18954, p-value < 2.2e-16			
	Ageing error model	2L	V = 37313, p-value = 4.85e-15	V = 52640, p-value = 0.002009		
		3L	V = 35248, p-value < 2.2e-16	V = 49274, p-value = 3.623e-05		
		4L	V = 32274, p-value < 2.2e-16	V = 53918, p-value = 0.007069		
h		5L	V = 35057, p-value < 2.2e-16	V = 55195, p-value = 0.02153		
U	Intermediate method	2L	V = 36636, p-value = 8.97e-16			
		3L	V = 35714, p-value < 2.2e-16			
		4L	V = 29036, p-value < 2.2e-16			
		5L	V = 32230, p-value < 2.2e-16			
с	Ageing error model	2L	V = 26437, p-value < 2.2e-16	V = 60299, p-value = 0.4719		
		3L	V = 26608, p-value < 2.2e-16	V = 62971, p-value = 0.9149		
		4L	V = 22487, p-value < 2.2e-16	V = 67331, p-value = 0.1455		
		5L	V = 23658, p-value < 2.2e-16	V = 68155, p-value = 0.08714		
	Intermediate method	2L	V = 24027, p-value < 2.2e-16			
		3L	V = 19659, p-value < 2.2e-16			
		4L	V = 14297, p-value < 2.2e-16			
		5L	V = 15527, p-value < 2.2e-16			



APPENDIX C. Bayesian fit of the growth model coupled with the ageing error model

FIG. C1. Marginal posterior distributions of the parameters of the growth model coupled with the ageing error model (black) compared with the prior distributions (grey)

	$L_{\infty}$	$k_1$	$k_2$	α	β	$t_0$
$L_{\infty}$	254.808	-0.93	-0.8	-0.29	-0.027	-0.43
<i>k</i> <sub>1</sub>	-0.593	0.002	0.8	0.21	0.093	0.67
<i>k</i> <sub>2</sub>	-5.527	0.014	0.189	0.48	-0.12	0.34
α	-7.568	0.001	0.034	0.026	-0.3	-0.074
β	-2.044	0.018	-0.259	-0.236	23.11	0.17
$t_0$	-0.601	0.002	0.013	-0.001	0.074	0.007

TABLE C1. Correlation-covariance matrix of growth parameters; numerals in bold represent the covariances