



Next generation sequencing yields complete mitogenomes of Leopard whipray (*Himantura leoparda*) and Blue-spotted stingray (*Neotrygon kuhlii*) (Chondrichthyes: Dasyatidae)

Kang-Ning Shen, Chih-Wei Chang, Shou-Yi Tsai, Shan-Chun Wu, Zi-Han Lin,
Yen-Fan Chan, Ching-Hung Chen, Chung-Der Hsiao, Philippe Borsa

► To cite this version:

Kang-Ning Shen, Chih-Wei Chang, Shou-Yi Tsai, Shan-Chun Wu, Zi-Han Lin, et al.. Next generation sequencing yields complete mitogenomes of Leopard whipray (*Himantura leoparda*) and Blue-spotted stingray (*Neotrygon kuhlii*) (Chondrichthyes: Dasyatidae) . Mitochondrial DNA Part A, 2016, 27, pp.2613-2614. 10.3109/19401736.2015.1041119 . ird-01312741v1


HAL Id: ird-01312741

<https://ird.hal.science/ird-01312741v1>

Submitted on 9 May 2016 (v1), last revised 9 May 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Next generation sequencing yields the complete
mitogenome~~S~~ of Leopard whipray (*Himantura leoparda*) and
Blue-spotted ~~maskray~~ (*Neotrygon kuhlii*) (Chondrichthyes:
Dasyatidae)**

Journal:	<i>Mitochondrial DNA</i>
Manuscript ID:	Draft
Manuscript Type:	Mitogenome Announcements
Date Submitted by the Author:	n/a
Complete List of Authors:	Shen, Kang-Ning; National Taiwan Ocean University, Center of Excellence for the Oceans Chang, Chih-Wei; National Museum of Marine Biology and Aquarium, Department of Biology Tsai, Shiou-Yi; Chung Yuan Christian University, Department of Bioscience Technology Wu, Shan-Chun; Chung Yuan Christian University, Department of Bioscience Technology Lin, Zi-Han; Chung Yuan Christian University, Department of Bioscience Technology Chan, Yen-Fan; Chung Yuan Christian University, Department of Bioscience Technology Chen, Ching-Hung; WeThink Biotech INC, Hsiao, Chung-Der; Chung Yuan Christian University, Department of Bioscience Technology Borsa, Philippe; Universitas Udayana,
Keywords:	Leopard whipray, Blue-spotted stingray, next generation sequencing, mitogenome

SCHOLARONE™
Manuscripts

1
2
3 MITOGENOME ANNOUNCEMENTS
4
5

6 Next generation sequencing yields the complete mitogenome^S of Leopard whipray
7 (Himantura leoparda) and Blue-spotted stingray (Neotrygon kuhlii) (Chondrichthyes:
8 Dasyatidae)
9
10

11
12
13
14 Kang-Ning Shen¹, Chih-Wei Chang^{2,3}, Shiou-Yi Tsai⁴, Shan-Chun Wu⁴, Zi-Han Lin⁴,
15
16 Yen-Fan Chan⁴, Ching-Hung Chen⁵, Chung-Der Hsiao^{4,*}, Philippe Borsa⁶
17
18
19

20
21 ¹ Center of Excellence for the Oceans, National Taiwan Ocean University, Keelung,
22 Taiwan
23
24

25 ² Department of Biology, National Museum of Marine Biology and Aquarium, Checheng,
26 Pingtung, Taiwan
27
28

29 ³ Graduate Institute of Marine Biodiversity & Evolutionary Biology, National Dong Hwa
30 University, Checheng, Pingtung, Taiwan
31
32

33 ⁴ Department of Bioscience Technology, Chung Yuan Christian University, Chung-Li,
34 Taiwan
35
36

37 ⁵ WeThink Biotech Inc., Taoyuan, Taiwan
38
39

40 ⁶ IRD-UMR 9220 eMtroPI c/o Universitas Udayana, Denpasar, Indonesia
41
42 Institut de recherche pour le développement,
43

44 * corresponding to : Chung-Der Hsiao, Department of Bioscience Technology, Chung
45 Yuan Christian University, 200 Chung-Pei Rd., Chung-Li, Taiwan 32023
46
47

48 e-mail address: cdhsiao@cycu.edu.tw
49
50

Abstract

The Leopard whipray (*Himantura leoparda*) and Blue-spotted stingray (*Neotrygon kuhlii*) are stingray distributed in the Indian and West Pacific Ocean and considered as species complex by morphological and molecular evidences. In this study, we used the next-generation sequencing method to decode the complete mitogenome of *H. leoparda* and *N. kuhlii*. The assembled mitogenome, consisting of 17,690 bp for *H. leoparda* and 17,974 bp for *N. kuhlii*, shows 78% identities each other. Both mitogenomes follow the typical vertebrate arrangement, including 13 protein coding genes, 22 transfer RNAs, 2 ribosomal RNAs genes and a non-coding control region of D-loop. D-loop contains 1,931 bp (*H. leoparda*) and 2,243 bp (*N. kuhlii*). length is located between tRNA-Pro and tRNA-Phe. The overall GC content is 40.3% for *H. leoparda* and 39.8% for *N. kuhlii*. The complete mitogenome of *H. leoparda* and *N. kuhlii* provides essential and important DNA molecular data for further phylogenetic and evolutionary analysis for stingray species complex.

Keywords: Leopard whipray, Blue-spotted stingray, mitogenome, next generation sequencing

The population genetics of stingrays are among the most poorly understood among vertebrates because of the limited geographic sampling and the occurrence of cryptic species (Beheregaray, 2008). The two targeted stingray species in this study ~~are classified~~ belong to in the family Dasyatidae which ~~encompasses~~ encompasses nine genera and about 70 species. A recent taxonomic revision of ocellated whipray species complex comprised at least four biological species mostly from the Coral Triangle (Arlyza et al., 2013), including three known species *Himantura leoparda*, *H. uarnak*, *H. undulata* and another newly described cryptic species *H. tutul* sp. nov. (Borsa et al., 2013). The ~~establishment~~ development of stingray mitogenome~~s~~ S is important to understand their population genetics, molecular phylogenetic, phylogeography and evolution ~~for stingray species complex~~.

Samples *H. leoparda* (voucher no. 329) and *N. kuhlii* (voucher no. 330) were collected from Taiwan and Cebu, the Philippines, respectively. The methods for genomic DNA extraction, library construction and next generation sequencing were followed by our previous publication (Shen et al., 2014). By using commercial software (Geneious V8, Auckland, New Zealand), about 0.16% (10,058 out of 6,280,529) and 0.02% raw reads (1,082 out of 5,005,674) were isolated from total genomic reads and *de novo* assembled to produce a single, circular form of complete mitogenome with about an average 185 X and 22 X coverage for *H. leoparda* and *N. kuhlii*, respectively. The assembled mitogenome of *H. leoparda* and *N. kuhlii*, consisting of 17,690 bp (GenBank: KR019776) and 17,974 bp (GenBank: KR019777), both showing 78% identities each other. In addition, the mitogenome of *N. kuhlii* decoded in this study show 97.6% identity to previously reported *N. kuhlii* species complex (Chen et al., 2014). D-loop contains 1,931 bp in *H. leoparda* and 2,243 bp in *N. kuhlii* and locates between tRNA-Pro and tRNA-Phe.

The protein coding, rRNA and tRNA genes of mitogenome were predicted by using DOGMA (Wyman et al., 2004), ARWEN (Laslett and Canback, 2008), MITOS (Bernt et al., 2013) and MitoAnnotator (Iwasaki et al., 2013) tools. For *H. leoparda*, six of 13

protein-coding genes is inferred to termination with an incomplete stop codon of either T- - (COX2, ND3, ND4 and CYTB) or TA- (ATP6 and COX3). For *N. kuhlii*, four of 13 protein-coding genes is inferred to termination with an incomplete stop codon of either T- - (COX2, ND3 and ND4) or TA- (ND2). The longest one is ND5 gene (1,848 bp for *H. leoparda* and 1,842 bp for *N. kuhlii*) in all protein coding genes, whereas the shortest is ATP8 gene (168 bp for both species). The two ribosomal RNA genes, 12S rRNA gene (968 bp for *H. leoparda* and 963 bp for *N. kuhlii*) and 16S rRNA gene (1,687 bp for *H. leoparda* and 1,701 bp for *N. kuhlii*), are located between tRNA-Phe and tRNA-Leu (UAA) and separated by tRNA-Val.

To validate the phylogenetic position of *H. leoparda* and *N. kuhlii*, we used MEGA6 (Tamura et al., 2013) software to construct a Maximum likelihood tree (with 500 bootstrap replicates) containing complete mitogenomes of 16 species derived from 11 different genus in Dasyatidae. *Chimaera monstrosa* derived from Chimaeridae was used as outgroup for tree rooting. Result shows *H. leoparda* is closely related to *H. granulata* and *N. kuhlii* is closely related to its species complex (Chen et al., 2014) with high bootstrap value supported (Figure 1). In conclusion, the complete mitogenome of *H. leoparda* and *N. kuhlii* are decoded for the first time in this study and provides essential and important DNA molecular data for further phylogenetic and evolutionary analysis for stingray species complex.

Declaration of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

Arlyza, I.S., Shen, K.N., Solihin, D.D., Soedharma, D., Berrebi, P., and Borsa, P. (2013).

1
2
3 Species boundaries in the *Himantura uarnak* species complex (Myliobatiformes:
4 Dasyatidae). Molecular phylogenetics and evolution 66, 429-435.
5 Beheregaray, L.B. (2008). Twenty years of phylogeography: the state of the field and the
6 challenges for the Southern Hemisphere. Molecular ecology 17, 3754-3774.
7 Bernt, M., Donath, A., Juhling, F., Externbrink, F., Florentz, C., Fritzsch, G., Putz, J.,
8 Middendorf, M., and Stadler, P.F. (2013). MITOS: improved de novo metazoan
9 mitochondrial genome annotation. Molecular phylogenetics and evolution 69, 313-319.
10 Borsa, P., Durand, J.D., Shen, K.N., Arlyza, I.S., Solihin, D.D., and Berrebi, P. (2013).
11 *Himantura tutul* sp. nov. (Myliobatoidei: Dasyatidae), a new ocellated whipray from the
12 tropical Indo-West Pacific, described from its cytochrome-oxidase I gene sequence.
13 Comptes rendus biologies 336, 82-92.
14 Chen, X., Xiang, D., Yu, J., Ding, W., and Zhang, S. (2014). Complete mitochondrial
15 genome of the blue-spotted stingray *Neotrygon kuhlii* (Myliobatiformes: Dasyatidae).
16 Mitochondrial DNA 25, 429-430.
17 Iwasaki, W., Fukunaga, T., Isagozawa, R., Yamada, K., Maeda, Y., Satoh, T.P., Sado, T.,
18 Mabuchi, K., Takeshima, H., Miya, M., et al. (2013). MitoFish and MitoAnnotator: a
19 mitochondrial genome database of fish with an accurate and automatic annotation
20 pipeline. Molecular biology and evolution 30, 2531-2540.
21 Laslett, D., and Canback, B. (2008). ARWEN: a program to detect tRNA genes in
22 metazoan mitochondrial nucleotide sequences. Bioinformatics 24, 172-175.
23 Shen, K.N., Yen, T.C., Chen, C.H., Li, H.Y., Chen, P.L., and Hsiao, C.D. (2014). Next
24 generation sequencing yields the complete mitochondrial genome of the flathead
25 mullet, *Mugil cephalus* cryptic species NWP2 (Teleostei: Mugilidae). Mitochondrial DNA,
26 1-2.
27 Tamura, K., Stecher, G., Peterson, D., Filipski, A., and Kumar, S. (2013). MEGA6:
28 Molecular Evolutionary Genetics Analysis version 6.0. Molecular biology and evolution
29 30, 2725-2729.
30 Wyman, S.K., Jansen, R.K., and Boore, J.L. (2004). Automatic annotation of organellar
31 genomes with DOGMA. Bioinformatics 20, 3252-3255.




Figure 1. Molecular phylogeny of *Himantura leoparda* and *Neotrygon kuhlii* in ~~Dasyatidae~~ based on complete mitogenome. The complete mitogenome was downloaded from GenBank and the phylogenetic tree is constructed by Maximum likelihood method with 500 bootstrap replicates. The gene's accession number for tree construction is listed as follows: *Dasyatis bennetti* (KC992792), *Pteroplatytrygon violacea* (KJ641617), *Taeniura meyeni* (JX827260), *Neotrygon kuhlii* (NC_021767; present study), *Neotrygon kuhlii* (KR019777), *Pastinachus atrus* (HG942172), *Himantura leoparda* (KR019776), *Himantura granulata* (KF751650), *Potamotrygon motoro* (KF709642), *Aetobatus flagellum* (KF482070), *Plesiobatis daviesi* (AY597334), *Mobula japanica* (JX392983), *Mobula eregoodootenkee* (KM361353), *Gymnura poecilura* (KJ617038) and *Chimaera monstrosa* (AJ310140).

Highlighted black: two new mitogenomes announced in present note.