



HAL
open science

How many genera of vascular plants are endemic to New Caledonia? A critical review based on phylogenetic evidence

Yohan Pillon, Laure Barrabé, Sven Buerki

► To cite this version:

Yohan Pillon, Laure Barrabé, Sven Buerki. How many genera of vascular plants are endemic to New Caledonia? A critical review based on phylogenetic evidence. *Botanical Journal of the Linnean Society*, 2017, 183 (2), pp.177-198. 10.1093/botlinnean/bow001 . ird-03644800

HAL Id: ird-03644800

<https://ird.hal.science/ird-03644800v1>

Submitted on 19 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

REVIEW

How many genera of vascular plants are endemic to New Caledonia? A critical review based on phylogenetic evidence

Yohan Pillon¹, Laure Barrabé² & Sven Buerki³

¹ Laboratoire de chimie des substances naturelles, Université de Limoges, 123 avenue Albert Thomas, 87060 Limoges cedex, France.

² Institut Agronomique néo-Calédonien (IAC)–Laboratoire de botanique et d'écologie végétale appliquées, Centre IRD, B.P. A5, 98800 Nouméa, Nouvelle-Calédonie.

³ Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

Running title: New Caledonian endemic genera

New Caledonia is a biodiversity hotspot located in the south-western Pacific, well known for its rich, unique and endangered flora. The island flora has a high level of endemism at the species level (75%), but also at the generic and family (three endemic) levels. We review here the taxonomic validity of the *c.* 100 endemic New Caledonian genera of vascular plants (13%) by using the monophyly criterion based on available phylogenetic data. As observed for other island floras, some of these genera were recovered nested in larger genera and are consequently likely to lose their rank. After a critical review, we concluded that the New Caledonian plant vascular flora contains between 62 and 91 endemic genera. This large variation in the number of endemic genera is mainly caused by a lack of DNA sequences (eight genera) and limited phylogenetic evidence. This work highlights gaps of knowledge that will have to be addressed to stabilize the taxonomy of the New Caledonian flora. Although this study shows that several genera are not monophyletic, New Caledonia still harbours more endemic genera than any other islands in the Pacific Ocean. Preliminary results indicate that the high level of endemism at higher taxonomic levels could be explained by an accumulation of relictual lineages, rather than adaptive radiations. Hypotheses explaining this phenomenon are provided in this study.

Keywords: adaptive radiation, biodiversity hotspots, biogeography, extinction, Pacific islands, relicts, ultramafic rocks.

Endemics can be either relicts—“left behind,” or they can be autochthonous—evolving their distinguishing characteristics, which place them in distinct genera or families, after arrival on islands. The two processes are not always different. Relicts may change after they have become island residents.

Sherwin Carlquist (1965: 305)

INTRODUCTION

New Caledonia is well known for its rich and unique flora and is the smallest biodiversity hotspot listed by Myers *et al.* (2000). This archipelago, located in the southwestern Pacific, has an unusual geological history and does not have a volcanic origin like most Pacific islands. New Caledonia detached from Australia during the late Cretaceous (*c.* 80 million years ago; Pelletier, 2006). During its drift eastward, New Caledonia was submerged under deep water as suggested by sediments (Cluzel, Chiron & Courme, 1998; Picard, 1999). Geological evidence strongly supports the emergence of New Caledonia at the Eocene (*c.* 37 million years ago) when a portion of oceanic crust was deposited over the island (Cluzel *et al.*, 1998; Pelletier, 2006). Weathering subsequently produced the ultramafic substrates, which are home to a peculiar flora (Virost, 1956; Jaffré, 1980; Isnard *et al.*, 2016). Thus in spite of its continental origin, the submergence of New Caledonia implies that the island was entirely re-colonized by terrestrial life after its Eocene emergence and it therefore fits the ‘Darwinian’ island hypothesis (Grandcolas *et al.*, 2008) comparable to most Pacific islands.

New Caledonia is not only unique because of its species diversity and rate of endemism, but also because of its unique diversity of gymnosperms and early diverging angiosperms (Morat *et al.*, 2012), many of which qualify as relicts (Grandcolas, Nattier &

Trewick, 2014). The island harbours three endemic families and 98 endemic genera of vascular plants, a number much greater than most other Pacific islands (Morat *et al.*, 2012). Recent reviews have suggested that the number of endemic genera on islands may be over-estimated for many reasons (Garnock-Jones, 2014), but a common pattern is that an endemic genus may be nested in a more widespread genus (Buerki *et al.*, 2013; Stuessy, König & Sepulveda, 2014). There is no clear universal definition of what a genus is; however, there are a number of criteria that can be applied to circumscribe them (see, e.g., Garnock-Jones, 2014, for a review), including monophyly, ease of recognition (morphologically), minimizing redundancy and time of divergence. Although some of these criteria cannot be applied objectively and the delimitation of a genus may be considered as arbitrary, they may actually be in many cases higher evolutionary significant units (Barraclough & Humphreys, 2015).

In this review, we aimed to evaluate the status of all of the endemic New Caledonian genera of vascular plants based on available phylogenetic evidence. We mostly investigated whether each genus fulfils the most objective criterion to define genera: monophyly. We do not discuss here morphological distinctiveness for each genus except in a few cases where phylogenetic evidence is ambiguous. This synthesis should allow a more objective comparison of the flora of New Caledonia with that of other Pacific islands, for which this work has already been done (e.g. New Zealand; Garnock-Jones, 2014). Identifying the most distinctive endemic genera is also paramount for the conservation of phylogenetic diversity (Mace, Gittleman & Purvis, 2003; Buerki *et al.*, 2015). It has long been accepted that the flora of New Caledonia is composed of ancient elements and the offspring of recent migrants (Virot, 1956: 373; Carlquist, 1965: 42), but their classification requires more scrutiny because supposedly ancient groups such as New Caledonian *Araucaria* Juss. may be more recent than previously thought (Kranitz *et al.*, 2014). Thus, we hoped to gain evidence on the prevailing patterns of endemism in New Caledonia: an ancient species repository or a hot-bed of recent

radiation (*sensu* Verboom *et al.*, 2009). We also hope that this review will trigger further taxonomic work on the flora of New Caledonia by raising potential issues on generic delimitations and also encourage the inference of a generic level phylogeny for this unique flora.

REVIEWING THE PHYLOGENETIC STATUS OF THE NEW CALEDONIAN ENDEMIC GENERA OF VASCULAR PLANTS

Endemic genera are shown in bold on first appearance. We follow here a linear sequence based on APG IV (2016) and consider all the endemic genera listed in Morat *et al.* (2012) and the few others described or re-instated after that publication. For non-monotypic genera, we indicated in parenthesis the number of species (a) included in the cited phylogenetic studies and the total number of species per genus (b) as follows: *genus* (a/b). Numbers of species in endemic genera are taken from Munzinger *et al.* (2016). All of the endemic genera formed a monophyletic clade in the cited phylogenetic studies unless otherwise stated.

FERNS

Gleicheniaceae

The monotypic ***Stromatopteris*** Mettenius is the only genus of fern endemic to New Caledonia. In molecular phylogenetic analyses including all genera of Gleicheniaceae, *Stromatopteris* is recovered with strong support as sister to the Indo-Pacific *Gleichenia* Sm. (Pryer *et al.*, 2004; Lehtonen, 2011).

GYMNOSPERMS

Podocarpaceae

The monotypic *Parasitaxus* de Laub. is the only known parasitic gymnosperm and is involved in an original type of parasitism with another podocarp [*Falcatifolium taxoides* (Brongn. & Gris) de Laub., also endemic to New Caledonia] that may also involve a fungus partner (Feild & Brodribb, 2005). In a molecular phylogenetic analysis including all genera of Podocarpaceae, *Parasitaxus* forms a moderately supported clade with *Manoao* Molloy (New Zealand) and *Lagarostrobos* Quinn (Tasmania), both monotypic, although their exact relationships are unresolved. *Parasitaxus* may be sister to the other two taxa (Sinclair *et al.*, 2002). A close relationship between these three genera was confirmed by Lu *et al.* (2014), but their relationships are not resolved.

Cupressaceae

Phylogenetic data have demonstrated that the monotypic *Neocallitropsis* Florin is nested in the Australian and New Caledonian *Callitris* Vent. (Pye, Gadek & Edwards, 2003). This lineage is the sister group to the New Caledonian *C. sulcata* Schltr. (*C. neocaledonica* Dümmer was not included). Consequently, the sole species, *N. pancheri* (Carrière) de Laub., was recently transferred to *Callitris* (Byng, 2015).

Taxaceae

The monotypic *Austrotaxus* Compton is the only representative of its family in the Southern Hemisphere. In a phylogenetic framework including all the genera of the family, Cheng *et al.* (2000) found that this genus was sister to the rest of tribe Taxeae (*Taxus* L., *Pseudotaxus* W.C.Cheng, from Eurasia, Malesia and North America) with strong support. This latter relationship was confirmed by Lu *et al.* (2014).

ANGIOSPERMS

AMBORELLALES, Amborellaceae

Amborella Baill. (monotypic) is now usually accepted as the sister lineage to all angiosperms (Qiu *et al.*, 1999; Soltis *et al.*, 2011), although a few analyses have recovered Nymphaeales as its sister group (e.g. Goremykin *et al.*, 2003). *Amborella trichopoda* Baill. is the first New Caledonian endemic plant to have had its genome entirely sequenced (Amborella Genome Project, 2013).

MAGNOLIIDS

LAURALES

Atherospermataceae

In a phylogenetic analysis including all genera of Atherospermataceae, the monotypic *Nemuaron* Baill. is recovered as sister to the Australian monotypic *Atherosperma* Labill. (Renner, Foreman & Murray, 2000).

Monimiaceae

Renner *et al.* (2010) recovered the monotypic *Kibaropsis* Vieill. ex Jérémie in clade III, which also includes the Oceanian *Levieria* Becc. and *Hedycarya* J.R.Forst. & G.Forst. Support in this clade is moderate, but *Hedycarya* appears polyphyletic, with *Kibaropsis* and *Levieria* nested inside it. *Kibaropsis* seems closer to the New Zealand *H. arborea* J.R.Forst. &

G.Forst. (bootstrap 65%) rather than the New Caledonian *H. cupulata* Baill. Generic delimitation in this group requires further investigations.

Lauraceae

Adenodaphne S.Moore (1/4) has been recovered as the sister group to two species of *Litsea* Lam. from Japan and China with moderate support (Chanderbali, van der Werff & Renner, 2001). Delimitation in *Litsea* (c. 400 species) and its allied genera requires clarifications (e.g. Fijridiyanto & Murakami, 2009). Since morphological confusions are possible between *Adenodaphne* and *Litsea*, also present in New Caledonia (Kostermans, 1974), their distinctiveness needs to be critically assessed.

MONOCOTS

LILIALES, Campynemataceae

Campynemataceae comprise only two genera, the Tasmanian monotypic *Campynema* Labill. and the New Caledonian *Campynemanthe* Baill. (1/3) that have been recovered as sister groups in phylogenetic analyses (Vinnersten & Bremer, 2001; Mennes *et al.*, 2015).

ASPARAGALES, Orchidaceae

The exact placement of some of the endemic genera (all terrestrial) is still somewhat uncertain, mostly because a number of related genera in this diverse family have not yet been included in any molecular phylogenetic analysis. This is particularly true for subtribe Goodyerinae (Chase *et al.*, 2015). *Clematepistephium* N.Hallé and *Eriaxis* Rchb.f. (both monotypic, tribe Vanilleae) are sister groups and together seem to be sister to the East Asian-New Guinean *Lecanorchis* Blume (Cameron, 2009). *Pachyplectron* Schltr. (1/3), has been

recovered as sister to the rest of tribe Goodyerinae (Alvarez-Molina & Cameron, 2009).

Gonatostylis Schltr. (1/2) seems to be sister to *Odontochilus* Blume (synonym *Pristiglottis* Cretz. & J.J.Sm.; Alvarez-Molina & Cameron, 2009), although the sampling of genera of Goodyerinae and *Odontochilus* spp. is limited in this study. The latter genus is found in Asia and the Pacific, including New Caledonia. *Gonatostylis* is also morphologically similar to the mycotrophic *Danhatchia* Garay & Christenson (Ormerod & Cribb, 2003) from New Zealand and Australia, which has never been included in any molecular phylogenetic studies.

Achlydosa M.AClem. & D.L.Jones (monotypic) has been found to be sister to the Australasian *Pterostylis* R.Br. (Cisternas *et al.*, 2012), the sole other member of subtribe Pterostylidinae (Chase *et al.*, 2015), also present in New Caledonia. *Coilochilus* Schltr. (monotypic) is sister to *Cryptostylis* R.Br. (Kores *et al.*, 2001), the sole other member of subtribe Cryptostylidinae (Chase *et al.*, 2015) from South-east Asia and the Pacific, including New Caledonia.

ARECALES, Arecaceae

The taxonomic updates by Pintaud & Baker (2008), associated with a complete generic-level phylogenetic analysis of the family (Baker *et al.*, 2009), resulted in a significant decrease in the number of palm genera in New Caledonia, placing in synonymy six endemic genera. More recently, *Pritchardiopsis* Becc. (monotypic) was placed in synonymy with *Saribus* Blume (Bacon & Baker, 2011; Bacon, Baker & Simmons, 2012), that occurs in South-east Asia and the Pacific. There are currently eight endemic palm genera recognized in New Caledonia, all members of tribe Areceae. Subtribe Clinospermatinae are endemic to New Caledonia and is composed of *Clinosperma* Becc. (4/4) and *Cyphokentia* Brongn (2/2). The study by Baker *et al.* (2011), including all of the species of Areceae, confirmed that the two genera are reciprocally monophyletic. Subtribe Archontophoenicinae include *Actinokentia* Dammer

(2/2), *Chambeyronia* Vieill. (2/2) and *Kentiopsis* Brongn. (4/4), the Australian *Archontophoenix* H.Wendl. & Drude, and the Papuan *Actinorhytis* H.Wendl. & Drude, but relationships and generic delimitations between the three New Caledonian genera and *Archontophoenix* remain unclear, including the monophyly of *Chambeyronia* and *Kentiopsis* (Domenech *et al.*, 2014). *Basselinia* Vieill. (3/14), *Burretiokentia* Pic.Serm. (2/5) and *Cyphophoenix* H.Wendl. ex Benth. & Hook.f (3/4) are placed in subtribe Basseliniinae. The phylogenetic study by Baker *et al.* (2011) suggested that *Cyphophoenix nucele* H.E.Moore could be sister to *Burretiokentia*, whereas two other *Cyphophoenix* spp, form a clade sister to the Melanesian genus *Physokentia* Becc. This study also indicates that three *Basselinia* spp. and the Lord Howe Island endemic *Hedyscepe* H.Wendl. & Drude form a polytomy, although *Hedyscepe* has been assigned to a different subtribe, Rhopalostylidinae. Further work is needed to clarify the taxonomic circumscriptions of the genera in subtribes Archontophoenicinae and Basseliniinae.

POALES, Poaceae

The dwarf bamboo *Greslania* Balansa (2/3) has been recovered with strong support as sister to an Australasian clade composed of *Cyrtochloa* S.Dransf., *Dinochloa* Buse, *Mullerochloa* K.M.Wong, *Neololeba* Widjaja, *Parabambusa* Widjaja, *Pinga* Widjaja and *Sphaerobambos* S.Dransf. (Chokthaweeapanich, 2014).

EUDICOTS

PROTEALES, Proteaceae

In a molecular phylogenetic tree including all genera of Proteaceae (Sauquet *et al.*, 2009), *Beaupreopsis* Virost (monotypic) formed a strongly supported clade with the Tasmanian genus

Cenarrhenes Labill. (monotypic) and the Malagasy genus *Dilobeia* Thou. (two species), and the last two are moderately supported as sister groups. *Beauprea* Brongn. & Gris (9/13) is monophyletic (He, Lamont & Fogliani, 2016) and is strongly supported as the sister lineage to tribe Proteeae (*Protea* L. and *Faurea* Harv. from Africa; Sauquet *et al.*, 2009). The two species of New Caledonian *Knightia* R.Br. are not related to the type species of this genus from New Zealand and are instead placed in an endemic genus *Eucarpha* (R.Br.) Spach (1/2) by Weston & Barker (2006), although the nomenclatural combinations have never been published. *Eucarpha* is strongly supported as the sister group to the eastern Australian *Triunia* L.A.S.Johnson & B.G.Briggs (Sauquet *et al.*, 2009). *Kermadecia* Brongn. & Gris (3/4) and *Sleumerodendron* Viot (monotypic) are part of a well-supported subtribe Gevuiniinae (Mast *et al.*, 2008) in which they form a well-supported clade with *Turrillia* A.C.Sm. (1/3, Fiji, Vanuatu). There is a weak support for a sister relationship between *Turrillia* and *Kermadecia*, which are in turn sister to *Sleumerodendron* (Mast *et al.*, 2008). The species of these three morphologically close genera (Viot, 1967; Smith & Haas, 1975) were all once placed in *Kermadecia* and further evidence for their reciprocal monophyly is desirable. *Virotia* L.A.S.Johnson & B.G.Briggs (2/6) is part of subtribe Virotiinae (Mast *et al.*, 2008) with *Athertonia* L.A.S.Johnson & B.G.Briggs (Queensland) and *Helicopsis* Sleumer (Indo-Malesia); their relationships are unresolved. *Garnieria* Brongn. & Gris (monotypic) is nested in the larger Australian genus *Persoonia* Michx. with the New Zealand endemic *Toronia* L.A.S.Johnson & B.G.Briggs and the south-western Australian *Acidonia* L.A.S.Johnson & B.G.Briggs (Weston & Porter, unpub. data, cited in Weston & Barker, 2006).

ROSIDS

FABALES

Fabaceae

Egan *et al.* (2016) found that *Arthroclianthus* Baill. (2/19) and *Nephrodesmus* Schindl. (2/5) were sister groups and reciprocally monophyletic in tribe Desmodieae.

Polygalaceae

Balgoya Morat & Meijden (monotypic) is generally placed in tribe Moutabeae and the study of Persson (2001), which included only a short *trnL* sequence for this genus, failed to resolve its phylogenetic position in the family or to confirm its placement in this tribe.

ROSALES

Rhamnaceae

The recently described *Jaffrea* H.C.Hopkins & Pillon (2/2, Hopkins *et al.*, 2015a) is sister to *Emmenosperma* F.Muell. (occurring in Malesia, Australia, New Caledonia and Fiji).

Moraceae

In a molecular phylogenetic analysis including 32 of 37 genera of Moraceae, Zerega *et al.* (2005) found strong support for a sister relationship between the New Caledonian endemic *Sparattosyce* Bureau (1/2) and the New Guinean *Antiaropsis* K.Schum (two species).

FAGALES, Myricaceae

In a molecular phylogenetic analysis including all genera of Myricaceae, Herbert *et al.* (2006) found strong support for a placement of *Canacomyrica* Guillaumin (monotypic) as the sister group to the rest of this widespread family.

CELASTRALES, Celastraceae

In a phylogenetic study including all genera of tribe Celastreae (Simmons *et al.*, 2008), *Salaciopsis* Baker f. (3/6) occupies an unresolved position between several multigeneric clades. *Menepetalum* Loes. (2/4) is sister to *Dinghousia* R.H.Archer from Queensland and *Dicarpellum* (Loes.) A.C.Sm. (2/4) is sister to *Hypsophila* F.Muell., also from Queensland (Coughenour *et al.*, 2010; Simmons *et al.*, 2012b). *Peripterygia* Loes. (monotypic) forms a weakly supported clade with the Australasian *Siphonodon* Griff. and *Crossopetalum* P.Browne (West Indies, Africa), but their exact relationships are unresolved (Simmons *et al.*, 2012a).

OXALIDALES, Cunoniaceae

In a phylogenetic study including all but two genera of Cunoniaceae, *Codia* J.R.Forst. & G.Forst. is recovered with strong support as the sister group to the Australian *Callicoma* Andrews (Hopkins, Rozefelds & Pillon, 2013) and is probably monophyletic (15/15, Pillon *et al.*, 2009a; Pillon pers. obs.). The phylogenetic position of *Hooglandia* McPherson & Lowry (monotypic) is unresolved somewhere among the early branches of the phylogenetic tree for Cunoniaceae (Sweeney, Bradford & Lowry, 2004; Hopkins *et al.*, 2013; Hopkins, Pillon & Hoogland, 2014). The latter genus has some morphological affinities with *Aistopetalum* Schltr. (McPherson & Lowry, 2004; H.C. Hopkins, pers. comm.), a genus of two species occurring in New Guinea that has not yet been included in any molecular phylogenetic study. *Pancheria* Brongn. & Gris (3/27) is placed in tribe Cunonieae and its relationship to *Cunonia*

L. (New Caledonia, southern Africa) and *Weinmannia* L. (pantropical) is unclear, although it might be sister to both (Bradford, 2002).

MALPIGHIALES

Clusiaceae

In a phylogenetic study including all but one genus of Clusiaceae, Ruhfel *et al.* (2011) recovered *Montrouziera* Pancher ex Planch. & Triana (2/6) as sister group to the South American *Platonia* Raf. with low support.

Salicaceae

In a phylogenetic study of former Flacourtiaceae (Alford, 2005), *Lasiochlamys* Pax & K.Hoffm. (1/11) was found to be nested in *Xylosma* G.Forst. (America, Asia and Pacific). However, because the single species sampled, *L. reticulata* Pax & K.Hoffm., was sister to a *Xylosma* sp. from New Caledonia (*X. vincentii* Guillaumin), the author thought there might be a possible misidentification or a possible plastid capture confounding the phylogenetic signal. *Lasiochlamys* differs from *Xylosma* in having petals (Lescot, 1980), but otherwise the two genera are similar and the presence/absence of petals has been shown to be highly homoplasious in this family (Alford, 2005).

Euphorbiaceae

In the analysis of Horn *et al.* (2012) including all the genera of tribe Euphorbieae, the monotypic *Neoguillauminia* Croizat is sister to the Australian *Calycopeplus* Planch. (five species) and these two genera are in turn sister to the large genus *Euphorbia* L. In the study of Tokuoka (2007), *Cocconerion* Baill. (1/2) is sister to *Baloghia* Endl. (Australia, New

Caledonia, Norfolk and Lord Howe Islands). The distinction between the two genera may warrant further investigations. *Bocquillonia* Baill. (1/14) is nested with strong support in the pantropical *Alchornea* Sw. (Wurdack, Hoffmann & Chase, 2005), although the latter study included a single species of *Bocquillonia* and two of *Alchornea*. Webster (1994) thought that most genera of subtribe Alchorneinae are only weakly separated from *Alchornea*. *Myricanthe* Airy Shaw (monotypic) has never been included in any molecular phylogenetic study and was classified as a close relative to *Cocconerion*, the Australian *Bertya* Planch. and the Bornean endemic *Borneodendron* Airy Shaw based on morphological similarities (Webster, 1994).

Picrodendraceae

In the analysis of Tokuoka & Tobe (2006) including only four genera of Picrodendraceae (out of c. 24), *Scagea* McPherson (1/2) is sister to the rest of the family. *Longetia* Baill. (monotypic) has not yet been included in any molecular phylogenetic studies.

MYRTALES, Myrtaceae

Carpolepis (J.W.Dawson) J.W.Dawson (3/3) is nested in *Metrosideros* Banks ex Gaertn. (Wright *et al.*, 2000; Pillon *et al.*, 2015), a genus widespread across the Pacific Islands (including New Caledonia), of which it is now considered a synonym (Pillon *et al.*, 2015). *Cloezia* Brongn. & Gris (1/6) is placed with strong support with tribe Tristanieae (Australasia, Biffin *et al.*, 2010), although its exact sister group has not been identified, but it may be sister to the entire tribe. In the analysis by Snow *et al.* (2011), *Kanakomyrtus* N.Snow (3/6, tribe Myrteae) is polyphyletic with its species mixed with those of the genera *Rhodomomyrtus* (DC.) Rchb. (South-east Asia and Pacific), *Pilidiostigma* Burret (Australasia) and *Octamyrtus* Diels (Malesia). The monotypic *Myrtastrum* Burret is part of the “Australasian group” of Myrteae and is the earliest diverging lineage in this clade (T.N.C. Vasconcelos, pers. comm.).

Arillastrum Pancher ex Baill. (monotypic) is well supported as sister to *Corymbia* K.D.Hill & L.A.S.Johnson + *Angophora* Cav. in Eucalypteae (Australasia) (Thornhill *et al.*, 2015). Tribe Xanthostemoneae is composed of three genera all represented in New Caledonia [*Pleurocayptus* Brongn. & Gris (two species), *Purpureostemon* Gugerli (monotypic) and the much larger *Xanthostemon* F.Muell. (also Australia, New Guinea, Philippines, eastern Indonesia and Solomon Islands)]. The first two have never been included in any molecular phylogenetic analysis and morphological data suggest that *Pleurocalyptus* could be nested in *Xanthostemon* (Wilson, 2011).

CROSSOSOMATALES, Strasburgeriaceae

The monospecific genus *Strasburgeria* Baill., once considered the sole member of Strasburgeriaceae, has been recovered as sister to the New Zealand endemic *Ixerba* A.Cunn. (Cameron, 2003) in the poorly known order Crossossomatales.

SAPINDALES

Sapindaceae

Sapindaceae are highly diverse in New Caledonia with four endemic genera: *Gongrodiscus* Radlk. (3/3), *Loxodiscus* Hook.f. (monotypic), *Podonephelium* Baill. (7/9), and *Storthocalyx* Radlk. (4/5). Buerki *et al.* (2012) have formally tested the monophyly and phylogenetic relationships of these endemic genera by including representatives into a generic level phylogenetic analysis of Sapindaceae (Buerki *et al.*, 2009). This study confirmed the monophyly of all the endemic genera, which belong to two different phylogenetic lineages: the *Cupania* L. group (Sapindoideae) and the *Dodonaea* Mill. group (Dodonaeoideae). In the *Cupania* group, *Gongrodiscus* and *Storthocalyx* were inferred as closely related together with

one species of the polyphyletic genus *Sarcotoechia* Radlk. from Australia; however, this relationship has to be clarified by sampling more DNA regions. *Podonephelium* belongs to another clade in the *Cupania* group and is sister to the widely distributed genus *Alectryon* Gaertn. (Malesia, Australia, New Caledonia, New Zealand and Hawaii). In the *Dodonaea* group, the monotypic *Loxodiscus* is supported as sister to a clade comprising *Harpullia* Roxb. (India, south-eastern China, Malesia to Australia, New Caledonia and Pacific Islands), *Cossinia* Comm. ex Lam. (Mascarenes, New Caledonia, eastern Australia, Fiji), *Diplopeltis* Endl. (north-western Australia) and *Dodonaea* (mainly in Australia, Malesia, New Guinea, Caribbean and Madagascar).

Rutaceae

Appelhans *et al.* (2014) found that *Comptonella* Baker f. (4/8), *Dutaillyea* Baill. (1/2) and *Picrella* Baill. (2/3) were nested in the larger and more widespread *Melicope* J.R.Forst. & G.Forst. (South-east Asia and Pacific), being close to the New Caledonian members of this genus. The authors suggested that *Dutaillioopsis* T.G.Hartley (monotypic) may also be nested in *Melicope*, although it has not been included in any molecular phylogenetic studies.

Oxanthera Montrouz. (2/5) is nested in the large genus *Citrus* L. (Bayer *et al.*, 2009).

Boronella Baill. (2/4) is nested in the Australian *Boronia* Sm. (Bayly *et al.*, 2013) and its species have now been transferred into this genus (Bayly *et al.*, 2015). *Crossosperma*

T.G.Hartley (1/2) has been recovered with strong support as sister to the Australian *Acradenia* Kippist (both species included). The relationship between *Myrtopsis* O.Hoffm. (2/8) and a number of Australian genera placed in tribe Boronieae, some of medium size [*Leionema* (F.Muell.) Paul G.Wilson, *Phebalium* Vent., *Philotheca* Rudge], are unresolved.

Neoschmidia T.G.Hartley (1/2) is recovered as sister to a diverse clade of Australasian genera including *Halfordia* F.Muell. and *Myrtopsis* (Bayly *et al.*, 2013).

MALVALES

Malvaceae

In a phylogenetic study including all the genera of subfamily Sterculioideae, Wilkie *et al.* (2006) recovered *Acropogon* Schltr. (2/25) as a member of the moderately supported *Brachychiton* Schott & Endl. clade. The relationships between *Acropogon*, *Brachychiton* (Australia-New Guinea), *Argyrodendron* (Endl.) Klotzsch (Australia) and *Franciscodendron* B.Hyland & Steenis (Australia) are unresolved. In a phylogenetic study including all eight genera of tribe Lasiopetaleae (Whitlock *et al.*, 2011), *Maxwellia* Baill. (monotypic) formed a strongly supported clade with *Hannafordia* F.Muell. (Australia) and a group of four genera (*Thomasia* J.Gay, *Lysiosepalum* F.Muell., *Lysiopetalum* K.Schum., *Guichenotia* J.Gay; all Australian), but their exact relationships are unresolved.

Thymelaeaceae

The phylogenetic relationships of *Deltaria* Steenis (monotypic) and *Solmsia* Baill. (1/2) with the genera *Lethedon* Spreng. (New Caledonia, Australia), *Gonystylus* Teijsm. & Binn. (South-east Asia) and *Arnhemia* Airy Shaw (Australia) in subfamily Gonystyloideae are unresolved (Beaumont *et al.*, 2009).

SANTALALES

Balanophoraceae

In a phylogenetic study of Balanophoraceae, *Hachettea* Baill. (monotypic) was recovered as sister to the New Zealand endemic *Dactylanthus* Hook.f. (Su *et al.*, 2015).

Santalaceae

In a phylogenetic analysis including all of the accepted genera of Santalaceae (Der & Nickrent, 2008), *Amphorogyne* Stauffer & Hurlim. (1/3) and *Daenikera* Hurlim. & Stauffer (monotypic) are retrieved as sister and inferred as sister to the rest of tribe Amphorogyneae. *Elaphanthera* N.Hallé (monotypic) was considered as a synonym of *Exocarpos* Labill. (South-east Asia and Pacific) by Der & Nickrent (2008), but the single species of this genus has not yet been included in a molecular phylogenetic framework.

ASTERIDS

ERICALES

Sapotaceae

Since the earliest monograph was published (Aubréville, 1967), New Caledonian Sapotaceae have been the subject of major generic rearrangements over the past few years (Bartish *et al.*, 2005; Swenson *et al.*, 2008; Swenson, Nylinder & Munzinger, 2013) and several genera once considered as endemic to New Caledonia have been placed in synonymy (*Corbassona* Aubrév., *Leptostylis* Benth., *Ochrothallus* Pierre ex Baill., *Pyriluma* Aubrév., *Sebertia* Pierre ex Engl. and *Trouettia* Pierre ex Baill.) or now include species from outside New Caledonia (*Pichonia* Pierre). The expanded *Pycnandra* Benth. has emerged as the largest endemic genus of New Caledonia with 59 described species. It is sister to the Australian *Niemeyera* F.Muell. (Swenson *et al.*, 2013) and a study including 60 of 68 presumed species (some undescribed) indicated that it is monophyletic (Swenson *et al.*, 2015).

Primulaceae

The recently described monotypic genus *Mangenotiella* M.Schmid (Schmid, 2012) has not yet been included in any phylogenetic study. There has been little phylogenetic work on woody Primulaceae (but see Strijk *et al.*, 2014), and *Mangenotiella* needs to be critically compared with *Tapeinosperma* Hook.f. (also present in New Caledonia) and *Discocalyx* Mez (South-east Asia to Pacific), the distinctness of which has been questioned (e.g. Smith, 1973).

Ericaceae

The phylogenetic relationships between *Cyathopsis* Brongn. & Gris (1/3) and a number of other genera of the Styphelieae are unresolved (Quinn *et al.*, 2005; Puente-Lelièvre *et al.*, 2015), although an analysis based on the nuclear ITS region alone suggests a weakly supported sister relationship with the Australian *Lissanthe* R.Br.

ICACINALES, Oncothecaceae

Oncotheca Baill. (1/2) is the only genus of Oncothecaceae. Oncothecaceae were not placed in any order in APG III (2009), because most phylogenetic studies had failed to clarify an exact position in lamiids (Bremer *et al.*, 2002; Cameron, 2003; González *et al.*, 2007; Soltis *et al.*, 2011; Byng *et al.*, 2014). A recent study using complete plastid genomes recovered this genus as sister to a narrowly defined Icacinaeae (Stull *et al.*, 2015) and it was placed in Icacinales in APG IV (2016) to reflect this finding.

GENTIANALES

Rubiaceae

Morierina Vieill. (2/2) and *Thiollierea* Montrouz. (12/12, previously included under *Bikkia*

Reinw., but reinstated by Barrabé *et al.*, 2011), form a well-supported clade, sister to an alliance of several Caribbean genera mainly occurring in Cuba with high support (*Ceratopyxis* Hook.f., *Eosanthe* Urb., *Phialanthus* Griseb. and *Schmidtottia* Urb.; Manns & Bremer, 2010). Although Manns & Bremer (2010) placed *Morierina montana* Vieill. as sister to two *Thiollierea* spp., uncertainties remain about the internal topology of the ‘*Morierina-Thiollierea*’ clade since an unpublished study, based on a complete species sampling and five DNA loci, has shown unresolved interspecific and intergeneric relationships (Mouly, pers. comm.). *Guettarda* L. has been revealed to be polyphyletic (Achille *et al.*, 2006). Of the 20 New Caledonian species, only the type species *G. speciosa* L. belongs to *Guettarda* s.s. The other species should be included in *Antirhea* Juss., *Guettardella* Benth., *Timonius* DC., *Tinadendron* Achille and a new endemic genus, ***Gea*** Achille ined. (2/6). Robust morphological characters support the last genus, composed of six species (Achille, pers. comm.), although this genus is unpublished and its monophyly has not been demonstrated. The monotypic *Normandia* Hook.f. was found to be nested in *Coprosma* J.R.Forst. & G.Forst. (Anderson, Rova & Andersson, 2001), but this study was based on an incorrect DNA sequence (Markey, Lord & Orlovich, 2004). Recently, *Normandia* has been excluded from this genus with strong support and placed as the possible sister-group of the large alliance formed by *Coprosma* and *Nertera* Gaertn. (Markey *et al.*, 2004; Cantley *et al.*, 2014).

Apocynaceae

In the analysis of Potgieter & Albert (2001), *Cerberiopsis* Vieill. ex Pancher & Sebert (1/3) is strongly supported as sister to *Cerbera* L. (Indian Ocean, Australasia, Pacific). Livshultz *et al.* (2007) found *Artia* Guillaumin (1/4) to be sister to the Australasian *Parsonsia* R.Br. noting the similarities between the two genera and acknowledging that only a few *Parsonsia* spp. were included in the phylogenetic analysis. The two genera are in turn nested in the

Neotropical *Prestonia* R.Br. Therefore, generic delimitations need to be further evaluated in this group. *Artia* was not considered as an endemic genus in Morat *et al.* (2012), but this was an error.

LAMIALES, Gesneriaceae

In a phylogenetic analysis including all of the genera of tribe Coronanthereae, Woo *et al.* (2011) found that *Depanthus* S.Moore (1/2) was either sister to the Australian *Lenbrassia* G.W.Gillett or to the Lord Howe island endemic *Negria* F.Muell.

AQUIFOLIALES, Stemonuraceae

Gastrolepis Tiegh. (two species) has not yet been included in any molecular phylogenetic analysis.

ASTERALES

Alseuosmiaceae

In a phylogenetic analysis published by Lundberg & Bremer (2003) *Platyspermation* Guillaumin (monotypic) was inferred as sister to the rest of the family (four out of five genera included). *Periomphale* Baill. (monotypic) was once synonymized with *Wittsteinia* F.Muell., a view latter rejected by Tirel & Jérémie (1996). Unpublished phylogenetic analyses support the latter conclusion (A. Mouly, pers. comm.).

Phellinaceae

Phelline Labill. (3/14) is the only genus of Phellinaceae, which form a clade with Alseuosmiaceae and Argophyllaceae in Asterales (Kårehed *et al.*, 1999). The relationships

between these three small and poorly known families are unclear. Phellinaceae were placed with strong support as sister to Argophyllaceae in the analysis of Tank & Donoghue (2010), but in the analysis of Soltis *et al.* (2011) including more DNA regions, they were placed as sister to a weakly supported clade including Alseuosmiaceae and Argophyllaceae.

PARACRYPHIALES, Paracryphiaceae

Paracryphia Baker f. (monotypic) was once considered as the sole member of Paracryphiaceae, which has been expanded to include *Sphenostemon* Baill. and *Quintinia* A.DC. (APG III, 2009). There is strong support for a sister-group relationship between *Paracryphia* and *Sphenostemon* from Australia, New Guinea and New Caledonia (Tank & Donoghue, 2010; Soltis *et al.*, 2011).

APIALES

Myodocarpaceae

Myodocarpaceae have a centre of diversity in New Caledonia and include only two genera, which are reciprocally monophyletic (Nicolas & Plunkett, 2009): the endemic New Caledonian *Myodocarpus* Brongn. & Gris (4/10) and *Delarbrea* Vieill. (Malesia to New Caledonia).

Apiaceae

In a phylogenetic study including 40 of the 42 genera previously placed in subfamily Hydrocotyloideae, Nicolas & Plunkett (2009) found that *Apiopetalum* Baill. (2/2) was sister to *Actinotus* Labill. (Australasian) in the newly circumscribed subfamily Mackinlayoideae.

DISCUSSION

How many endemic genera of vascular plants occur in New Caledonia?

Current phylogenetic evidence suggests that the monophyly of 30 New Caledonian endemic genera still needs to be confirmed or has been challenged (Table 1) and eight are clearly nested in larger, more widespread genera; some have been synonymized since Morat *et al.* (2012). In most problematic cases, the endemic genera are nested in a larger more widespread genus (e.g. *Neocallitropsis* nested in *Callitris*, and *Comptonella*, *Dutaillyea* and *Picrella* in *Melicope*) or their distinction from such genera needs further testing (e.g. *Lasiochlamys* vs. *Xylosma*, *Kibaropsis* vs. *Hedycarya*). This latter situation mirrors previous results obtained from the New Zealand flora (Garnock-Jones, 2014) and matches the expectations of Stuessy *et al.* (2014). In several cases, the evidence that an endemic genus is nested in a larger one has been known for a long time (e.g. *Oxanthera* nested in *Citrus*), but the synonymy and the necessary taxonomic changes are still pending. There are currently few cases of endemic genera that are polyphyletic (the recently described *Kanakomyrtus* and potentially *Chambeyronia* and *Kentiopsis*). Although many species in endemic genera are yet to be included in phylogenetic studies to confirm monophyly, the small size of most endemic genera makes polyphyly unlikely. In a few cases, some endemic genera may be better merged, although further evidence may still be required (e.g. *Thiollierea/Morierina*, *Chambeyronia/Actinokentia/Kentiopsis*, *Kermadecia/Sleumerodendron*). The distinctiveness and monophyly of eight genera remain to be evaluated by molecular analyses.

The flora of New Caledonia is still incompletely known as indicated by its increasing number of known endemic species over time: 2423 (Jaffré *et al.*, 2001), 2519 (Morat *et al.*, 2012) and 2544 (Munzinger *et al.*, 2016). Many important families in New Caledonia have not yet been monographed in the local flora “*Flore de la Nouvelle-Calédonie*” or elsewhere. As for other tropical areas, herbaria around the world probably house a large number of New

Caledonian species awaiting description (Bebber *et al.*, 2010) and this may include a few novel genera. Future phylogenetic works may also reveal new endemic genera to be resurrected or described, e.g. the recently segregated *Jaffrea* (Rhamnaceae) or, in Sapindaceae, the New Caledonian species of the Palaeotropical *Arytera* and *Cupaniopsis* appear to be phylogenetically distinct (Buerki *et al.*, 2012). As increasing knowledge is gathered regarding the phylogenetic relationships of the endemic genera, critical morphological comparison with their closest relatives becomes easier. Some of the monotypic genera may lose their rank to reduce redundancy similar to the expanded concepts of Strasburgeriaceae and Paracryphiaceae (Angiosperm Phylogeny Group III, 2009) that reduced the number of endemic families from five to three.

Current evidence indicates that between 62 and 91 genera endemic to New Caledonia probably satisfy the criterion of monophyly (Table 1) and this probably represents a conservative estimate of the actual number of endemic genera. These genera have on average 4.7–4.9 species, with a median of 2; 26–35 of them are monotypic and they encompass 12.0–16.7% of the endemic flora of New Caledonia.

Comparison of the New Caledonian endemic generic flora with those of other Pacific islands

New Zealand may have between 28 and 44 endemic genera after considering phylogenetic evidence (Garnock-Jones, 2014). Eastern Polynesia (French Polynesia and the Cook Islands) has 12 endemic genera according to Meyer (2004); however, the endemic *Sclerotheca* A.DC. and *Apetahia* Baill. (Campanulaceae) may be merged since *Sclerotheca* is nested in *Apetahia* (Givnish *et al.*, 2009; Pillon *et al.* unpublished) and are morphologically similar (Florence, 1997). The endemic *Tahitia* Burret (Malvaceae) may be considered as a synonym of *Christiana* DC. (Kubitzki, 1995) and *Plakothira* J.Florence (Loasaceae) is nested in *Klaprothia* Kunth (Weigend *et al.*, 2004). Samoa has only two endemic genera according

to Whistler (2011): *Sarcopygme* Setch. & Christoph. (Rubiaceae) and *Solfia* Rechinger (Arecaceae). However, recent phylogenetic evidence challenged their distinctiveness so that the first genus is now a synonym of *Morinda* L. (Razafimandimbison & Bremer, 2011) and the second is a synonym of *Balaka* Becc. (Zona & Baker, 2014). Fiji has seven endemic genera according to *Flora Vitiensis Nova* (Smith, 1996): *Amaroria* A.Gray (Simaroubaceae), *Degeneria* I.W.Bailey & A.C.Sm. (Degeneriaceae), *Gillespiea* A.C.Sm., *Hedstromia* A.C.Sm., *Squamellaria* Becc., *Sukunia* A.C.Sm. (Rubiaceae) and *Pimia* Seem. (Malvaceae). *Amaroria* is nested in *Soulamea* Lam. (Clayton *et al.*, 2007) and *Pimia* is a synonym of *Commersonia* J.R.Forst. & G.Forst. (Dorr & Cheek, 2011). *Gillespiea* has never been included in any molecular study, *Hedstromia* belongs to a large *Psychotria* L. clade the limits of which remain unclear and requires further work (Barrabé *et al.*, 2014) and *Squamellaria* is a peculiar lineage of subtribe Hydnohytinae and is now considered to be present on Vanuatu and the Solomon Islands (Chomicki & Renner, 2016). *Sukunia*, sometimes considered as a synonym of *Atractocarpus* Schltr. & K.Krause (Puttock & Quinn, 1999), was viewed as distinct more recently (Mouly *et al.*, 2014). Note that Degeneriaceae are endemic to Fiji. There is no conspectus for the flora of Vanuatu, but it seems to host only one endemic genus: *Carpoxylon* H.Wendl. & Drude (Arecaceae). We still lack a flora treatment for the Solomon islands, but Thorne (1969) listed five endemic genera in this region: *Allowoodsonia* Markgr. (Apocynaceae), *Cassidispermum* Hemsl. (Sapotaceae), *Kajewskiella* Merr. & L.M.Perry (Rubiaceae), *Rehderophoenix* Burret and *Strongylocaryum* Burret (Arecaceae). *Cassidispermum* is thought to be a synonym of *Burckella* Pierre (Govaerts *et al.*, 2015), *Rehderophoenix* is a synonym of *Drymophloeus* Zipp. and *Strongylocaryum* is a synonym of *Ptychosperma* Labill. (Govaerts *et al.*, 2016). The Hawaiian islands have 35 endemic vascular plant genera (Wagner, Herbst & Sohmer, 1999; Palmer, 2003), a number that needs to be

revised and should be considered as an upper estimate. Stuessy *et al.* (2014) suggested that only 22 may be recognized after careful examination.

Most Pacific archipelagos have only a few endemic genera, the number tending to decrease with the progress of phylogenetic analyses. Larger numbers are found in New Caledonia, New Zealand and Hawaii. Even when considering lower estimates, New Caledonia still has by far the largest number of endemic genera in the Pacific region. It also has three endemic families (and one order, Amborellales); only Fiji also has an endemic family.

Why are there so many endemic genera in New Caledonia?

New Caledonia has a rich and unique flora (Morat *et al.*, 2012) with 3389 species, of which 75.1% are endemic (Munzinger *et al.*, 2016). It can therefore also be expected to have a higher number of endemic genera. These two levels of endemism may share a number of identical causes. New Caledonia is geographically isolated, has a relatively large surface (c. 18,600 km²) and is chiefly composed of a single large island (rather than many small ones) and might therefore be expected to harbour more species than an archipelago of equivalent surface, but composed of a greater number of smaller islands (Diamond, 1975). It also lies entirely in the tropical belt, compared to, e.g., New Zealand, which is at a higher latitude and can be expected to have a comparatively less species-rich flora (Gaston, 2000). Beyond such trivial explanations, there may be other factors that have favoured a greater accumulation of endemic genera in New Caledonia than in any other Pacific islands.

Endemic genera as relicts

Endemic genera may be relicts of larger and once widespread clades, in which species from outside the island may have been driven to extinction and New Caledonia may have

offered particular conditions to accumulate many such relicts. Indeed, 26 to 35 of these genera are monotypic and most of them have few species (only eight to 12 have ten or more species), suggesting that they may be declining globally rather than actively diversifying. Several endemic genera have a sister group in remote areas (South Africa, South America, Caribbean) that do not seem to be intuitive source areas for the New Caledonian flora. Such biogeographical disjunctions may be easier to explain by extinctions in areas adjacent to New Caledonia.

Although its biodiversity may not be as old as it was previously estimated, New Caledonia is still much older than any other islands in the Pacific region besides New Zealand (Table 2). Communities on New Caledonia may have had sufficient time to reach saturation as indicated by slowing-down of diversification and density-dependent diversification observed in several groups (Espeland & Murienne, 2011; Pillon, 2012; Swenson *et al.*, 2015). There is, however, not yet any evidence from molecular dating (Pillon, 2012) or the fossil record (which is sparse and requires a critical review; Guillaumin, 1919) supporting the continuous coverage of vegetation since the emergence of the island in the Eocene. Nevertheless, rainforests may have been continuously present for a long period of time, including through the Quaternary climatic oscillations (Pintaud, Jaffré & Puig, 2001; Poncet *et al.*, 2013; Pouteau *et al.*, 2015). In contrast, nearby Australia, which probably served as a source for most of the island flora (Morat, 1993) went through dramatic climatic changes during the Tertiary, particularly aridification, that may have driven to extinction many plant lineages, most notably in the rainforest (Holloway, 1979; Byrne *et al.*, 2011; Crisp & Cook, 2013; Sniderman, Jordan & Cowling, 2013), some of them being putative close relatives of New Caledonian species. Indeed, a number of genera restricted to New Caledonia nowadays have a fossil record in Australia and/or New Zealand: *Beauprea* (Pocknall & Crosbie, 1988), *Codia* (Barnes & Hill, 1999), *Amphorogyne*, *Paracryphia* and *Phelline* (Pole, 2010). New

Caledonian conifers also suffered lower extinction rates than in other areas (Condamine, Leslie & Antonelli, 2016).

New Caledonia is well known for its ultramafic substrates that offer challenging conditions for plant growth and host the unique ‘maquis’ vegetation (Isnard *et al.*, 2016). We found that the majority of the endemic genera (59–85) are found on ultramafic substrates and 25–36 of the endemic genera are restricted to them (or almost so), whereas only three to six avoid them (Table 1). Ultramafic substrates cover about one third of New Caledonia and host about half of the indigenous flora; 36% of the species are restricted to this substrate (Isnard *et al.*, 2016). Therefore, the number of endemic genera restricted to these substrates is unexpectedly high regarding their surface and species richness. Ultramafic substrates probably covered the entire island as it emerged (Pelletier, 2006) and probably acted as ecological filter in the making of this island flora (Pillon *et al.*, 2010). This double insularity may have protected the earliest settlers by limiting the rate of immigration (Jaffré *et al.*, 1987), slowing down the turnover in this community and allowing relicts to survive. The high diversity of conifers in New Caledonia, often considered as less competitive than angiosperms, and their strong bias towards ultramafic substrates (Jaffré, Munzinger & Lowry, 2010) and the limited number of invasive species recorded on such soils (Meyer *et al.*, 2006) support this hypothesis.

Endemic genera as the product of adaptive radiations

Adaptive radiation on islands implies the diversification of one colonizer into an array of species occupying a variety of niches and differing by morphological and physiological traits associated with those niches (Schluter, 2000). Carlquist (1965: 134) cited only a few examples of adaptive radiations in New Caledonia: *Araucaria*, *Dacrydium* Lamb, *Podocarpus* L’Hér. ex Pers. (all relatively small radiations of conifers), *Phyllanthus* L. (multiple

colonization events, Kathriarachchi *et al.*, 2006) and *Tieghemopanax* R.Vig. (now included in *Polyscias* J.R.Forst. & G.Forst., see Eibl, Plunkett & Lowry, 2001). When morphological differences are sufficient, the offspring of an adaptive radiation may be treated as different genera, but this does not seem to be common in New Caledonia. There are currently only four to six pairs of sister endemic genera: *Clematepistephium* & *Eriaxis*, *Clinosperma* & *Cyphokentia*, *Amphorogyne* & *Daenikera*, *Arthroclianthus* & *Nephrodesmus* and, possibly, *Morierina* & *Thiollierea* and *Kermadecia* & *Sleumerodendron*. There is no clade of three or more endemic genera (but see *Pycnandra* below).

Eriaxis and *Clematepistephium* (both monotypic) have remarkably different habits. The first is an erect herb not exceeding 50 cm (inflorescence excluded) mostly found in maquis, whereas the second is a long rainforest vine reaching several meters (Hallé, 1977). *Clinosperma* (four species) and *Cyphokentia* (two species) differ by the presence/absence of membranous ramenta and didymous/not didymous anthers (Pintaud & Baker, 2008), but these cannot be easily connected with putative ecological differences. *Amphorogyne* spp. (three) are root parasitic chlorophyllous shrubs and trees reaching several meters, whereas *Daenikera* (monotypic) is a parasitic, aphyllous, purplish shrub not exceeding 60 cm (Hallé, 1988). *Nephrodesmus* (19 species) and *Arthroclianthus* (five species) differ by the position of the corolla wings (cover the keel in *Arthroclianthus* vs. do not cover the keel in *Nephrodesmus*), and their fruit morphology (sustained by a stipe vs. not sustained by a stipe; H. Vandrot, comm. pers.). These latter genera do not differ in terms of ecology, although *Arthroclianthus* has a greater ecological amplitude. *Morierina* (two species) and *Thiollierea* (12 species) differ in flower morphology (tubular with long and reflexed lobes vs. campanulate with short lobes), inflorescence structure (straight and subtending on a stiff axis with leafy bracts vs. pendant and generally on a flexible axis without leafy bracts) and type of anthers (not twisted vs. twisted, Barrabé *et al.*, 2011), suggesting a shift in pollination syndrome (moths vs. birds,

Motley, Wurdack & Delprete, 2005). The first is mostly found in rainforests, whereas the second is generally in maquis; both are restricted to ultramafic substrates. *Kermadecia* (four species) and *Sleumerodendron* (one species) and *Turrillia* (Vanuatu & Fiji) are similar vegetatively and ecologically and their flowers differ only in small points (Virot, 1967; Smith & Haas, 1975).

These pairs of sister genera account only for a small number of species (24 at most for *Arthroclianthus-Nephrodesmus*). The morphological and ecological differences are relatively modest in three cases (*Clinosperma & Cyphokentia*, *Arthroclianthus & Nephrodesmus*, *Kermadecia & Sleumerodendron*) and do not obviously qualify as adaptive radiations. The fact that two are relatively species-poor (*Clematapistephium & Eriaxis*, *Amphorogyne & Daenikera*) suggests that they are relicts rather than actively diversifying lineages. This contrasts with Hawaii that has several multi-generic plant radiations: the silversword alliance (three genera, 28 species; Baldwin, 1997), lobeliads (six genera, 126 species; Givnish *et al.*, 2009) and mints (three genera, 60 species, Lindqvist & Albert, 2002). The Macaronesian islands have a large radiation of Crassulaceae (four genera, 63 species; Mort *et al.*, 2002). Adaptive radiations seem to have contributed little to the formation of endemic genera in New Caledonia. The largest monophyletic clade in New Caledonia is a group of *Psychotria* (c. 78 species, Barrabé *et al.*, 2014) with modest morphological diversity that is not easily correlated with the ecology of the different species. The largest endemic genus, *Pycnandra*, is composed of six subgenera (Swenson *et al.*, 2015), most of which were once recognized as genera. However, apart from subgenus *Lepostylis*, the delimitation of these subgenera has changed considerably since the time when they were ranked as genera (Swenson *et al.*, 2008). The homoplasy of most morphological characters explored so far (Swenson *et al.*, 2015) probably made the classification of this group complicated. Since the subgenera of *Pycnandra* are difficult to characterize morphologically (Swenson & Munzinger, 2009, 2016; Swenson *et al.*,

2015) and do not show clear ecological differences, *Pycnandra* does not seem to be comparable to the multigeneric adaptive radiations from Hawaii or the Macaronesian islands cited above. With the current evidence at hand, most New Caledonian plant lineages may be considered as mere “radiation” or, in a few cases, “explosive radiation” (e.g. *Psychotria*) rather than “adaptive radiation” (Givnish, 2015).

It remains unclear why the New Caledonian flora has few dramatic examples of adaptive radiation, but it may be linked to the fact that the vegetation is largely composed of woody plant species (Virot, 1956), whereas most convincing examples of adaptive radiation in plants are herbaceous or ancestrally herbaceous (Givnish, 2010). “Increased woodiness” is a commonly observed syndrome on islands, whereby herbaceous ancestors become shrubs or trees after colonizing islands [Carlquist, 1965 (chapter 8), 1974 (chapter 10)], but this is almost unseen in New Caledonia and the rest of Melanesia (Carlquist, 1974: 409). Many large herbaceous families are under-represented in New Caledonia, including Asteraceae, Campanulaceae, Cyperaceae, Lamiaceae and Poaceae (Pillon *et al.*, 2010). The paucity of Asteraceae is particularly striking, especially on the ultramafic soils (Carlquist, 1965: 42), whereas they are well represented in many island floras, producing several remarkable adaptive radiations (Carlquist, 1965: 204). When New Caledonia emerged, it was probably entirely covered with ultramafic substrate which may have filtered the successful immigrants (Pillon *et al.*, 2010) and primarily woody plants may be more competitive than herbs on ultramafic substrates.

The relative ancient age of New Caledonia may also explain the lack of dramatic adaptive radiation by allowing enough time for a diverse cohort of immigrants to reach the island. The diversity of Asteraceae on islands may be explained by their ability to disperse long distances and their weediness and evolutionary lability (Carlquist, 1965: 204), whereas trees may be poor dispersers (Carlquist, 1965: 189). As time passes, even the least likely

dispersal might happen and, for example, new immigrant tree species may outcompete those that evolved from herbaceous colonizers. The available data from molecular dating suggest that arrivals of colonizers were spread in time rather than being synchronous and close to the putative emergence time of New Caledonia (Pillon, 2012). On New Caledonia, we may be observing a mature flora, in which continuous immigration has progressively thinned out the earlier adaptive radiations. Such processes have not yet taken place in the other more recent Pacific islands.

Despite its complex environmental gradients, New Caledonia may not offer ecological conditions as diverse as other islands, particularly in terms of climate. Hawaiian islands display a diverse range of climatic conditions (Carlquist, 1980: 63), some extreme, offering considerable opportunities for adaptive radiations. New Caledonia lies entirely in the tropical belt and its highest peak (Mt Panié, 1628 m) is lower than that of several Pacific archipelagos (Table 2). Although its mountain tops host some distinctive plant communities that may be exposed to frost (Bradford & Jaffré, 2004; Munzinger, McPherson & Lowry, 2008), those are still dense and largely shrubby and do not display distinctive vegetative forms such as large rosette and cushion plants.

It has long been recognized that a striking ecological feature of New Caledonia is the diversity of its soils (Jaffré, 1993) and this could have been a major driver for plant diversification (Pillon *et al.*, 2009a,b). However this might have been translated into physiological diversity more than morphological attributes and hence a rather cryptic mode of adaptive radiation (e.g. Pillon *et al.*, 2014).

Finally, New Caledonia also displays few dramatic examples of plant-pollinator co-evolution that could compare to, for example, Hawaiian lobeliads and Hawaiian honeycreepers (Carlquist, 1965: 131). There is an interesting case of mutualism reported between the large genus *Phyllanthus* and *Epicephala* moths (Kawakita & Kato, 2004), but it

did not result in spectacular variations in flower morphology. Pollination syndromes have not been thoroughly investigated in New Caledonia (but see Carpenter, Read & Jaffré, 2003; Kato & Kawakita, 2004; Donovan *et al.*, 2013; Hopkins *et al.*, 2015b) and some patterns might have been overlooked. It is possible that the pollination guild of New Caledonia is more depauperate than elsewhere (Carpenter *et al.*, 2003; Pauly & Munzinger, 2003). Also, most of the lineages in which zygomorphic flowers are found are under-represented in the flora of New Caledonia: monocots, Asterales, Fabales and Lamiales (Pillon *et al.*, 2010), and this may have reduced the opportunities for specific plant-pollinator interactions.

CONCLUSIONS

New Caledonia may have a total of between 62 and 91 endemic genera. Many of these genera have still not been included in molecular phylogenetic studies and the delimitation of others needs further work, and some other genera may be revealed in the future. This number of endemic genera is far greater than in any other Pacific archipelago. Endemic genera may be the result of extinctions outside the island, evolutionary changes on the island or both. In the case of New Caledonia, the first phenomenon may have been the preponderant force. The relative ancient age of the island, a putative climatic stability and a possible protective role of ultramafic substrate against new colonizers may have been key factors in the accumulation of relict lineages. In contrast, evidence for adaptive radiations is limited and this may be linked to the modest contribution of primarily herbaceous lineages to the composition of flora or the limited ecological contrasts on the island beside the diversity of soils. Beyond the now famous *Amborella*, New Caledonia hosts a large number of unique relicts of conservation significance, as can be expected in old, climatically buffered, infertile landscapes (OCBILs, Hopper, 2009; Hopper, Silveira & Fiedler, 2016). Endemic genera may be in some cases the sole survivor of ancient lineages and the age and phylogenetic diversity of these taxa will be

discussed in a forthcoming paper. It can be anticipated that this will magnify the conservation significance of the ultramafic substrates of New Caledonia, which host most endemic genera and where the preservation of the native ecosystems is most conflicted (Pascal *et al.*, 2008). Updated conservation assessments for these unique lineages are urgently needed to set conservation priorities and preserve these unique branches of the tree of life.

ACKNOWLEDGEMENTS

We thank Bill Baker, Mark Chase, Ashley Egan, Hervé Jourdan, Eve Lucas, Jean-Yves Meyer, Arnaud Mouly, Laurence Ramon, Tim Utteridge, Hervé Vandrot and Thais Vasconcelos for comments or for sharing unpublished results with us, Vanessa Hequet for providing a picture of *Loxodiscus*, and Carlos García-Verdugo, Phil Garnock-Jones and an anonymous reviewer for insightful reviews on a previous version of this manuscript.

REFERENCES

- Achille F, Motley TJ, Lowry PP, Jérémie J. 2006.** Polyphyly in *Guettarda* L. (Rubiaceae, Guettardeae) based on nrDNA ITS sequence data. *Annals of the Missouri Botanical Garden* **93**: 103–121.
- Alford MA. 2005.** *Systematics studies in Flacourtiaceae*. PhD thesis. Cornell University.
- Alvarez-Molina A, Cameron KM. 2009.** Molecular phylogenetics of Prescotttiinae *s.l.* and their close allies (Orchidaceae, Cranichideae) inferred from plastid and nuclear ribosomal DNA sequences. *American Journal of Botany* **96**: 1020–1040.
- Amborella Genome Project. 2013.** The *Amborella* genome and the evolution of flowering plants. *Science* **342**: 1241089.
- Anderson CL, Rova JHE, Andersson L. 2001.** Molecular phylogeny of the tribe Anthospermeae (Rubiaceae): systematic and biogeographic implications. *Australian Systematic Botany* **14**: 231-244.
- APG III. 2009.** An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society* **161**: 105–121.

- APG IV. 2016.** An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* **181**: 1–20.
- Appelhans MS, Wen J, Wagner WL. 2014.** A molecular phylogeny of *Acronychia*, *Euodia*, *Melicope* and relatives (Rutaceae) reveals polyphyletic genera and key innovations for species richness. *Molecular Phylogenetics and Evolution* **79**: 54–68.
- Aubréville A. 1967.** Sapotacées. In: Aubréville A, ed. *Flore de la Nouvelle-Calédonie et Dépendances*. Paris: Muséum National d'Histoire Naturelle, 1–168.
- Bacon CD, Baker WJ. 2011.** *Saribus* resurrected. *Palms* **55**: 109–116.
- Bacon CD, Baker WJ, Simmons MP. 2012.** Miocene dispersal drives island radiations in the palm tribe Trachycarpeae (Arecaceae). *Systematic Biology* **61**: 426–442.
- Baker WJ, Savolainen V, Asmussen-Lange CB, Chase MW, Dransfield J, Forest F, Harley MM, Uhl NW, Wilkinson M. 2009.** Complete generic-level phylogenetic analyses of Palms (Arecaceae) with comparisons of supertree and supermatrix approach. *Systematic Biology* **58**: 240–256.
- Baker WJ, Norup MV, Clarkson JJ, Couvreur TLP, Dowe JL, Lewis CE, Pintaud JC, Savolainen V, Wilmot T, Chase MW. 2011.** Phylogenetic relationships among arecoid palms (Arecaceae: Arecoideae). *Annals of Botany* **108**: 1417–1432.
- Baldwin BG. 1997.** Adaptive radiation of the Hawaiian silversword alliance: congruence and conflict of phylogenetic evidence from molecular and non-molecular investigations. In: Givnish TJ, Sytsma KJ, eds. *Molecular evolution and adaptive radiation*. Cambridge: Cambridge University Press, 103–128.
- Barnes RW, Hill RS. 1999.** Macrofossils of *Callicoma* and *Codia* (Cunoniaceae) from Australian Cainozoic sediments. *Australian Systematic Botany* **12**: 647–670.
- Barrabé L, Mouly A, Lowry PP, Munzinger J. 2011.** Reinstatement of the endemic New Caledonian genus *Thiollierea* Montrouz. (Rubiaceae) necessitated by the polyphyly of *Bikkia* Reinw. as currently circumscribed. *Adansonia* **33**: 115–134.
- Barrabé L, Maggia L, Pillon Y, Rigault F, Mouly A, Davis AP, Buerki S. 2014.** New Caledonian lineages of *Psychotria* (Rubiaceae) reveal different evolutionary histories and the largest documented plant radiation for the archipelago. *Molecular Phylogenetics and Evolution* **71**: 15–35.
- Barraclough TG, Humphreys AM. 2015.** The evolutionary reality of species and higher taxa in plants: a survey of post-modern opinion and evidence. *New Phytologist* **207**: 291–296.
- Bartish IV, Swenson U, Munzinger J, Anderberg AA. 2005.** Phylogenetic relationships among New Caledonian Sapotaceae (Ericales): molecular evidence for generic polyphyly and repeated dispersal. *American Journal of Botany* **92**: 667–673.
- Bayer RJ, Mabberley DJ, Morton CM, Miller CH, Sharma IK, Pfeil BE, Rich S, Hitchcock R, Sykes S. 2009.** A molecular phylogeny of the orange subfamily (Rutaceae: Aurantioideae) using nine cpDNA sequences. *American Journal of Botany* **96**: 668–685.

- Bayly MJ, Holmes GD, Forster PI, Cantrill DJ, Ladiges PY. 2013.** Major clades of Australasian Rutoideae (Rutaceae) Based on *rbcL* and *atpB* sequences. *PloS one* **8**: e72493.
- Bayly MJ, Duretto MF, Holmes GD, Forster PI, Cantrill DJ, Ladiges PY. 2015.** Transfer of the New Caledonian genus *Boronella* to *Boronia* (Rutaceae) based on analyses of cpDNA and nrDNA. *Australian Systematic Botany* **28**: 111-123.
- Beaumont AJ, Edwards TJ, Manning J, Maurin O, Rautenbach M, Motsi MC, Fay MF, Chase MW, van der Bank M. 2009.** *Gnidia* (Thymelaeaceae) is not monophyletic: taxonomic implications for Thymelaeoideae and a partial new generic taxonomy for *Gnidia*. *Botanical Journal of the Linnean Society* **160**: 402–417.
- Bebber DP, Carine MA, Wood JRI, Wortley AH, Harris DJ, Prance GT, Davidse G, Paige J, Pennington TD, Robson NKB, Scotland RW. 2010.** Herbaria are a major frontier for species discovery. *Proceedings of the National Academy of Science of the USA* **107**: 22169–22171.
- Biffin E, Lucas EJ, Craven LA, da Costa IR, Harrington MG, Crisp MD. 2010.** Evolution of exceptional species richness among lineages of fleshy-fruited Myrtaceae. *Annals of Botany* **106**: 79–93.
- Bradford JC. 2002.** Molecular phylogenetics and morphological evolution in Cunoniaceae (Cunoniaceae). *Annals of the Missouri Botanical Garden* **89**: 491–503.
- Bradford JC, Jaffré T. 2004.** Plant species microendemism and conservation of montane maquis in New Caledonia: two new species of *Pancheria* (Cunoniaceae) from the Roche Ouaième. *Biodiversity and Conservation* **13**: 2253–2273.
- Breitwieser I, Brownsey PJ, Garnock-Jones PJ, Perrie LR, Wilton A. 2012.** Phylum Tracheophyta, vascular plants. In: Gordon D, ed. *The New Zealand Inventory of Biodiversity: a Species 2000 Symposium Review*. Christchurch: University of Canterbury Press, 411–459.
- Bremer B, Bremer K, Heidari N, Erixon P, Olmstead RG, Anderberg AA, Källersjö M, Barkhordarian E. 2002.** Phylogenetics of asterids based on 3 coding and 3 non-coding chloroplast DNA markers and the utility of non-coding DNA at higher taxonomic levels. *Molecular Phylogenetics and Evolution* **24**: 274–301.
- Buerki S, Forest F, Acevedo-Rodriguez P, Callmender MW, Nylander JAA, Harrington M, Sanmartin I, Küpfer P, Alvarez N. 2009.** Plastid and nuclear DNA markers reveal intricate relationships at subfamilial and tribal levels in the soapberry family (Sapindaceae). *Molecular Phylogenetics and Evolution* **51**: 238–258.
- Buerki S, Forest F, Callmender MW, Lowry PP, Devey DS, Munzinger J. 2012.** Phylogenetic inference of New Caledonian lineages of Sapindaceae: molecular evidence requires a reassessment of generic circumscriptions. *Taxon* **61**: 109–119.
- Buerki S, Devey DS, Callmender MW, Phillipson PB, Forest F. 2013.** Spatio-temporal history of the endemic genera of Madagascar. *Botanical Journal of the Linnean Society* **171**: 304–329.

Buerki S, Callmander MW, Bachman S, Moat J, Labat JN, Forest F. 2015. Incorporating evolutionary history into conservation planning in biodiversity hotspots. *Philosophical Transactions of the Royal Society B-Biological Sciences* **370**: 20140014.

Byng JW. 2015. *The gymnosperms handbook: a practical guide to extant families and genera of the world*. Hertford: Plant Gateway Ltd.

Byng JW, Bernardini B, Joseph JA, Chase MW, Utteridge TMA. 2014. Phylogenetic relationships of Icacinaceae focusing on the vining genera. *Botanical Journal of the Linnean Society* **176**: 277–294.

Byrne M, Steane DA, Joseph L, Keates DK, Jordan GJ, Crayn D, Aplin K, Cantrill DJ, Cook LG, Crisp MD, Keogh JS, Melville J, Moritz C, Porch N, Sniderman JM, Sunnucks P, Weston PH. 2011. Decline of a biome: evolution, contraction, fragmentation, extinction and invasion of the Australian mesic zone biota. *Journal of Biogeography* **38**: 1635–1656.

Cameron KM. 2003. On the phylogenetic position of the New Caledonian endemic families Paracryphiaceae, Oncothecaceae, and Strasburgeriaceae: a comparison of molecules and morphology. *The Botanical Review* **68**: 428–443.

Cameron KM. 2009. On the value of nuclear and mitochondrial gene sequences for reconstructing the phylogeny of vanilloid orchids (Vanilloideae, Orchidaceae). *Annals of Botany* **104**: 377–385.

Cantley JT, Swenson NG, Markey A, Keeley SC. 2014. Biogeographic insights on Pacific *Coprosma* (Rubiaceae) indicate two colonizations to the Hawaiian Islands. *Botanical Journal of the Linnean Society* **174**: 412–424.

Carlquist S. 1965. *Island life*. New York: American Museum of Natural History.

Carlquist S. 1974. *Island biology*. New York: Columbia University Press.

Carlquist S. 1980. *Hawaii a natural history*. Lawai: National Tropical Botanical Garden.

Carpenter RJ, Read J, Jaffré T. 2003. Reproductive traits of tropical rain-forest trees in New Caledonia. *Journal of Tropical Ecology* **19**: 351–365.

Chanderbali AS, van der Werff H, Renner SS. 2001. Phylogeny and historical biogeography of Lauraceae: evidence from the chloroplast and nuclear genomes. *Annals of the Missouri Botanical Garden* **88**: 104–134.

Chase MW, Cameron KM, Freudenstein JV, Pridgeon AM, Salazar G, van der Berg C, Schuiteman A. 2015. An updated classification of Orchidaceae. *Botanical Journal of the Linnean Society* **177**: 151–174.

Cheng Y, Nicolson RG, Tripp K, Chaw SM. 2000. Phylogeny of Taxaceae and Cephalotaxaceae genera inferred from chloroplast *matK* gene and nuclear rDNA ITS region. *Molecular Phylogenetics and Evolution* **14**: 353–365.

Chokthaweeapanich H. 2014. *Phylogenetics and evolution of the Paleotropical woody bamboos (Poaceae: Bambusoideae: Bambuseae)*. PhD thesis. Iowa State University.

- Chomicki G, Renner SS. 2016.** Evolutionary relationships and biogeography of the ant-epiphytic genus *Squamellaria* (Rubiaceae: Psychotrieae) and their taxonomic implications. *PloS one* **11**: e0151317.
- Cisternas MA, Salazar GA, Verdugo G, Novoa P, Calderon X, Negritto MA. 2012.** Phylogenetic analysis of Chloraeinae (Orchidaceae) based on plastid and nuclear DNA sequences. *Botanical Journal of the Linnean Society* **168**: 258–277.
- Clayton JW, Fernando ES, Soltis PS, Soltis DE. 2007.** Molecular phylogeny of the tree-of-heaven family (Simaroubaceae) based on chloroplast and nuclear markers. *International Journal of Plant Science* **168**: 1325–1339.
- Cluzel D, Chiron MD, Courme MD. 1998.** Upper Eocene unconformity and pre-obduction events in New Caledonia. *Comptes-Rendus de l'Académie des Sciences. Science de la terre et des planètes* **327**: 485–491.
- Condamine FL, Leslie AB, Antonelli A. 2016.** Ancient islands acted as refugia and pumps for conifer diversity. *Cladistics*: in press.
- Coughenour JM, Simmons MP, Lombardi JA, Cappa JJ. 2010.** Phylogeny of Celastraceae subfamily Salacioideae and tribe Lophopetaleae inferred from morphological characters and nuclear and plastid genes. *Systematic Botany* **35**: 358–367.
- Crisp MD, Cook LG. 2013.** How was the Australian flora assembled over the last 65 million years? A molecular phylogenetic perspective. *Annual Review of Ecology, Evolution, and Systematics, Vol 44* **44**: 303–324.
- Der JP, Nickrent DL. 2008.** A molecular phylogeny of Santalaceae (Santalales). *Systematic Botany* **33**: 107–116.
- Diamond JM. 1975.** The island dilemma: lessons of modern biogeographic studies for the design of natural reserves. *Biological Conservation* **7**: 129–146.
- Domenech B, Asmussen-Lange CB, Baker WJ, Alapetite E, Pintaud JC, Nadot S. 2014.** A phylogenetic analysis of palm subtribe Archontophoenicinae (Arecaceae) based on 14 DNA regions. *Botanical Journal of the Linnean Society* **175**: 469–481.
- Donovan BJ, Munzinger J, Pauly A, McPherson G. 2013.** Flower-visiting records of the native bees of New Caledonia. *Annals of the Missouri Botanical Garden* **99**: 19–43.
- Dorr LJ, Cheek MR. 2011.** The identity and typification of *Pimia* Seem. (Sterculiaceae). *Kew Bulletin* **66**: 629–631.
- Egan AN, Vatanparast M, Cagle W. 2016.** Parsing polyphyletic *Pueraria*: delimiting distinct evolutionary lineages through phylogeny. *Molecular Phylogenetics and Evolution*: **104**: 44–59.
- Eibl JM, Plunkett GM, Lowry PP. 2001.** Evolution of *Polyscias* sect. *Tieghemopanax* (Araliaceae) based on nuclear and chloroplast DNA sequence data. *Adansonia* **23**: 23–48.
- Espeland M, Muriene J. 2011.** Diversity dynamics in New Caledonia: towards the end of the museum model? *BMC Evolutionary Biology* **11**: 254.

- Ewart A. 1988.** Geological history of the Fiji-Tonga-Samoan region of the SW Pacific, and some palaeogeographic and biogeographic implications. In: Duffels JP, ed. *The cicadas of the Fiji, Samoa and Tonga Islands, their taxonomy and biogeography*. (Homoptera, Cicadoidea). Leiden: Scandinavian Science Press, 15–23.
- Feild TS, Brodribb TJ. 2005.** A unique mode of parasitism in the conifer coral tree *Parasitaxus ustus* (Podocarpaceae). *Plant, Cell and Environment* **28**: 1316–1325.
- Fijridiyanto IA, Murakami N. 2009.** Phylogeny of *Litsea* and related genera (Laureae-Lauraceae) based on analysis of *rpb2* gene sequences. *Journal of Plant Research* **122**: 283–298.
- Florence J. 1997.** New species of *Plakothira* (Loasaceae), *Melicope* (Rutaceae), and *Apetahia* (Campanulaceae) from the Marquesas Islands. *Allertonia* **7**: 238–253.
- Garnock-Jones PJ. 2014.** Evidence-based review of the taxonomic status of New Zealand's endemic seed plant genera. *New Zealand Journal of Botany* **52**: 163–212.
- Gaston KJ. 2000.** Global patterns in biodiversity. *Nature* **405**: 220–227.
- Givnish TJ, William KC, Mast AR, Paterson TB, Theim TJ, Hipp AL, Henss JM, Smith JF, Wood KR, Sytsma KJ. 2009.** Origin, adaptive radiation and diversification of the Hawaiian lobeliads (Asterales: Campanulaceae). *Proceedings of the Royal Society of London B* **276**: 407–416.
- Givnish TJ. 2010.** Ecology of plant speciation. *Taxon* **59**: 1326–1366.
- Givnish TJ. 2015.** Adaptive radiation versus ‘radiation’ and ‘explosive diversification’: why conceptual distinctions are fundamental to understanding evolution. *New Phytologist* **207**: 297–303.
- González F, Betancur J, Maurin O, Freudenstein JV, Chase MW. 2007.** Metteniusaceae, an early-diverging family in the lamiid clade. *Taxon* **56**: 795–800.
- Goremykin V, Hirsch-Ernst KI, Wölfl S, Hellwig FH. 2003.** Analysis of the *Amborella trichopoda* chloroplast genome sequence suggests that *Amborella* is not a basal angiosperm. *Molecular Biology and Evolution* **20**: 1499–1505.
- Govaerts R, Harvey Y, Jessup L, Pennington TD, Wink W 2015.** *World checklist of Sapotaceae*. Kew: Royal Botanic Gardens. <http://apps.kew.org/wcsp/>.
- Govaerts R, Dransfield J, Zona SF, Hodel DR, Henderson A. 2016.** *World checklist of Areaceae*. Facilitated by the Royal Botanic Gardens, Kew. <http://apps.kew.org/wcsp/>.
- Grandcolas P, Murienne J, Robillard T, Desutter-Grandcolas L, Jourdan H, Guilbert E, Deharveng L. 2008.** New Caledonia: a very old Darwinian island? *Philosophical Transactions of the Royal Society of London B* **363**: 3309–3317.
- Grandcolas P, Nattier R, Trewick S. 2014.** Relict species: a relict concept? *Trends in Ecology & Evolution* **29**: 655–663.

- Guillaumin A. 1919.** Notes de paléobotanique néo-calédonienne. *Revue générale de botanique* **31**: 273–276.
- Gustafson R, Herbst DR & Rundel PW. 2014.** *Hawaiian plant life: vegetation and flora*. Honolulu: University of Hawai‘i Press.
- Hallé N. 1977.** Orchidacées. In: Aubréville A, Leroy JF, eds. *Flore de la Nouvelle-Calédonie et dépendances*. Paris: Muséum National d’Histoire Naturelle, 1–565.
- Hallé N. 1988.** Santalaceae. In: Morat P, MacKee HS, eds. *Flore de la Nouvelle-Calédonie et dépendances*. Paris: Muséum National d’Histoire Naturelle, 99–152.
- He T, Lamont BB & Fogliani B. 2016.** Pre-Gondwanan-breakup origin of *Beauprea* (Proteaceae) explains its historical presence in New Caledonia and New Zealand. *Science Advances* **2**: e1501648–e1501648.
- Herbert J, Chase MW, Möller M, Abbott RJ. 2006.** Nuclear and plastid DNA sequences confirm the placement of the enigmatic *Canacomyrica monticola* in Myricaceae. *Taxon* **55**: 349–357.
- Holloway JD. 1979.** *A survey of the Lepidoptera, biogeography, and ecology of New Caledonia*. The Hague, Boston: W. Junk.
- Hopkins HCF, Rozefelds AC, Pillon Y. 2013.** *Karrabina* gen. nov (Cunoniaceae), for the Australian species previously placed in *Geissois*, and a synopsis of genera in the tribe Geissoieae. *Australian Systematic Botany* **26**: 167–185.
- Hopkins HCF, Pillon Y, Hoogland RD. 2014.** *Cunoniaceae: Flore de la Nouvelle-Calédonie, volume 26*. Paris: Publications scientifiques du Muséum; Marseille: IRD.
- Hopkins HCF, Pillon Y, Stacy EA, Kellermann J. 2015a.** *Jaffrea*, a new genus of Rhamnaceae endemic to New Caledonia, with notes on *Alphitonia* and *Emmenosperma*. *Kew Bulletin* **70**: 42.
- Hopkins HCF, Bradford JC, Donovan B, Pillon Y, Munzinger J, Fogliani B. 2015b.** Floral biology of the Cunoniaceae in New Caledonia and the role of insects, birds and geckos as potential pollinators. *Kew Bulletin* **70**: 8.
- Hopper SD. 2009.** OCBIL theory: towards an integrated understanding of the evolution, ecology and conservation of biodiversity on old-climatically buffered, infertile landscapes. *Plant and Soil* **322**: 49–86.
- Hopper SD, Silveira FAO, Fiedler PL. 2016.** Biodiversity hotspots and OCBIL theory. *Plant and Soil* **403**: 167–216.
- Horn JW, van Ee BW, Morawetz JJ, Riina R, Steinmann VW, Berry PE, Wurdack KJ. 2012.** Phylogenetics and the evolution of major structural characters in the giant genus *Euphorbia* L. (Euphorbiaceae). *Molecular Phylogenetics and Evolution* **63**: 305–326.
- Isnard S, L’huillier L, Rigault F, Jaffré T. 2016.** How did the ultramafic soils shape the flora of the New Caledonian hotspot? *Plant and Soil* **403**: 53–76.

- Jaffré T. 1980.** *Etude écologique du peuplement végétal des sols dérivés de roches ultrabasiqes en Nouvelle-Calédonie.* Paris: ORSTOM.
- Jaffré T, Morat P, Veillon JM, MacKee HS. 1987.** Changements dans la végétation de la Nouvelle-Calédonie au cours du Tertiaire: la végétation et la flore des roches ultrabasiqes. *Adansonia* **4**: 365–391.
- Jaffré T. 1993.** The relationship between ecological diversity and floristic diversity in New Caledonia. *Biodiversity Letters* **1**: 82–87.
- Jaffré T, Morat P, Veillon JM, Rigault F, Dagostini G. 2001.** *Composition and characterisation of the native flora of New Caledonia.* Nouméa: Institut de Recherche pour le Développement.
- Jaffré T, Munzinger J, Lowry PP. 2010.** Threats to the conifer species found on New Caledonia's ultramafic massifs and proposals for urgently needed measures to improve their protection. *Biodiversity and Conservation* **19**: 1485–1502.
- Kårehed J, Lundberg J, Bremer B, Bremer K. 1999.** Evolution of the Australasian families Alseuosmiaceae, Argophyllaceae, and Phellinaceae. *Systematic Botany* **24**: 660–682.
- Kathriarachchi H, Samuel R, Hoffmann P, Mlinarec J, Wurdack KJ, Ralimanana, Stuessy TF, Chase MW. 2006.** Phylogenetics of tribe Phyllanthae (Phyllanthaceae; Euphorbiaceae *sensu lato*) based on nrITS and plastid *matK* DNA sequence data. *American Journal of Botany* **93**: 637–655.
- Kato M & Kawakita A. 2004.** Plant-pollinator interactions in New Caledonia influenced by introduced honey bees. *American Journal of Botany* **91**: 1814–1827.
- Kawakita A & Kato M. 2004.** Evolution of obligate pollination mutualism in New Caledonian *Phyllanthus* (Euphorbiaceae). *American Journal of Botany* **91**: 410–415.
- Kores PJ, Molvray M, Weston PH, Hopper SD, Brown AP, Cameron KM, Chase MW. 2001.** A phylogenetic analysis of Diurideae (Orchidaceae) based on plastid DNA sequence data. *American Journal of Botany* **88**: 1903–1914.
- Kostermans AJGH. 1974.** Lauracées. In: Aubréville A, Leroy JF, eds. *Flore de la Nouvelle-Calédonie et dépendances.* Paris: Muséum National d'Histoire Naturelle, 1–123.
- Kranitz ML, Biffin E, Clark A, Hollingsworth ML, Ruhsam M, Gardner MF, Thomas P, Mill RR, Ennos RA, Gaudeul M, Lowe, AJ, Hollingsworth PM. 2014.** Evolutionary diversification of New Caledonian *Araucaria*. *PloS one* **9**: e110308.
- Kubitzki K. 1995.** *Asterophorum* and *Tahitia* congeneric with *Christiana*. *Botanische Jahrbücher für Systematik Pflanzengeschichte und Pflanzengeographie* **116**: 537–542.
- Lehtonen S. 2011.** Towards resolving the complete fern Tree of Life. *PLoS one* **6**: e24851.
- Lescot M. 1980.** Flacourtiacées. In: Aubréville A, Leroy JF, eds. *Flore de la Nouvelle-Calédonie et Dépendances.* Paris: Muséum National d'Histoire Naturelle, 3–134.

- Lindqvist C, Albert VA. 2002.** Origin of the Hawaiian endemic mints within North American *Stachys* (Lamiaceae). *American Journal of Botany* **89**: 1709–1724.
- Livshultz T, Middleton DJ, Endress ME, Williams JK. 2007.** Phylogeny of Apocynoideae and the APSA clade (Apocynaceae *s.l.*). *Annals of the Missouri Botanical Garden* **94**: 324–359.
- Lu Y, Ran JH, Guo DM, Yang ZY, Wang XQ. 2014.** Phylogeny and divergence times of gymnosperms inferred from single-copy nuclear genes. *PLoS one* **9**: e107679.
- Lundberg J, Bremer K. 2003.** A phylogenetic study of the order Asterales using one morphological and three molecular data sets. *International Journal of Plant Science* **164**: 553–578.
- Mace GM, Gittleman JL, Purvis A. 2003.** Preserving the tree of life. *Science* **300**: 1707–1709.
- Manns U, Bremer B. 2010.** Towards a better understanding of intertribal relationships and stable tribal delimitations within Cinchonoideae *s.s.* (Rubiaceae). *Molecular Phylogenetics and Evolution* **56**: 21–39.
- Markey AS, Lord JM, Orlovich DA. 2004.** *Coprosma talbrockiei*: an oddball sheds light on the Coprosminae. Abstract of paper presented at SYSTANZ meeting, Whakapapa Village, 2004.
- Mast AR, Willis CL, Jones EH, Downs KM, Weston PH. 2008.** A smaller *Macadamia* from a more vagile tribe: inference of phylogenetic relationships, divergence times, and diaspore evolution in *Macadamia* and relatives (tribe Macadamieae; Proteaceae). *American Journal of Botany* **95**: 843–870.
- Maury R, Legendre C, Chauvel C, Guille G, Blais S, Guillou H, Rossi P. 2014.** An atypical hotspot chain. In: Meyer JM, Claridge E, eds. *Terrestrial biodiversity of the Austral Islands, French Polynesia*. Paris: Muséum National d’Histoire naturelle, 21–37.
- McPherson G, Lowry PP. 2004.** *Hooglandia*, a newly discovered genus of Cunoniaceae from New Caledonia. *Annals of the Missouri Botanical Garden* **91**: 260–265.
- Mennes CB, Lam VKY, Rudall PJ, Lyon SP, Graham SW, Smets EF, Mercks VSFT. 2015.** Ancient Gondwana break-up explains the distribution of the mycoheterotrophic family Corsiaceae (Liliales). *Journal of Biogeography* **42**: 1123–1136.
- Meyer JY. 2004.** Threat of invasive alien plants to native flora and forest vegetation of eastern Polynesia. *Pacific Science* **58**: 357–375.
- Meyer JY, Loope LL, Sheppard A, Munzinger J, Jaffré T. 2006.** Les plantes envahissantes et potentiellement envahissantes dans l’archipel néo-calédonien : première évaluation et recommandations de gestion. In: Beauvais ML, Coleno A, Jourdan H, eds. *Les espèces envahissantes dans l’archipel néo-calédonien, un risque environnemental et économique majeur*. Paris: IRD éditions.
- Morat P. 1993.** Our knowledge of the flora of New Caledonia: endemism and diversity in relation to vegetation types and substrates. *Biodiversity Letters* **1**: 72–81.

- Morat P, Jaffré T, Tronchet F, Munzinger J, Pillon Y, Veillon JM, Chalopin M. 2012.** The taxonomic reference base Florical and characteristics of the native vascular flora of New Caledonia. *Adansonia* **34**: 179–221.
- Mort ME, Soltis DE, Soltis PS, Francisco-Ortega J, Santos-Guerra A. 2002.** Phylogenetics and evolution of the Macaronesian clade of Crassulaceae inferred from nuclear and chloroplast sequence data. *Systematic Botany* **27**: 271–288.
- Motley TJ, Wurdack KJ, Delprete PG. 2005.** Molecular systematics of the Catesbaeeae-Chiococceae complex (Rubiaceae): flower and fruit evolution and biogeographic implications. *American Journal of Botany* **92**: 316–329.
- Mouly A, Kainulainen K, Persson C, Davis AP, Wong KM, Razafimandimbison SG, Bremer B. 2014.** Phylogenetic structure and clade circumscriptions in the Gardenieae complex (Rubiaceae). *Taxon* **63**: 801–818.
- Munzinger J, Morat P, Jaffré T, Gâteblé G, Pillon Y, Tronchet F, Veillon JM, Chalopin M. 2016.** FLORICAL: checklist of the vascular indigenous flora of New Caledonia vers. 22.IV.2016. <http://www.botanique.nc/herbier/florical>.
- Munzinger J, McPherson G, Lowry PP. 2008.** A second species in the endemic New Caledonian genus *Gastrolepis* (Stemonuraceae) and its implications for the conservation status of high-altitude maquis vegetation: coherent application of the IUCN Red List criteria is urgently needed in New Caledonia. *Botanical Journal of the Linnean Society* **157**: 775–783.
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J. 2000.** Biodiversity hotspots for conservation priorities. *Nature* **403**: 853–858.
- Nicolas AN, Plunkett GM. 2009.** The demise of subfamily Hydrocotyloideae (Apiaceae) and the re-alignment of its genera across the entire order Apiales. *Molecular Phylogenetics and Evolution* **53**: 134–151.
- Ormerod P, Cribb PJ. 2003.** *Danhatchia*. In: Pridgeon AM, Cribb PJ, Chase MW, Rasmussen FN, eds. *Genera Orchidacearum volume 3: Orchidoideae (part two), Vanilloideae*. Oxford: Oxford University Press, 82–85.
- Palmer DD. 2003.** *Hawai'i's ferns and fern allies*. Honolulu: University of Hawai'i Press.
- Pascal M, Richer de Forges B, Le Guyader H, Simberloff D. 2008.** Mining and other threats to the New Caledonia biodiversity hotspots. *Conservation Biology* **22**: 498–499.
- Pauly A, Munzinger J. 2003.** Contribution à la connaissance des Hyménoptères Apoidea de Nouvelle-Calédonie et de leurs relations avec la flore butinée. *Annales de la Société Entomologique de France* **39**: 153–166.
- Pelletier B. 2006.** Geology of the New Caledonia region and its implications for the study of the New Caledonian biodiversity. In: Payri CE, Richer de Forges B, eds. *Compendium of marine species from New Caledonia*. Nouméa: IRD, 17–30.
- Persson C. 2001.** Phylogenetic relationships in Polygalaceae based on plastid DNA sequences from the *trnL-F* region. *Taxon* **50**: 763–779.

Picard M. 1999. *L'archipel néo-calédonien, 300 millions d'années pour assembler les pièces d'un puzzle géologique*. Nouméa: Centre de Documentation Pédagogique de Nouvelle-Calédonie.

Pillon Y, Munzinger J, Amir H, Hopkins HCF, Chase MW. 2009a. Reticulate evolution on a mosaic of soils: diversification of the New Caledonian endemic genus *Codia* (Cunoniaceae). *Molecular Ecology* **18**: 2263–2275.

Pillon Y, Hopkins HCF, Munzinger J, Amir H, Chase MW. 2009b. Cryptic species, gene recombination and hybridization in the genus *Spiraeanthemum* (Cunoniaceae) from New Caledonia. *Botanical Journal of the Linnean Society* **161**: 137–152.

Pillon Y, Munzinger J, Amir H, Lebrun M. 2010. Ultramafic soils and species sorting in the flora of New Caledonia. *Journal of Ecology* **98**: 1108–1116.

Pillon Y. 2012. Time and tempo of diversification in the flora of New Caledonia. *Botanical Journal of the Linnean Society* **170**: 288–298.

Pillon Y, Hopkins HCF, Rigault F, Jaffré T, Stacy EA. 2014. Cryptic adaptive radiation in tropical forest trees in New Caledonia. *The New phytologist* **202**: 521–30.

Pillon Y, Lucas E, Johansen JB, Sakishima T, Hall B, Geib SM, Stacy EA. 2015. An expanded *Metrosideros* (Myrtaceae) to include *Carpolepis* and *Tepualia* based on nuclear genes. *Systematic Botany* **40**: 782–790.

Pintaud JC, Baker WJ. 2008. A revision of the palm genera (Arecaceae) of New Caledonia. *Kew Bulletin* **63**: 61–73.

Pintaud JC, Jaffré T, Puig H. 2001. Chorology of New Caledonian palms and possible evidence of Pleistocene rain forest refugia. *Compte Rendus de l'Académie des Sciences de Paris, Sciences de la vie* **324**: 453–463.

Pocknall DT, Crosbie YM. 1988. Pollen morphology of *Beauprea* (Proteaceae): modern and fossil. *Review of Palaeobotany and Palynology* **53**: 305–327.

Pole M. 2010. Was New Zealand a primary source for the New Caledonian flora? *Alcheringa* **34**: 61–74.

Poncet V, Munoz F, Munzinger J, Pillon Y, Gomez C, Couderc M, Tranchant-Dubreuil C, Hamon S, de Kochko A. 2013. Phylogeography and niche modelling of the relict plant *Amborella trichopoda* (Amborellaceae) reveal multiple Pleistocene refugia in New Caledonia. *Molecular ecology* **22**: 6163–78.

Potgieter K, Albert VA. 2001. Phylogenetic relationships within Apocynaceae *s.l.* based on *trnL* intron and *trnL-F* spacer sequences and propagule characters. *Annals of the Missouri Botanical Garden* **88**: 523–549.

Pouteau R, Trueba S, Feild TS, Isnard I. 2015. New Caledonia: a Pleistocene refugium for rain forest lineages of relict angiosperms. *Journal of Biogeography* **42**: 2062–2077.

Pryer KM, Schuettpelz E, Wolf PG, Schneider H, Smith AR, Cranfill R. 2004. Phylogeny and evolution of ferns (monilophytes) with a focus on the early leptosporangiate divergences. *American Journal of Botany* **91**: 1582–1598.

Puente-Lelièvre C, Hislop M, Harrington M, Brown EA, Kuzmina M, Crayn DM. 2015. A five-marker molecular phylogeny of the Styphelieae (Epacridoideae, Ericaceae) supports a broad concept of Styphelia. *Australian Systematic Botany* **28**: 368–387.

Puttock CF, Quinn CJ. 1999. Generic concepts in Australian Gardenieae (Rubiaceae): a cladistic approach. *Australian Systematic Botany* **12**: 181.

Pye MG, Gadek PA, Edwards KJ. 2003. Divergence, diversity and species of the Australasian *Callitris* (Cupressaceae) and allied genera : evidence from ITS sequence data. *Australian Systematic Botany* **16**: 505–514.

Qiu YL, Lee J, Bernasconi-Quadroni F, Soltis DE, Soltis PS, Zanis M, Zimmer EA, Chen Z, Savolainen V, Chase MW. 1999. The earliest angiosperms: evidence from mitochondrial, plastid and nuclear genomes. *Nature* **402**: 404–407.

Quinn CJ, Brown EA, Heslewood MM, Crayn DM. 2005. Generic concepts in Styphelieae (Ericaceae): the *Cyathodes* group. *Australian Systematic Botany* **18**: 439–454.

Razafimandimbison SR, Bremer B. 2011. Nomenclatural changes and taxonomic notes in the tribe Morindeae (Rubiaceae). *Adansonia* **33**: 283–309.

Renner SS, Strijk JS, Strasberg D, Thébaud C. 2010. Biogeography of the Monimiaceae (Laurales): a role for East Gondwana and long-distance dispersal, but not West Gondwana. *Journal of Biogeography* **37**: 1227–1238.

Renner SS, Foreman DB, Murray D. 2000. Timing transantarctic disjunction in the Atherospermataceae (Laurales): evidence from coding and non coding chloroplast sequences. *Systematic Biology* **49**: 579–591.

Ruhfel BR, Bittrich V, Bove CP, Gustafsson MHG, Philbrick CT, Rutishauser R, Xi Z, Davis CC. 2011. Phylogeny of the clusioid clade (Malpighiales): evidence from the plastid and mitochondrial genomes. *American Journal of Botany* **98**: 306–325.

Sauquet H, Weston PH, Anderson CJ, Barker NP, Cantrill DJ, Mast AR, Savolainen V. 2009. Contrasted patterns of hyperdiversification in Mediterranean hotspots. *Proceedings of the National Academy of Science of the USA* **106**: 221–225.

Schluter D. 2000. *The ecology of adaptive radiation*. Oxford: Oxford University Press.

Schmid M. 2012. Contribution to the knowledge of Primulaceae (ex Myrsinaceae) of New Caledonia. III. *Tapeinosperma* Hook.f. and *Mangenotiella* gen. nov. genera. *Adansonia* **34**: 279–341.

Simmons MP, Cappa JJ, Archer RH, Ford AJ, Eichstedt D, Clevinger CC. 2008. Phylogeny of the Celastreae (Celastraceae) and the relationships of *Catha edulis* (qat) inferred from morphological characters and nuclear and plastid genes. *Molecular Phylogenetics and Evolution* **48**: 745–757.

- Simmons M, Bacon CD, Cappa JJ, McKenna MJ. 2012a.** Phylogeny of Celastraceae subfamilies Cassinoideae and Tripterygioideae inferred from morphological characters and nuclear and plastid loci. *Systematic Botany* **37**: 456–467.
- Simmons MP, McKenna MJ, Bacon CD, Yakobson K, Cappa JJ, Archer RH, Ford AJ. 2012b.** Phylogeny of Celastraceae tribe Euonymeae inferred from morphological characters and nuclear and plastid genes. *Molecular Phylogenetics and Evolution* **62**: 9–20.
- Sinclair WT, Mill RR, Gardner MF, Woltz P, Jaffré T, Preston ML, Hollingsworth ML, Ponge A, Möller M. 2002.** Evolutionary relationships of the New Caledonian heterotrophic conifer *Parasitaxus ustus* (Podocarpaceae), inferred from chloroplast *trnL-F* intron/spacer and nuclear ITS2 sequences. *Plant Systematics and Evolution* **233**: 79–104.
- Smith AC. 1973.** Studies of Pacific Island plants, XXV. the Myrsinaceae of the Fijian region. *Journal of the Arnold Arboretum* **54**: 228–292.
- Smith AC. 1996.** *Flora Vitiensis nova: a new flora of Fiji (spermatophytes only)*. Lawai: National Tropical Botanical Garden.
- Smith AC, Haas JE. 1975.** Studies of Pacific Island plants. XXIX. *Bleasdalea* and related genera of Proteaceae. *American Journal of Botany* **62**: 133.
- Sniderman JMK, Jordan GJ, Cowling RM. 2013.** Fossil evidence for a hyperdiverse sclerophyll flora under a non-Mediterranean-type climate. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 3423–8.
- Snow N, McFadden J, Evans TM, Salywon AM, Wojciechowski MF, Wilson PG. 2011.** Morphological and molecular evidence of polyphyly in *Rhodomyrtus* (Myrtaceae: Myrteae). *Systematic Botany* **36**: 390–404.
- Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF, Refulio-Rodriguez NF, Walker JB, Moore MJ, Carlswald BS, Bell CD, Latvis M, Crawley S, Black C, Diouf D, Xi Z, Rushworth CA, Gitzendanner MA, Sytsma KJ, Qiu YL, Hilu KW, Davis CC, Sanderson MJ, Beaman RS, Olmstead RG, Judd WS, Donoghue MJ, Soltis PS. 2011.** Angiosperm phylogeny: 17 genes, 640 taxa. *American Journal of Botany* **98**: 704–730.
- Strijk JS, Bone RE, Thébaud C, Buerki S, Fritsch PW, Hodkinson TR, Strasberg D. 2014.** Timing and tempo of evolutionary diversification in a biodiversity hotspot: Primulaceae on Indian Ocean islands. *Journal of Biogeography* **41**: 810–822.
- Stuessy TF, König C, Sepulveda PL. 2014.** Paraphyly and endemic genera of oceanic islands: implications for conservation. *Annals of the Missouri Botanical Garden* **100**: 50–78.
- Stull GW, Duno de Stefano R, Soltis DE, Soltis PS. 2015.** Resolving basal lamiid phylogeny and the circumscription of Icacinaceae with a plastome-scale data set. *American Journal of Botany* **102**: 1794–1813.
- Su HJ, Hu JM, Anderson FE, Der JP, Nickrent DL. 2015.** Phylogenetic relationships of Santalales with insights into the origins of holoparasitic Balanophoraceae. *Taxon* **64**: 491–506.

Sweeney PW, Bradford JC, Lowry PP. 2004. Phylogenetic position of the New Caledonian endemic genus *Hooglandia* (Cunoniaceae) as determined by maximum parsimony analysis of chloroplast DNA. *Annals of the Missouri Botanical Garden* **91**: 266–274.

Swenson U, Lowry PP, Munzinger J, Rydin C, Bartish IV. 2008. Phylogenetic and generic limits in the *Niemeyera* complex of New Caledonian Sapotaceae: evidence of multiple origins of the anisomerous flower. *Molecular Phylogenetics and Evolution* **49**: 909–929.

Swenson U, Munzinger J, Lowry PP, Cronholm B, Nylinder S. 2015. Island life–classification, speciation and cryptic species of *Pycnandra* (Sapotaceae) in New Caledonia. *Botanical Journal of the Linnean Society* **179**: 57–77.

Swenson U, Munzinger J. 2009. Revision of *Pycnandra* subgenus *Pycnandra* (Sapotaceae), a genus endemic to New Caledonia. *Australian Systematic Botany* **22**: 437–465.

Swenson U, Munzinger J. 2016. Five new species and a systematic synopsis of *Pycnandra* (Sapotaceae), the largest endemic genus in New Caledonia. *Australian Systematic Botany* **29**: in press.

Swenson U, Nylinder S, Munzinger J. 2013. Towards a natural classification of Sapotaceae subfamily Chrysophylloideae in Oceania and Southeast Asia based on nuclear sequence data. *Taxon* **62**: 746–770.

Tank DC & Donoghue MJ. 2010. Phylogeny and phylogenetic nomenclature of the Campanulidae based on an expanded sample of genes and taxa. *Systematic Botany* **35**: 425–441.

Thorne RF. 1969. Floristic relationships between New Caledonia and the Solomon Islands. *Philosophical Transactions of the Royal Society B-Biological Sciences* **255**: 595–602.

Thornhill AH, Ho SYW, Külheim C, Crisp MD. 2015. Interpreting the modern distribution of Myrtaceae using a dated molecular phylogeny. *Molecular Phylogenetics and Evolution* **93**: 29–43.

Tirel C, Jérémie J. 1996. Alseuosmiaceae. In: Morat P, ed. *Flore de la Nouvelle-Calédonie et Dépendances*. Paris: Muséum National d’Histoire Naturelle, 100–106.

Tokuoka T. 2007. Molecular phylogenetic analysis of Euphorbiaceae *sensu stricto* based on plastid and nuclear DNA sequences and ovule and seed character evolution. *Journal of Plant Research* **120**: 511–522.

Tokuoka T, Tobe H. 2006. Phylogenetic analyses of Malpighiales using plastid and nuclear DNA sequences, with particular reference to the embryology of Euphorbiaceae *sens. str.* *Journal of Plant Research* **119**: 599–616.

Trewick SA, Paterson AM, Campbell HJ. 2007. Hello New Zealand. *Journal of Biogeography* **34**: 1–17.

Verboom GA, Archibald JK, Bakker FT, Bellstedt DU, Conrad F, Dreyer LL, Forest F, Galley C, Goldblatt P, Henning JF, Mummenhoff K, Linder HP, Muasya M, Oberlander KC, Savolainen V, Snijman DA, van der Niet T, Nowell TL, 2009. Origin and

diversification of the Greater Cape flora: ancient species repository, hot-bed of recent radiation, or both? *Molecular Phylogenetics and Evolution* **51**: 44–53.

Vinnersten A, Bremer K. 2001. Age and biogeography of major clades in Lilliales. *American Journal of Botany* **88**: 1695–1703.

Viot R. 1956. *La végétation canaque*. Paris: Faculté des Sciences de l'université de Paris.

Viot R. 1967. Protéacées. In: Aubréville A, ed. *Flore de la Nouvelle-Calédonie et Dépendances*. Paris: Muséum National d'Histoire Naturelle, 1–254.

Wagner WL, Herbst DR, Sohmer SH. 1999. *Manual of the flowering plants of Hawai'i*. Honolulu: Bernice Pauahi Bishop Museum Special Publication.

Webster GL. 1994. Synopsis of the genera and suprageneric taxa of Euphorbiaceae. *Annals of the Missouri Botanical Garden* **81**: 33–144.

Weigend M, Gottschling M, Hoot S, Ackermann M. 2004. A preliminary phylogeny of Loasaceae subfam. Loasoideae (Angiospermae: Cornales) based on *trnL* sequence data, with consequences for systematics and historical biogeography. *Organisms Diversity & Evolution* **4**: 73–90.

Weston PH, Barker NP. 2006. A new suprageneric classification of the Proteaceae, with an annotated checklist of genera. *Telopea* **11**: 314–344.

Whistler A. 2011. *The rare plants of Samoa*. Apia: Conservation International.

Whitlock BA, Hale AM, Indorf JL, Wilkins CF 2011. Polyphyly of *Rulingia* and *Commersonia* (Lasiopetalae, Malvaceae s.l.). *Australian Systematic Botany* **24**: 215–225.

Wilkie P, Clark A, Pennington RT, Cheek M, Bayer C, Wilcock CC. 2006. Phylogenetic relationships within the subfamily Sterculioideae (Malvaceae/Sterculiaceae-Sterculieae) using the chloroplast gene *ndhF*. *Systematic Botany* **31**: 160–170.

Wilson PG. 2011. Myrtaceae. In: Kubitzki K, ed. *The families and genera of vascular plants. Volume X. Sapindales, Cucurbitales, Myrtaceae*. Heidelberg: Springer-Verlag, 212–271.

Woo VL, Funke MM, Smith JF, Lockhart PJ, Garnock-Jones PJ. 2011. New World origins of southwest Pacific Gesneriaceae: multiple movements across and within the South Pacific. *International Journal of Plant Science* **172**: 434–457.

Wright SD, Keeling DJ, Ashton FG, Dawson JW, Gardner RC. 2000. Phylogenetic analyses of New Caledonian *Metrosideros* and *Carpolepis* (Myrtaceae) from nrDNA (ITS) sequences. *Australian Systematic Botany* **13**: 919–926.

Wurdack KJ, Hoffmann P, Chase MW. 2005. Molecular phylogenetic analysis of uniovulate Euphorbiaceae (Euphorbiaceae *sensu stricto*) using plastid *rbcL* and *trnL-F* DNA sequences. *American Journal of Botany* **92**: 1397–1420.

Zerega NJC, Clement WL, Datwyler SL, Weiblen GD. 2005. Biogeography and divergence times in the mulberry family (Moraceae). *Molecular Phylogenetics and Evolution* **37**: 402–416.

Zielske S, Haase M. 2014a. When snails inform about geology: Pliocene emergence of islands of Vanuatu indicated by a radiation of truncatelloidean freshwater gastropods (Caenogastropoda: Tateidae). *Journal of Zoological Systematics and Evolutionary Research* **52**: 217–236.

Zielske S, Haase M. 2014b. New insights into tateid gastropods and their radiation on Fiji based on anatomical and molecular methods (Caenogastropoda: Truncatelloidea): Tateid Gastropods on Fiji. *Zoological Journal of the Linnean Society* **172**: 71–102.

Zona S, Baker WJ. 2014. *Solfia* transferred to *Balaka*. *Palms* **58**: 191–192.

Table 1. Phylogenetic status of the vascular plant endemic genera endemic to New Caledonia. Column 1. Distinctiveness has been confirmed by phylogenetic studies with thorough sampling (all closely related genera have been included). Available data suggest that the genus is monophyletic (or in several cases monotypic or species-poor) and it seems unlikely that it is nested in another one. Column 2. All current evidence support the distinctiveness of the genus, although sampling of related genera may not be exhaustive or resolution in phylogenetic relationships may be low, or the phylogenetic study is not published yet. Column 3. There is some evidence that the genus may not be monophyletic or may be nested in another one or that the genus is closely related to a large genus that has not been thoroughly sampled yet, also present in New Caledonia or not, for which delimitation may be problematic, and in which the candidate genus could be nested. Column 4. The genus has been found to be nested in another genus with strong support or has been synonymized since the publication of Morat *et al.* (2012). Column 5. The genus has never been included in any molecular phylogenetic study and there are no DNA sequences available in GenBank. Underlined genera: restricted to ultramafic substrates (or almost so). Highlighted genera: never found on ultramafic substrates (or almost so). Numbers in parenthesis are species numbers according to Munzinger *et al.* (2016).

Status confirmed		Likely	Need further work	To be synonymized	Never investigated
<u>Achlydosa</u> (1) <u>Amborella</u> (1) <u>Amphorogyne</u> (3) <u>Apiopetalum</u> (2) <u>Arillastrum</u> (1) <u>Austrotaxus</u> (1) <u>Beauprea</u> (13) <u>Beaupreopsis</u> (1) <u>Campynemanthe</u> (3) <u>Canacomyrca</u> (1) <u>Clematepistephium</u> (1) <u>Clinosperma</u> (4) <u>Cloezia</u> (6) <u>Codia</u> (15) <u>Coilochilus</u> (1) <u>Cyphokentia</u> (2) <u>Daenikera</u> (1) <u>Depanthus</u> (2) <u>Dicarpellum</u> (4) <u>Eriaxis</u> (1) <u>Eucarpha</u> (2) <u>Gongrodiscus</u> (3) <u>Hachettea</u> (1) <u>Hooglandia</u> (1)	<u>Jaffrea</u> (2) <u>Loxodiscus</u> (1) <u>Menepetalum</u> (4) <u>Montrouziera</u> (6) <u>Myodocarpus</u> (10) <u>Myrtastrum</u> (1) <u>Nemuaron</u> (1) <u>Neoguillauminia</u> (1) <u>Neoschmidia</u> (2) <u>Oncotheca</u> (2) <u>Pachyplectron</u> (3) <u>Pancheria</u> (27) <u>Paracryphia</u> (1) <u>Parasitaxus</u> (1) <u>Phelline</u> (14) <u>Platyspermatum</u> (1) <u>Podonephelium</u> (9) <u>Pycnandra</u> (59) <u>Salaciopsis</u> (6) <u>Sparattosyce</u> (2) <u>Storthocalyx</u> (5) <u>Strasburgeria</u> (1) <u>Stromatopteris</u> (1) <u>Viotia</u> (6)	<u>Acropogon</u> (25) <u>Arthroclianthus</u> (19) <u>Balgoya</u> (1) <u>Cerberiopsis</u> (3) <u>Crossosperma</u> (2) <u>Cyathopsis</u> (3) <u>Gonatosstylis</u> (2) <u>Greslania</u> (3) <u>Maxwellia</u> (1) <u>Normandia</u> (1) <u>Nephrodesmus</u> (5) <u>Pripterygia</u> (1) <u>Periomphale</u> (1) <u>Scagea</u> (2)	<u>Actinokentia</u> (2) <u>Adenodaphne</u> (4) <u>Artia</u> (4) <u>Basselinia</u> (14) <u>Bocquillonina</u> (14) <u>Burretiokentia</u> (5) <u>Chambeyronia</u> (2) <u>Cocconerion</u> (2) <u>Cyphophoenix</u> (4) <u>Deltaria</u> (1) <u>Gea</u> (6) <u>Kanakomyrtus</u> (6) <u>Kentiopsis</u> (4) <u>Kermadecia</u> (4) <u>Kibaropsis</u> (1) <u>Lasiochlamys</u> (11) <u>Morierina</u> (2) <u>Myrtopsis</u> (8) <u>Sleumerodendron</u> (1) <u>Solmsia</u> (2) <u>Thiollierea</u> (12)	<u>Comptonella</u> (8) <u>Dutaillyea</u> (2) <u>Garnieria</u> (1) <u>Oxanthera</u> (5) <u>Picrella</u> (3)	<u>Dutailliopsis</u> (1) <u>Elaphanthera</u> (1) <u>Gastrolepis</u> (2) <u>Longetia</u> (1) <u>Mangenotiella</u> (1) <u>Myricanthe</u> (1) <u>Pleurocalyptus</u> (2) <u>Purpureostemon</u> (1)
				Synonymized since Morat <i>et al.</i> (2012)	
				<u>Boronella</u> (4) <u>Carpolepis</u> (3) <u>Neocallitropsis</u> (1) <u>Pritchardiopsis</u> (1)	

Table 2. Statistics for different archipelagos in the Pacific Ocean.

Archipelago	New Caledonia	New Zealand	Solomon Islands	Vanuatu	Fiji	Samoa	French Polynesia and Cook Islands	Hawaiian Islands
Surface (km ²)	18,576	268,021	28,400	12,190	18,274	3,030	3,916	16,636
Age of biota (Myr)	37 ^a	25 ^b	?	5 ^c	7-16 ^d	13,5 ^e	≈18 ^f	mostly ≤5 ^g
Highest elevation (m)	1,628	3,754	2,332	1,879	1,325	1,858	2,241	4,207
Number of vascular plant species	3,389 ^h	2,221 ⁱ	?	?	1,315 ^j	550 ^k	651 ^m	1,207 ⁿ
Specific endemism (%)	75.1 ⁸	78 ⁱ	?	?	62 ^j	31 ^k	70.9 ^m	88 ⁿ
Number of endemic genera	61-91	28-44	?2	1	1-5	0	≈ 9	22-35

^aGrandcolas *et al.* (2008), ^bTrewick *et al.* (2007), ^cZielske & Haase (2014a), ^dZielske & Haase (2014b), ^eEwart (1988), ^fMaury *et al.* (2014), ^gPrice & Clague, ^hMunzinger *et al.*, ⁱBreitwieser *et al.* (2012), ^jHeads, ^kangiosperms only, Whistler (2011), ^mFrench Polynesia only, Florence in Morat *et al.* (2012), ⁿGustafson *et al.* (2014).

Figure 1. Some representatives of New Caledonia endemic genera: A. *Loxodiscus coriaceus* Hook.f. (Sapindaceae), B. *Pancheria robusta* Guillaumin (Cunoniaceae), C. *Platyspermatium crassifolium* Guillaumin (Alseuosmiaceae), D. *Thiollierea kaalaensis* (N.Hallé & Jérémie) Barrabé & Mouly, E. *Hachettea austrocaledonica* Baill. (Balanophoraceae), F. *Campynemanthe viridiflora* Baill. (Campynemataceae). Photograph credits: V. Hequet (A), Y. Pillon (B, C, D, F), L. Barrabé (E).

