



HAL
open science

Fate of the insecticide chlordecone under methanogenic conditions

Hervé Macarie, Yoan Labrousse, Aurélie Amic, Nicolas Cimetiere, Alain Soler, Sébastien Bristeau, Christophe Mouvet

► **To cite this version:**

Hervé Macarie, Yoan Labrousse, Aurélie Amic, Nicolas Cimetiere, Alain Soler, et al.. Fate of the insecticide chlordecone under methanogenic conditions. XIV Latin American Workshop and Symposium on Anaerobic Digestion (DAAL XIV), Instituto de Ingeniería - Universidad Nacional Autónoma de México (II-UNAM), campus Juriquilla, Oct 2023, Querétaro, Mexico. ird-04433493

HAL Id: ird-04433493

<https://ird.hal.science/ird-04433493v1>

Submitted on 16 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**XIV Latin American Workshop and
Symposium on Anaerobic Digestion
Querétaro, Mexico, October 23-27, 2023**

CONFERENCE PROCEEDINGS

Oral Presentations

The DAAL XIV is organized by the Engineering Institute of
the National Autonomous University of Mexico

Scientific Committee

Germán Buitrón (Chairman)
Universidad Nacional Autónoma de México,
Mexico

Carlos A. Chernicharo,
Universidade Federal de Minas Gerais, Brazil

Sergio Francisco Aquino
Universidade Federal de Minas Gerais, Brazil

Lourdinha Florencio
Universidade Federal de Pernambuco, Brazil

André Bezerra dos Santos
Universidade Federal do Ceará, Brazil

Marcelo Zaiat
Universidade de Sao Paulo, Brazil

Gustavo Ribeiro
Universidad Estatal de Sao Paulo, Brazil

Alexandra Cerón Vivas
Universidad Pontificia Bolivariana, Colombia

Juan S Arcila Henao
Universidad de Manizales, Colombia

Rolando Chamy
Pontificia Universidad Católica de Valparaíso, Chile

David Jeison
Pontificia Universidad Católica de Valparaíso, Chile

Gladys Vidal
Universidad de Concepción, Chile

Lorna Guerreo
Universidad Técnica Federico Santa María, Chile

Jorge Rodríguez
Khalifa University of Science and Technology,
United Arab Emirates

José Luis Sanz
Universidad Autónoma de Madrid, Spain

Juan Lema
Universidad de Santiago de Compostela, Spain

Sara Pérez Elvira
Universidad de Valladolid, Spain

Jim Field
Arizona University, United States

Jorge Arreola Vargas
Texas A&M University, United States

Jean Philippe Steyer
INRAe, France

Nicolas Bernet
INRAe, France

Thierry Ribeiro
Institut Polytechnique LaSalle Beauvais, France

Hervé Macarie
Institut de Recherche pour le Développement,
France

Jules van Lier
Delft University of Technology, Holland

Fons Stams
Wageningen University, Holland

Adalberto Noyola
Universidad Nacional Autónoma de México,
Mexico

Francisco Cervantes
Universidad Nacional Autónoma de México,
Mexico

Iván Moreno
Universidad Nacional Autónoma de México,
Mexico

Contents

Oral presentation program	1
Session A1. Microbial ecology in anaerobic digestion.....	2
Session B1. Modelling and control	17
Session A2. Wastewater, sludge and organic wastes treatment	33
Session B2. Industrial wastewater	51
Session A3. Emerging technologies for sludge digestion and organic wastes	68
Session B3. Beyond biogas: hydrogen, bioelectrochemical systems, recovery of value-added chemicals	86
Session A4. Removal of emerging contaminants.....	104
Session B4. Beyond biogas: hydrogen, bioelectrochemical systems, recovery of value-added chemicals	121
Session A5. Post-treatment and conditioning of anaerobic effluent, biogas and digestate	139
Session B5. Circular economy, life cycle analysis and full scale experiences	157

**Session A5. Post-treatment and
conditioning of anaerobic effluent, biogas
and digestate**

Room A. 11:00 – 13:00 hours

Thursday 26

Session A5. Post-treatment and conditioning of anaerobic effluent, biogas and digestate

Chair: David Jeison

Time	Abstract ID	Title
11:00 - 11:15	8029	Effects of thermal hydrolysis process on the colloidal fraction of digestate and viscosity of reject water. <u>Javier Pavez</u> , Leslie Paiz, Jules van Lier, Merle de Kreuk. Netherlands
11:15 – 11:30	8199	Stating the quality of an acidogenic digestate. Betizabeth Arellano, Rosalinda Campuzano, Carmen Fajardo, <u>Oscar Monroy</u> . Mexico
11:30 – 11:45	8216	Start-up and Operation of an Anaerobic Filter for a Simultaneous Autotrophic-Heterotrophic Denitrification System. Andrea Barahona, <u>Lorna Guerrero</u> , Cesar Huiliñir, Rolando Chamy, Miguel Palacios. Chile
11:45 – 12:00	8221	Filtration of an anaerobic effluent by a self-forming dynamic membrane submerged in anaerobic granular sludge at pilot scale. Andrea Rodríguez-Medina, Eliel Aguirre-Cañete, Adalberto Noyola. Mexico

Program Oral presentations

12:00 – 12:15	8231	Feasibility of the utilization of liquid digestate from the AD of sewage sludge as a nutrient supply for the biomethanation of biogas with exogenous H₂ in a biotrickling filter.
		José Manuel García, Federico Ferrari, María del Mar Micó, María Fdz-Polanco, <u>Israel Díaz</u> .
		Spain
12:15 – 12:30	8191	Effect of the pH on the performance of anaerobic piggery wastewater treatment coupled with membrane-based ammonia extraction.
		<u>Fanny Rivera</u> , Pedro Prádanos, Antonio Hernández, Laura Palacio, Raúl Muñoz.
		Spain
12:30 – 12:45	8281	Microaeration with waste gas from desorption chamber: integration of processes in UASB reactor for recovery and treatment of methane and hydrogen sulfide.
		Tamile Marinho, <u>Cláudio Souza</u> .
		Brazil
12:45 – 13:00	8970	Fate of the insecticide chlordecone under methanogenic conditions.
		<u>Hervé Macarie</u> , Yoan Labrousse, Aurélie Amic, Nicolas Cimetiere, Alain Soler, Bristeau Sébastien, Mouvet Christophe.
		France

Fate of the insecticide chlordecone under methanogenic conditions

**Hervé Macarie^{1,2}, Yoan Labrousse^{2,3}, Aurélie Amic¹, Nicolas Cimetiere⁴, Alain Soler⁵,
Bristeau Sébastien⁶, Mouvet Christophe⁶**

¹ IRD, Aix Marseille Univ, Avignon Univ, CNRS, IMBE, Marseille, France. herve.macarie@ird.fr

² IRD, IMBE, Campus Agro-environnemental Caraïbe (CAEC), Martinique, France

³ IRD, Aix Marseille Univ, LPED, Marseille, France, ⁴ Ecole Nationale Supérieure de Chimie de Rennes, France

⁵ CIRAD, UPR GECO, Saint-Pierre, Réunion, France, ⁶ BRGM, Orléans, France

Highlights

- A natural methanogenic biomass can remove 1 or 2 chlorine atoms from chlordecone
- Chlordecone dechlorination by methanogenic biomass is less efficient than adsorption
- Digester sludge is not a good source of chlordecone degraders for soil bioremediation

Keywords: mono-hydrochlordecone; chlordecol; 5b-hydrochlordecone; organochlorine; anaerobiosis

Introduction

Chlordecone (CLD) is an organochlorine insecticide of raw formula $C_{10}Cl_{10}O$, used from 1972 to 1993 (date of its ban) to control the banana weevil populations in French West Indies (FWI). By the time of writing, i.e. three decades after its ban use, CLD is still present in the FWI soils and has led to widespread contamination of the other FWI environmental compartments and associated food resources, resulting in a widespread impregnation of the West Indian population, with proven health consequences. Despite a series of measures to prevent CLD from reaching the consumer's plate, it continues to circulate actively at least among the population living in the most contaminated areas. A definitive solution to the problem would be to destroy the stock of CLD still present in soil. Of all the options available to achieve this goal, microbial degradation is generally an economically attractive option, as well as being potentially applicable *in situ*. Until the late 2010's, because of its structure, CLD was considered non-biodegradable but the work of Dolfing et al (2012) showed that there was no thermodynamic impediment to its partial transformation (e.g dechlorination) or ultimate mineralization, and that microorganisms capable of catalyzing these reactions should/could exist. Old literature data (Jablonski et al., 1996) suggested that all methanogens would be enzymatically equipped to attack CLD and that natural sources of these organisms such as anaerobic digesters could eventually be used for bioaugmentation in a soil remediation process. This possibility was reinforced by the monitoring of CLD concentrations at the inlet and outlet of 2 FWI industrial anaerobic digesters treating contaminated rum vinasses, which showed that 53 to 91% of the CLD present in the vinasses was not found at the outlet of the digesters and had apparently not accumulated in their biomass (Macarie et al., 2013). These conclusions remained fragile, however, due to i) the difficulty of quantifying biomass in reactors of several hundred m^3 , ii) the low concentration of CLD in vinasses (a few $\mu g/L$), and iii) their complex composition, which make it impossible to track any transformation products, including chlorides. The objective of this work was to reproduce at lab scale the operation of an anaerobic digester fed with a simple culture medium containing CLD in order to make a precise CLD mass balance and confirm or not the capacity of any methanogenic mixed culture to degrade it.

Material and Methods

The experiments were done at ambient temperature with a 2 L glass UASB reactor covered by an aluminum film which was inoculated with a known quantity of granular methanogenic sludge from one of the two industrial reactors previously studied. The reactor was fed continuously for 81 days at a constant hydraulic retention time and volumetric loading rate through PTFE tubing with a simple low-chloride aqueous culture medium containing 2 mg CLD/L in dissolved form and the nutrients required for the methanization process (sources of C, N, P, S, trace metals). CLD was not added to the culture medium until one week after the start of the operation. Two categories of parameters were followed, those necessary to control the operation of the system and confirm the good functioning of the methanogenic population (COD inlet/outlet, pH, sludge redox potential, methane production, feed rate) and those necessary to evaluate CLD degradation (concentration of CLD and some of its known

degradation intermediates at the reactor inlet, outlet and in the reactor sludge and chloride concentration at the outlet). The quantification of CLD and known intermediates was done by GC-MS but also LC-MS for the reactor sludge. GC-MS was additionally used to search for the eventual presence of unknown transformation products. Chloride was determined by ionic chromatography

Results and Discussion

The parameters monitored confirmed that the system performed well under methanogenic conditions: very negative redox potential ($E_{Ag/AgCl} = -345$ to -434.2 mV) characteristic of this type of environment, total COD removal of $96.2 \pm 2.5\%$ coupled with a methane yield very close to the theoretical yield of 0.35 LCH₄/g COD_{degraded}. Remarkably, reactor performance remained stable after the addition of 2 mg CLD/L to the culture medium indicating the absence of CLD toxicity at this concentration for all the microbial partners required for the conversion of carbonaceous substrates to CH₄. In addition to CLD (2.21 ± 0.39 mg/l measured), chlordecol (17.2 ± 3.9 µg/l) and 5b-hydro-CLD (1.47 ± 0.1 µg/l) were detected in the culture medium. They correspond to impurities present in the analytical standard-grade CLD used to prepare it. A removal of over 99% of CLD and 5b-hydro-CLD and over 95% of chlordecol between the reactor inlet and outlet was obtained from the start of the CLD feed. This reduction was not associated with the accumulation of degradation intermediates in the liquid phase, with the exception of a monohydro-CLD (CLD-1Cl) different from 5b-hydro-CLD whose concentration however did not exceed 3.15 µg/l, representing barely 0.17 mol% of the CLD supplied to the reactor. Traces of 2 dihydro-CLD (CLD-2Cl) were also detected at the reactor outlet, but due to a lack of available analytical standards, they could not be quantified. Chloride analysis in the aqueous phase at the reactor outlet was found to be within the background noise of that of the low chloride content deionized water used (0.23 ± 0.13 mg/L) to prepare the medium, and well below that (1.43 mg Cl-/L) corresponding to 100% dechlorination of 2 mg CLD/L, thus demonstrating the absence of CLD transformation into more dechlorinated products not detected by GC-MS or its full mineralization. In order to reach a definitive conclusion on the fate of CLD in the reactor, its biomass was recovered and its CLD content quantified. Analysis on this matrix proved difficult, but finally, after much trial and error, the last method used (SPME¹-GC-MS combined to standard addition on fresh biomass), showed that at least 69% of the CLD brought to the reactor had accumulated in its biomass, and that the CLD-1Cl also detected represented 4.3 mol% of the fed CLD.

Although the CLD mass balance did not close (25% of the CLD fed to the system was not recovered at the exit of the reactor, in its sludge or in the form of the transformation products quantified), the previous results show that the main mechanism of CLD removal in the reactor was due to adsorption on the biomass. The presence of CLD-1Cl and CLD-2Cl not detected at the entrance of the system confirms however that the methanogenic biomass used had the capacity to dechlorinate CLD, but that this capacity was very low despite massive inoculation of the reactor. They therefore suggest that the microbial group capable of dechlorinating CLD was probably not methanogenic and represented a very tiny proportion of the microbial consortium. Although the results suggest a slow capacity for enrichment of this population over time (CLD-1Cl concentration increased at the outlet from 0.031 µg/L on day 11 to 3.157 µg/L on day 81), they also show that the possibility of using digester sludge for bioaugmentation in a soil remediation process to achieve CLD degradation would not be viable.

Acknowledgements.

This work was financially supported by INRA DEMICHLORD programme and FEDER Martinique 2007-2013.

References

- Dolfing, J., Novak, I., Archelas, A., Macarie, H. (2012). Gibbs free energy of formation of chlordecone and potential degradation products: implications for remediation strategies and environmental fate. *Environ Sci Technol*, 46, 8131-8139.
- Jablonski, P.E., Pheasant, D.J., Ferry, J.G. (1996). Conversion of kepone by *Methanosarcina thermophila*. *FEMS Microbiol. Lett.* **139**, 169-173.
- Macarie, H., Labrousse, Y., Sastre-Conde, I., Bristeau, S., Mouvet, C. (2013). Further evidence of biodegradability of the POP chlordecone under methanogenic conditions. In: CD ROM Proceedings 13th World Congress on Anaerobic Digestion: recovering (bio)resources for the world, J.M. Lema, F. Fdez-Polanco, M. Carballa, J. Rodriguez, S. Suárez (Eds.), 25-28 Jun 2013, Santiago de Compostela, Spain, ISBN 978-84-695-7756-1, 2 pages.

¹ SPME: Solid Phase Micro Extraction