



**HAL**  
open science

## Modification of REE distribution of ordinary chondrites from Atacama (Chile) and Lut (Iran) hot deserts: Insights into the chemical weathering of meteorites

Hamed Pourkhorsandi, Massimo d’Orazio, Pierre Rochette, Millarca Valenzuela, Jérôme Gattacceca, Hassan Mirnejad, Brad Sutter, Aurore Hutzler, Maria Aboulahris

### ► To cite this version:

Hamed Pourkhorsandi, Massimo d’Orazio, Pierre Rochette, Millarca Valenzuela, Jérôme Gattacceca, et al.. Modification of REE distribution of ordinary chondrites from Atacama (Chile) and Lut (Iran) hot deserts: Insights into the chemical weathering of meteorites. *Meteoritics and Planetary Science*, 2017, 52 (9), pp.1843-1858. 10.1111/maps.12894 . ird-04812133

**HAL Id: ird-04812133**

**<https://ird.hal.science/ird-04812133v1>**

Submitted on 8 Dec 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## Modification of REE distribution of ordinary chondrites from Atacama (Chile) and Lut (Iran) hot deserts: Insights into the chemical weathering of meteorites

Hamed POURKHORSANDI<sup>1\*</sup> , Massimo D'ORAZIO<sup>2</sup>, Pierre ROCHETTE<sup>1</sup> ,  
Millarca VALENZUELA<sup>3</sup>, Jérôme GATTACCECA<sup>1</sup> , Hassan MIRNEJAD<sup>4,5</sup>, Brad SUTTER<sup>6</sup>,  
Aurore HUTZLER<sup>7</sup>, and Maria ABOULHRIS<sup>1,8</sup>

<sup>1</sup>CNRS, Aix-Marseille Univ., IRD, Coll. France, CEREGE, Aix-en-Provence, France

<sup>2</sup>Dipartimento di Scienze della Terra, Università di Pisa, Via S. Maria 53, I-56126 Pisa, Italy

<sup>3</sup>Instituto de Astrofísica, Pontificia Universidad Católica de Chile, Vicuña Mackena 4860, Macul, Santiago, Chile

<sup>4</sup>Department of Geology, Faculty of Sciences, University of Tehran, Tehran 14155-64155, Iran

<sup>5</sup>Department of Geology and Environmental Earth Sciences, Miami University, Oxford, Ohio 45056, USA

<sup>6</sup>Jacobs-NASA, Johnson Space Center, Houston, Texas 77058, USA

<sup>7</sup>Natural History Museum, Burgring 7, A-1010 Vienna, Austria

<sup>8</sup>Département de Géologie, Faculté des Sciences, Université Hassan II Casablanca, Casablanca, Morocco

\*Corresponding author. E-mail: pourkhorsandi@cerege.fr

(Received 04 June 2016; revision accepted 10 April 2017)

---

**Abstract**—The behavior of rare earth elements (REEs) during hot desert weathering of meteorites is investigated. Ordinary chondrites (OCs) from Atacama (Chile) and Lut (Iran) deserts show different variations in REE composition during this process. Inductively coupled plasma–mass spectrometry (ICP-MS) data reveal that hot desert OCs tend to show elevated light REE concentrations, relative to OC falls. Chondrites from Atacama are by far the most enriched in REEs and this enrichment is not necessarily related to their degree of weathering. Positive Ce anomaly of fresh chondrites from Atacama and the successive formation of a negative Ce anomaly with the addition of trivalent REEs are similar to the process reported from Antarctic eucrites. In addition to REEs, Sr and Ba also show different concentrations when comparing OCs from different hot deserts. The stability of Atacama surfaces and the associated old terrestrial ages of meteorites from this region give the samples the necessary time to interact with the terrestrial environment and to be chemically modified. Higher REE contents and LREE-enriched composition are evidence of contamination by terrestrial soil. Despite their low degrees of weathering, special care must be taken into account while working on the REE composition of Atacama meteorites for cosmochemistry applications. In contrast, chondrites from the Lut desert show lower degrees of REE modification, despite significant weathering signed by Sr content. This is explained by the relatively rapid weathering rate of the meteorites occurring in the Lut desert, which hampers the penetration of terrestrial material by forming voluminous Fe oxide/oxyhydroxides shortly after the meteorite fall.

---

### INTRODUCTION

As soon as a meteoroid enters the Earth's atmosphere, it is subjected to terrestrial alteration (e.g., Bland et al. 2006). The meteorites which are used for cosmochemical studies are being collected immediately or thousands to several hundred thousands of years after their fall. It has been documented that the mineralogy and chemical

compositions of these samples can be affected over a human lifetime (Socki et al. 1991; Bland et al. 1998; Pillinger et al. 2013). Alteration occurs on even shorter time scales of less than decades in the laboratory environment (Velbel 2014) or tens of days in a humid natural environment (Bischoff et al. 2011). Initial investigations on the meteorite weathering processes commenced just after the recognition of a high number of

weathered hot desert samples (e.g., Buddhue 1939; Olsen and Fuchs 1967). Since that time, the discovery of large numbers of meteorites from cold and hot deserts has drawn the attention of researchers to document the effects of weathering on meteorites to avoid any inaccurate interpretation on their geochemical compositions.

The main purposes of meteorite weathering studies are (1) understanding the weathering process occurring within a meteorite and its relationship to the finding place of the meteorite, and (2) identifying the chemical and mineralogical modification during this process, to account for eventual bias in cosmochemical studies (Bland et al. 2006). Several studies have focused on the effect of weathering on the chemical and mineralogical compositions of meteorites from Antarctic and rare hot desert achondrites (White et al. 1967; Gooding 1982; Koeberl and Cassidy 1989; Torigoye-Kita et al. 1995; Swindle et al. 1998; Llorca et al. 2013). Considering their importance and also higher abundance, these studies on the ordinary chondrites (OCs) have high importance, compared to achondrites.

Among the trace elements which are used in geochemical and cosmochemical studies of meteorites, rare earth elements (REEs) have a significant importance. Variations in the abundance of REEs are diagnostic proxies to study evaporation/condensation processes in the nebula (Boynton 1975), parent-body aqueous alteration (Inoue et al. 2009), thermal metamorphism (Murrell and Burnett 1983), magmatic differentiation (Boynton et al. 1976), and terrestrial weathering (Crozas et al. 2003). Rare earth elements mobilization during terrestrial weathering has been studied by Mittlefehldt and Lindstrom (1991) on Antarctic eucrites and by Crozas et al. (2003) on hot desert achondrites. Rare earth element sensitivity to weathering may also affect Nd isotopic composition (Crozas and Wadhwa 2001).

Despite the high number of OCs, limited studies have been conducted on their REE mobilization during terrestrial weathering. Al-Kathiri et al. (2005) suggested a LREE enrichment in OCs from Oman, without presenting HREE concentrations. Saunier et al. (2010) mentioned the high amount of LREE in two highly weathered H5 and L6 OCs.

The goals of this study were to observe the possible differences between the REE contents of meteorites from different hot deserts (in particular Atacama and Lut) and to study the process of REE mobilization during weathering and its relationship to the finding region of the meteorite.

## STUDIED METEORITES

In order to investigate the possible effects of meteorite recovery site on the concentrations of REE

during weathering, OCs from two hot deserts with markedly different climatic settings were selected. The characteristics of the studied samples are described in the forthcoming subsections.

### Ordinary Chondrites from Lut Desert

With an annual precipitation of  $\leq 28$  mm  $\text{yr}^{-1}$  (in the marginal regions) and maximum ground temperature of up to  $70^\circ\text{C}$  (in the desert) (Djamali et al. 2011; Mildrexler et al. 2011) along with the increasing number of meteorite finds (Pourkhorsandi and Mirnejad 2013; Pourkhorsandi et al. 2016), the Lut desert in Iran has the characteristics of a suitable place for preserving meteorites.

For this study, we selected a set of OCs from different regions of Lut whose characteristics and finding places are presented in Table 1 and Fig. 1a. The analyzed samples comprise meteorites from LL, L, and H groups with petrologic types varying between three and six. Intense oxidation of Fe-Ni metal grains ( $>95\%$  of them) is the prevalent characteristic of these meteorites (Fig. 2a). According to the method of Wlotzka (1993), weathering degrees of these samples are between W3 and W5. Iron oxides/oxyhydroxides occur as individual patches, veins (along and/or perpendicular to cracks), and lamellar structures occupying former Fe-Ni metal, troilite, and silicate locations. Reflected light optical microscopy reveals the alteration of troilite to pyrite/marcasite—rather than Fe oxides/oxyhydroxides—in some of these samples.

### Ordinary Chondrites from Atacama Desert

The central depression of the Atacama, the driest and oldest hot desert on Earth (Hartley et al. 2005), is responsible for the preservation of very old surfaces containing unique meteorite accumulation areas (e.g., San Juan, and El Médano). Other meteorite dense collection areas (DCAs) occur in the domain of the coastal range, known as La Yesera and Pampa de Mejillones DCAs, with a different climatic condition, due to the presence of oceanic aerosols and morning mist coming from the Pacific Ocean. Extreme aridity and the surface stability of the Atacama lower the rate of chemical weathering of meteorites compared to those reported from other hot deserts (Gattacceca et al. 2011)—in this case Lut. Unlike the relatively low number of recovered meteorites from Lut, the Atacama hosts the highest number of meteorites per surface area reported so far (Gattacceca et al. 2011; Hutzler et al. 2016).

In this work, we used samples previously classified and described by Valenzuela (2011) from La Yesera and Pampa de Mejillones DCAs, which are located near to

Table 1. Meteorites studied in this work.

Meteorite	Type	Weathering degree	Terrestrial age (ka) <sup>a</sup>
Ordinary chondrites from Atacama			
La Yesera 01	H6	W3	10.79 ± 1.56
La Yesera 02	LL5	W2	25.44 ± 4.45
La Yesera 03	L4	W4	16.98 ± 2.47
La Yesera 04	L6	W3	34.07 ± 1.92
Pampa (a)	L6	W2	25.08 ± 1.46
Pampa (b)	L4/5	W3	21.29 ± 2.45
Pampa (c)	L4	W4	13.89 ± 2.08
Pampa (d)	L5	W2/3	14.18 ± 1.91
Pampa (g)	L5	W3	14.34 ± 1.62
PdM 02	H5	W3	3.86 ± 1.36
PdM 04	L6	W4/5	23.98 ± 4.40
PdM 07	L6	W4	>27.9
PdM 10	L5	W3	18.06 ± 1.90
PdM 11	L5	W5	4.35 ± 1.34
PdM 12	H4/5	W5	>34.3
PdM 14	L/LL4-6	W2	12.54 ± 1.95
San Juan 01	L5	W2	>28.1
San Juan 02	H6	W2	19.44 ± 1.69
EM 49	H4	W3	1890 ± 80
Estacion Imilac	H5	W1	–
Cobija	H6	W1	19.7 ± 4.20
Rencoret	H6	W3	25.3 ± 6.4
Lutschaunig's Stone	L6	W1	9.2 ± 1.6
CeC 06	L6	W3	2345 ± 80
Paposo	LL6	W2	>49.2
Ordinary chondrites from Lut			
Lut 01	H5	W3	–
Lut 03	L3	W3	–
Lut 06	LL3	W3	–
Lut 08	H4	W5	–
Lut 09	H4	W4	–
Kerman 01	H5	W4	–
Kerman 02	L6	W3	–
Kerman 03	L5	W2	–
Shahdad	H5	W4	–
Ordinary chondrites from Sahara			
Aridal 06	H6	W4	–

<sup>a</sup>Terrestrial age data from Valenzuela (2011) by <sup>14</sup>C, and Hutzler (2015) by <sup>36</sup>Cl/<sup>41</sup>Ca. PdM, Pampa de Mejillones; EM, El Médano; CeC, Caleta el Cobre.

the Pacific coast (Fig. 1b) and are affected by the considerable presence of coastal fog, which means a higher amount of NaCl in the regional soils. Most of these samples were found on sand dunes, in the context of Pleistocene sandstones with paleo-coastal lines, with the presence of carbonates in the soils.

To compare the chemical behavior of meteorites from different regions of the same desert, samples from San Juan and El Médano DCAs were chosen. These two regions are different from the coastal regions, as they are situated in much drier climatic conditions. Volcanic rocks

cover the surface of these areas, with remarkable presence of gypsum and anhydrite as caliche layers.

Similar to the Lut meteorites, OCs from LL, L, and H groups are included (Table 1). The selected meteorites exhibit weathering degrees between W1 and W5 with a mode at W3. In comparison to meteorites from other hot deserts, Atacama samples have long terrestrial ages. For instance, based on the <sup>36</sup>Cl/<sup>41</sup>Ca terrestrial age determinations, Hutzler (2015) presented ages of 1900 ± 80 ka and 2590 ± 100 ka for El Médano 049 and Caleta el Cobre 006 meteorites, respectively. Gattacceca et al. (2011) and Valenzuela (2011) also reported relatively old terrestrial ages (see Table 1 and see the Discussion section).

In addition to these meteorites, an OC from the Sahara (Aridal 006) was selected for analysis and comparison to these measurements.

## METHODOLOGY

To prepare the powder used for measuring the abundances of REE in bulk meteorites, samples were examined under a stereomicroscope to check for the presence of any terrestrial material (e.g., cemented or enclosed sand grains inside cracks, fusion crust, desert varnish, and other sedimentary crusts). Each of these materials shows different geochemical behaviors and their contamination can affect the analysis result. For instance, desert varnish tends to show high REE contents (Goldsmith et al. 2014). Thus, care was taken to separate the mentioned products from the exterior and interior of the meteorites. Samples were taken from the interior and from the freshest parts of the meteorites, away from the fusion crust (where present) and the mentioned terrestrial products. To have a representative aliquot of whole rock composition, approximately 1.5–2 g of each meteorite was finely hand-ground in an agate mortar and then carefully homogenized.

Inductively coupled plasma–mass spectrometry (ICP–MS) instruments (VG PQII Plus STE and Perkin-Elmer NexION<sup>®</sup> 300x) at the Dipartimento di Scienze della Terra of the Università di Pisa were used to determine the trace elements (herein REE, Sr, and Ba) contents of the meteorite samples. About 50–100 mg of each powder was dissolved in a mixture of HF and HNO<sub>3</sub> on a hot plate at ~120° C inside screw-top perfluoroalkoxy (PFA) vessels. It is possible that this dissolution procedure could incompletely dissolve celestine (SrSO<sub>4</sub>) or barite (BaSO<sub>4</sub>), which might eventually occur in the samples. However, we always obtained very clear final sample solutions. The sample solutions were then spiked with Rh, Re, and Bi as internal standards (20 ng mL<sup>-1</sup> in the final solutions) and diluted to 50 mL in polypropylene flasks. In each step of sample preparation, Mill-Q<sup>®</sup> purified water




Fig. 1. a) Map of the central Lut Desert and the find locations of the meteorites studied. b) Map of the Atacama Desert and the location of the dense collection areas (DCAs) from which meteorites were used in this study.

(18.2 M cm), and ultrapure HF and HNO<sub>3</sub> were used. The correction procedure included 1) blank subtraction, 2) instrumental drift correction using internal standardization and repeated (every five samples) analysis of a drift monitor, and 3) oxide-hydroxide interference correction. The geochemical reference samples with basaltic composition WS-E, PM-S, and BIR-1, and the Allende chondrite reference sample (USNM 3529, split 20, and position 22) were dissolved and analyzed along with the unknown samples to check the accuracy of the results. In Table 2, the results are reported for the four analyzed reference sample with literature values, and the detection limits for each analyte were calculated as six times the standard deviation of the blank counts. The analytical precision is between 5 and

10% RSD for elements with concentrations >0.5 μg g<sup>-1</sup> and between 10 and 20% RSD for elements with concentrations <0.5 μg g<sup>-1</sup>.

To investigate the effect of surface soil chemical composition and to observe any possible significant anomaly, representative soil samples collected from a hilltop close to Estacion Catalina (25°12.673'S, 69°41.894'W) were used in this study. This region which occurs in the Catalina DCA is composed of igneous rocks with the presence of caliche layers and has very similar lithology to San Juan and El Médano DCAs. Samples containing gravel and soils were collected from depths of 0 to 0.5 cm and 0.5 to 5 cm. The gravels and soils (<2 mm) were then sieved and separated for subsequent analyses. Soil concentrate samples were pulverized to




Fig. 2. Mosaic pictures of (a) Kerman 001 and (b) El Médano 049 prepared in reflected polarized light show the different weathering types of the meteorites. Kerman 001 contains clear parallel veins filled with weathering products and represents much stronger weathering than El Médano 049 whose Fe-Ni metal and troilite are not completely altered. (Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).)

85% passing 75  $\mu\text{m}$  in tungsten-carbide mill. Total chemistry for major and trace elements was determined using ICP-MS and ICP-AES after a lithium metaborate fusion of each soil sample at the ALS Geochemistry<sup>©</sup>.

Rare earth elements anomalies represent deviations from neighboring elements based on the expected change in REE abundances as a function of atomic number and ionic radii (Lipin and McKay 1989). To calculate Ce and Eu anomalies, the values were normalized to the corresponding OC group (LL, L, and H) by using the following equations (Dauphas and Pourmand 2015):

$$\text{Ce/Ce}^* = \text{Ce}_N / (\text{La}_N^{0.48} \times \text{Pr}_N^0 \text{Pr}_N^{0.52}) \quad (1)$$

$$\text{Eu/Eu}^* = \text{Eu}_N / (\text{Sm}_N^{0.45} \times \text{Gd}_N^0 \text{Gd}_N^{0.55}) \quad (2)$$

where  $N$  stands for the normalization of REE to mean OC composition.

The contents of Sr and Ba have been shown to be suitable chemical proxies to evaluate the amount of chemical weathering and differentiate the meteorite collection regions (Al-Kathiri et al. 2005; Folco et al. 2007; Zurfluh et al. 2012).

## RESULTS

### Meteorite Samples

Rare earth elements, Sr, and Ba concentrations of the meteorites studied are reported in Table 3. The REE data of each group are normalized to the corresponding mean composition of Wasson and Kallemeyn (1988). Masuda–Coryell diagrams of the studied meteorites together with the data of OCs from Oman (Al-Kathiri et al. 2005), UAE (Hezel et al. 2011), Sahara desert (Folco et al. 2007; Saunier et al. 2010), and Europe (Folco et al. 2007) are shown in Fig. 3. Both enrichment and depletion elemental patterns

compared to the corresponding OC mean composition can be seen in the diagrams.

$\Sigma\text{REE}$  values in the Atacama and Lut samples have a range of 3.55–5.77 (avg. 4.46; std. 0.79) ppm and 2.98–3.76 (avg. 3.42; std. 0.30) ppm for H, 2.97–6.45 (avg. 3.97; std. 0.88) ppm and 3.26–2.93 (avg. 3.63; std. 0.28) ppm for L, and 3.21–9.82 (avg. 5.95; std. 2.81) ppm (just for Atacama) for LL OCs, respectively (Table 3). These data show a higher amount of REEs in Atacama meteorites than those of Lut.

A closer look at the normalized REE patterns (Fig. 3) reveals an enrichment in La for some samples. This can be up to  $2.64 \times \text{H}$  in El Médano 049, with an average amount of  $1.79 \times \text{H}$  ( $n = 8$ ) and  $1.22 \times \text{H}$  ( $n = 5$ ) for Atacama and Lut meteorites, respectively. Likewise, La is enriched up to  $2.98 \times \text{L}$  in Caleta el Cobre 006, which is an L OC. Average  $\text{La}_N$  amounts of Atacama and Lut samples are  $1.40 \times \text{L}$  ( $n = 14$ ) and  $1.15 \times \text{L}$  ( $n = 3$ ), respectively. Among the studied meteorites, Paposo exhibits a remarkable La enrichment ( $4.05 \times \text{LL}$ ).

The general shape of the normalized spider diagrams (Fig. 3) indicates the occurrence of a fractionation between light REEs (LREEs) and heavy REEs (HREEs). Figure 4 displays the values of  $\text{La}_N/\text{Lu}_N$  (as an indicative of LREE/HREE fractionation) versus  $\text{La}_N$  for H and L OCs from different regions. Most of the samples show LREE/HREE ratios of more than 1. The Atacama and Sahara samples tend to represent higher degrees of REE fractionation than Lut and UAE samples. The  $\text{Gd}_N/\text{Yb}_N$  (as an indicator of MREE/HREE fractionation) ratios for the studied samples are mostly less than 1, but as the values are near the error zone, they will not be discussed further.

Cerium and Eu show both negative and positive anomalies (Table 3, Fig. 3). In H OCs, the variation range of  $\text{Ce/Ce}^*$  and  $\text{Eu/Eu}^*$  values for Atacama samples (0.60–1.69 and 0.56–1.08, respectively) is

Table 2. Concentrations ( $\mu\text{g g}^{-1}$ ) of Sr, Ba, and REE in the four geochemical reference materials analyzed by ICP-MS along with the unknown samples. Literature values for these reference materials are also reported. The last column reports the detection limits ( $\mu\text{g g}^{-1}$ ) of the used analytical method.

Element	Allende (this work)	Allende (literature) <sup>a</sup>	BIR-1		WS-E (this work)	WS-E (literature) <sup>c</sup>	PM-S (this work)	PM-S (literature) <sup>c</sup>	Detection limit
			(this work)	BIR-1 (literature) <sup>b</sup>					
Sr	14.5	12 ± 3	122	108	450	410 ± 5	307	280 ± 5	0.5
Ba	4.5	4 ± 1	<10	7	342	338 ± 6	139	148 ± 3.2	0.6–10
La	0.49	0.52 ± 0.04	0.62	0.62	27.6	27 ± 1.1	2.63	2.8 ± 0.17	0.02
Ce	1.24	1.33 ± 0.08	1.90	1.95	60	61 ± 1.4	6.5	6.8 ± 1.25	0.06
Pr	0.21	0.21 ± 0.01	0.39	0.38	8.0	7.8 ± 0.4	1.05	1.08 ± 0.16	0.007
Nd	1.01	0.99 ± 0.03	2.45	2.5	33.3	33 ± 0.7	5.4	5.5 ± 0.25	0.03
Sm	0.33	0.34 ± 0.02	1.10	1.1	8.8	8.8 ± 0.3	1.78	1.75 ± 0.05	0.02
Eu	0.09	0.11 ± 0.01	0.53	0.54	2.31	2.25 ± 0.04	1.04	1.07 ± 0.04	0.05
Gd	0.39	0.42 ± 0.02	1.96	1.85	7.4	7.2 ± 0.23	2.05	2 ± 0.1	0.004
Tb	0.07	0.081 ± 0.010	0.38	0.36	1.10	1.1 ± 0.04	0.35	0.36 ± 0.02	0.002
Dy	0.46	0.42 ± 0.03	2.78	2.5	6.5	6 ± 0.16	2.18	2 ± 0.1	0.005
Ho	0.10	0.10 ± 0.01	0.59	0.57	1.18	1.2 ± 0.06	0.42	0.42 ± 0.03	0.006
Er	0.29	0.29 ± 0.01	1.81	1.7	3.17	3 ± 0.11	1.19	1.1 ± 0.06	0.01
Tm	0.05	0.0572	0.26	0.26	0.42	0.43 ± 0.03	0.16	0.17 ± 0.01	0.003
Yb	0.31	0.30 ± 0.02	1.70	1.65	2.56	2.5 ± 0.1	0.96	1 ± 0.05	0.007
Lu	0.04	0.052 ± 0.006	0.26	0.26	0.36	0.37 ± 0.01	0.14	0.15 ± 0.01	0.002

<sup>a</sup>Jarosewich et al. (1987).

<sup>b</sup>Govindaraju (1994).

<sup>c</sup>Govindaraju (1995).

broader than those of Lut (0.92–0.96 and 0.96–1.29, respectively). In general, no pronounced difference in the amounts of Ce/Ce\* and Eu/Eu\* between different OC groups exists. A remarkable positive Ce anomaly is evident in Cobija (H6) and Lutschaunig's Stone (L6) (Figs. 3a and 3b). Note that these two meteorites are the least weathered samples (W1) studied so far from the Atacama.

The negative Tm anomaly among the analyzed samples (Fig. 3) is similar to the Tm anomaly in the CI-normalized patterns described from ordinary chondrites (e.g., Dauphas and Pourmand 2015). However, it is not the case here for OC-normalized samples. It should be noted that the seemingly more negative anomalies in H OCs relative to that of L OCs are not real but occurred as the result of the normalization errors.

Logarithmic plot of Sr versus Ba for the OCs from Atacama, Oman, and Sahara is shown in Fig. 5. Different chemical trends are particularly more pronounced among those from the Atacama and Oman.

### Atacama Soil Samples

The major and trace element composition of the analyzed soil samples are reported in Table 4. As reflected in their SiO<sub>2</sub> content, the parent rock material is basaltic/andesitic but the soil consists of more acidic

(granitic) material than the gravels, which is transported by eolian processes from neighboring lithologies. Both CI-normalized REE patterns of gravels and <2 mm fractions of the soils are fractionated, showing LREE enrichments and a relatively flat HREE (Fig. 6). The LREE enrichment in the <2 mm soil is higher than the parent rock gravels ( $\text{La}_N/\text{Lu}_N$  mean values 9.1 versus 6.7), which is related to the weathering of the parent rock and the higher mobility of LREE into the soils during rock weathering process (Kabata-Pendias and Pendias 2001). The same process explains the negative Eu anomaly of the soils compared to the gravels and more resistance of Eu hosting grains during weathering. However, the chemical differences mentioned between the bedrock and the soil could also affect these patterns.

Comparison of the analyzed samples with the REE composition of upper crust composition (UCC) (Kemp and Hawkesworth 2004) and mean terrestrial soil composition references (Govindaraju 1994; Kabata-Pendias and Pendias 2001) shows their similarity, which implies a well-mixed texture from various bedrocks. In contrast to the Atacama meteorites that generally show higher REE concentrations compared to the other hot desert meteorites, the soils show REE chemical compositions comparable to the UCC and mean soil composition and without any significant compositional difference.

Table 3. Strontium, Ba, and bulk REE composition ( $\mu\text{g g}^{-1}$ ) and the calculated ratios of the studied meteorites.

Meteorite	Sr	Ba	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	$\Sigma\text{REE}$	Ce*/Ce*	Eu*/Eu*	Tm*/Tm*	La <sub>N</sub> /		Gd <sub>N</sub> /		Sr/ Ba	
																					Lu <sub>N</sub>	Sm <sub>N</sub>	Yb <sub>N</sub>	Ba		
La Yessera 01	30.6	38	0.76	1.63	0.22	1.01	0.27	<0.05	0.31	0.05	0.38	0.08	0.23	0.03	0.22	0.03	5.3	0.92	0.56	0.86	2.27	1.73	0.98	0.79	0.79	
La Yessera 02	26.3	50	0.54	1.54	0.18	0.81	0.23	0.10	0.31	0.05	0.39	0.08	0.25	0.04	0.25	0.03	4.8	1.08	1.28	1.13	1.73	1.52	0.90	0.53	0.53	
La Yessera 03	37	12.6	0.45	1.12	0.17	0.86	0.26	<0.05	0.29	0.05	0.38	0.09	0.27	0.04	0.25	0.05	4.3	0.90	0.58	0.92	1.04	1.07	0.83	2.99	2.99	
La Yessera 04	12.4	7.5	0.31	0.81	0.13	0.61	0.20	0.07	0.25	0.05	0.31	0.08	0.21	0.03	0.21	0.03	3.3	0.93	1.03	0.87	0.93	0.97	0.84	1.67	1.67	
Pampa (a)	18.1	4.1	0.30	0.85	0.12	0.61	0.21	0.10	0.24	0.05	0.33	0.08	0.22	0.03	0.23	0.03	3.4	0.99	1.41	0.88	1.16	0.93	0.75	4.41	4.41	
Pampa (b)	21.1	5.0	0.29	0.75	0.12	0.52	0.23	0.07	0.23	0.04	0.29	0.07	0.19	0.03	0.20	0.03	3.06	0.91	0.97	0.93	1.14	0.81	0.84	4.21	4.21	
Pampa (c)	31.8	14.6	0.40	0.94	0.15	0.70	0.23	0.09	0.29	0.05	0.35	0.08	0.26	0.04	0.25	0.04	3.9	0.86	1.06	0.89	1.17	1.10	0.81	2.18	2.18	
Pampa (d)	21.3	4.8	0.26	0.70	0.11	0.59	0.19	<0.05	0.24	0.04	0.31	0.07	0.21	0.03	0.19	0.03	3.0	0.95	0.75	0.92	0.87	0.84	0.87	4.39	4.39	
Pampa (g)	38.6	42	0.55	1.08	0.17	0.82	0.26	<0.05	0.32	0.05	0.35	0.09	0.26	0.03	0.23	0.04	4.2	0.81	0.55	0.83	1.39	1.31	0.99	0.92	0.92	
PdM 02	17.6	5.9	0.43	1.22	0.17	0.73	0.23	0.07	0.25	0.05	0.31	0.08	0.23	0.03	0.23	0.03	4.1	1.05	0.95	0.84	1.45	1.19	0.76	2.96	2.96	
PdM 04	23.6	8.3	0.39	0.97	0.14	0.68	0.21	0.08	0.25	0.05	0.31	0.07	0.20	0.03	0.21	0.03	3.6	0.92	1.07	0.97	1.24	1.16	0.85	2.82	2.82	
PdM 07	42	66	0.66	1.13	0.20	0.92	0.25	0.10	0.32	0.06	0.37	0.08	0.25	0.03	0.23	0.03	4.6	0.70	1.13	0.88	1.91	1.65	0.98	0.64	0.64	
PdM 10	26.1	5.3	0.29	0.88	0.12	0.55	0.21	0.08	0.26	0.05	0.31	0.07	0.19	0.03	0.23	0.03	3.3	1.06	1.10	0.85	0.97	0.90	0.80	4.91	4.91	
PdM 11	11.3	5.1	0.32	0.82	0.12	0.62	0.22	0.07	0.27	0.05	0.33	0.08	0.24	0.04	0.22	0.03	3.4	0.94	0.94	1.02	1.19	0.90	0.85	2.19	2.19	
PdM 12	39	21.3	0.43	1.07	0.14	0.68	0.20	0.07	0.25	0.05	0.33	0.07	0.23	0.03	0.21	0.02	3.8	1.00	1.08	0.84	1.69	1.36	0.80	1.85	1.85	
PdM 14	18.4	12.5	0.27	0.92	0.12	0.59	0.19	<0.05	0.25	0.05	0.30	0.08	0.200	0.03	0.19	0.03	3.2	1.11	0.74	1.08	0.92	0.88	0.94	1.47	1.47	
San Juan 01	71	52	0.57	1.17	0.18	0.77	0.24	0.11	0.27	0.05	0.34	0.07	0.20	0.03	0.21	0.03	4.3	0.83	1.35	1.00	1.77	1.48	0.89	1.37	1.37	
San Juan 02	39	9.4	0.38	0.91	0.14	0.69	0.18	<0.05	0.24	0.05	0.32	0.08	0.24	0.03	0.21	0.03	3.5	0.92	0.77	0.75	1.12	1.28	0.80	4.24	4.24	
EM 49	45	270	0.78	1.34	0.25	1.22	0.31	0.10	0.44	0.07	0.49	0.10	0.30	0.04	0.27	0.04	5.8	0.69	0.91	0.82	1.87	1.56	1.13	0.17	0.17	
Estacion	11.8	9.8	0.44	1.03	0.16	0.71	0.21	0.07	0.27	0.05	0.34	0.08	0.24	0.03	0.24	0.03	3.9	0.90	0.97	0.82	1.35	1.29	0.78	1.20	1.20	
Imilac																										
Cobija	15.8	6.2	0.33	1.48	0.12	0.62	0.19	0.07	0.25	0.04	0.33	0.07	0.21	0.03	0.19	0.03	4.0	1.69	1.05	0.85	1.03	1.08	0.89	2.57	2.57	
Rencoret	26.1	12.9	0.68	1.62	0.22	1.01	0.27	0.09	0.32	0.05	0.35	0.08	0.25	0.04	0.24	0.03	5.2	0.97	0.97	0.81	2.04	1.58	0.89	2.01	2.01	
Lutschaunig's	10.4	4.2	0.35	1.83	0.14	0.69	0.22	0.07	0.30	0.05	0.38	0.09	0.25	0.03	0.23	0.03	4.7	1.84	0.92	0.89	1.06	0.98	0.92	2.48	2.48	
Stone																										
CeC 06	72	326	0.92	1.89	0.28	1.22	0.32	0.10	0.40	0.07	0.48	0.10	0.31	0.04	0.28	0.04	6.4	0.85	0.87	0.90	2.46	1.80	1.01	0.22	0.22	
Paposo	41	70	1.48	3.49	0.50	1.88	0.46	0.10	0.48	0.08	0.52	0.11	0.31	0.05	0.30	0.05	9.8	0.88	0.68	0.99	2.92	2.06	1.15	0.58	0.58	
Lut 01	62	<10	0.30	0.72	0.11	0.50	0.18	0.08	0.23	0.04	0.30	0.06	0.20	0.03	0.20	0.03	2.98	0.92	1.28	0.82	1.02	1.06	0.80	6.2	6.2	
Lut 03	91	<10	0.34	0.81	0.12	0.58	0.18	0.07	0.26	0.04	0.33	0.07	0.20	0.03	0.19	0.03	3.26	0.90	1.01	0.94	1.20	1.20	0.94	9.1	9.1	
Lut 06	126	20.8	0.37	0.92	0.14	0.68	0.23	0.08	0.29	0.06	0.37	0.08	0.24	0.04	0.26	0.03	3.8	0.86	1.01	0.96	1.10	1.04	0.82	6.1	6.1	
Lut 08	72	<10	0.31	0.79	0.12	0.62	0.20	0.07	0.26	0.05	0.32	0.07	0.20	0.03	0.20	0.03	3.28	0.96	0.96	0.86	1.01	0.99	0.87	7.1	7.1	
Lut 09	497	14.3	0.33	0.84	0.12	0.62	0.19	0.07	0.25	0.05	0.31	0.07	0.20	0.03	0.20	0.03	3.3	0.95	1.08	0.86	1.03	1.08	0.85	3.4	3.4	
Kerman 01	165	<10	0.38	0.93	0.14	0.70	0.21	0.08	0.30	0.05	0.36	0.08	0.22	0.03	0.22	0.03	3.7	0.93	0.99	0.87	1.16	1.12	0.96	16.5	16.5	
Kerman 02	33	<10	0.35	0.90	0.14	0.66	0.22	0.08	0.29	0.05	0.38	0.08	0.24	0.03	0.23	0.04	3.7	0.91	1.00	0.89	1.07	1.02	0.88	3.3	3.3	
Kerman 03	41	<10	0.43	1.00	0.14	0.70	0.22	0.08	0.30	0.05	0.37	0.08	0.24	0.03	0.24	0.04	3.9	0.92	1.06	0.87	1.27	1.23	0.89	4.1	4.1	
Shahdad	141	<10	0.47	1.04	0.14	0.66	0.20	0.07	0.25	0.05	0.33	0.07	0.20	0.03	0.21	0.03	3.8	0.93	1.07	0.85	1.63	1.46	0.79	14.1	14.1	
Aridal 06	17.9	23.7	0.29	0.70	0.10	0.49	0.16	0.08	0.21	0.04	0.28	0.06	0.18	0.03	0.18	0.03	2.84	0.94	1.49	0.82	1.01	1.14	0.80	0.76	0.76	

PdM, Pampa de Mejillones; EM, El Médano; CeC, Caleta el Cobre.

## DISCUSSION

Our results show that terrestrial weathering may significantly modify the whole rock REE distribution of OCs from hot deserts. However, this modification varies among different deserts. Atacama OCs have higher concentrations of REE and their compositions have been affected to a greater extent than those of Lut.

### REE Mobilization During Meteorite Weathering: A Concise Review

Some of the first plausible evidence for REE mobilization during terrestrial weathering of meteorites was reported by Masuda et al. (1977) and Masuda and Tanaka (1978) on samples from Antarctica. These authors, however, considered the differences in the abundance of REE between the internal and external portions of meteorite samples to be intrinsic and primary rather than produced by terrestrial effects. Further studies on Antarctic achondrites attributed the modified REE compositions to cold-desert weathering (Shimizu et al. 1983; Floss and Crozaz 1991; Mittlefehldt and Lindstrom 1991; Shinonaga et al. 1994; Kagi and Takahashi 1998; Swindle et al. 1998). The most prevalent results of weathering effects are LREE-enriched patterns with notable Ce (and in some cases Eu) anomalies produced by aqueous leaching. In a process proposed by Mittlefehldt and Lindstrom (1991) and confirmed by Floss and Crozaz (1991), during the exposure of a meteorite to atmosphere and its heating by the Sun at the surface in Antarctica, surrounding meltwater can penetrate the meteorite through cracks and fractures. In equilibrium with atmospheric CO<sub>2</sub>, this fluid becomes a weak carbonic acid solution which could leach the minerals such as Ca-phosphates and causes removal of trivalent REEs. In addition to being in +3 state, Ce and Eu occur as tetravalent and divalent ions, respectively. As Ce<sup>+4</sup> is less soluble than the +3 REEs and Eu is largely retained in the +2 state in plagioclase (which is relatively unaffected by weathering), these two elements are preferentially retained in the sample and thus show their corresponding anomalies. Another study by Kagi and Takahashi (1998) showed a correlation between Ce anomalies and absorbed water content in Antarctic lunar meteorites that demonstrated the terrestrial origin of these anomalies.

Following the studies on the Antarctic meteorites, attempts were made to investigate the possible weathering effects on hot desert achondrites (both in mineral and whole rocks). In order to monitor the weathering of recently fallen meteorites, Barrat et al. (1999) analyzed fragments of Tatahouine diogenite

recovered just 63 yr after its fall. By comparing the REE composition with those of the fragments recovered immediately after the fall, elevated amounts of LREE were found in one of the fragments. This study revealed the disturbance of trace element composition of meteorites in relatively short time spans in a terrestrial environment. Crozaz et al. (2003) evaluated the mineral composition of different achondrite groups by secondary ion mass spectrometry (SIMS) and attributed the elevated levels of LREE in olivine and low-Ca pyroxene to terrestrial contamination. However, Barrat et al. (2003) did not find any significant weathering effects on whole rock REE composition of eucrites from the Sahara desert. A similar conclusion was made during a study of NWA 4872 brachinite by Hyde et al. (2014).

### Effects of Weathering on the REE Composition of Atacama and Lut OCs

The aliquots analyzed in our work were from a large mass (about 2 g) of well-mixed powder, so the formation of the observed REE patterns by a nugget effect of REE-rich components is unlikely. Deviations of more than ±10% from the mean composition of the corresponding meteorite groups assure us that the enrichments/depletions are significant.

Unlike the Atacama OCs, those from Lut do not show profound changes in ΣREE (Table 3). Chemical modification of a meteorite during weathering, including changes in REE content, is controlled by several factors such as primary chemical composition, modal composition, degree of recrystallization, shock stage, terrestrial age, and locality (Crozaz et al. 2003). Rare earth elements which are mostly concentrated in Ca-phosphates and silicates (Ebihara and Honda 1984) are released during the weathering of these minerals. This process is facilitated by formation of sulfuric acid inside the meteorite as a result of troilite (FeS) oxidation in contact with meteoric water (e.g., Bland et al. 2006). With the development of primary mineral dissolution, REEs tend to concentrate in veins filled with iron oxides/oxyhydroxides (Thiagarajan and Aeolus Lee 2004).

In contrast to meteorites from Antarctica, which release their REE during weathering on ice, an increase in REE concentration is expected for hot desert meteorites as a consequence of residing on a relatively REE-rich soil surface (Crozaz and Wadhwa 2001). Once a meteorite with its relatively high initial porosity falls on a hot desert surface, soil salts dissolved by water infiltrate into it by capillary forces triggered by temperature fluctuations (Zurfluh 2012). In addition, wind activity, burial in soil, and desert varnish




Fig. 3. Normalized REE spider diagrams of (a) H, (b) L, and (c) LL ordinary chondrites. For comparison, the data of OCs from Oman (the average values for different age classes) (Al-Kathiri et al. 2005), UAE (Hezel et al. 2011), Sahara desert (Folco et al. 2007; Saunier et al. 2010), and Europe (Folco et al. 2007) are used along the data of this work. PdM is the abbreviated form of Pampa de Mejillones. (Color figure can be viewed at wileyonlinelibrary.com.)




Fig. 4.  $La_N/Lu_N$  versus  $La_N$  for H and L OCs from different regions. For comparison, the data of OCs from Oman (Al-Kathiri et al. 2005), UAE (Hezel et al. 2011), Sahara desert (Folco et al. 2007; Saunier et al. 2010), and Europe (Folco et al. 2007) are used along the data of this work. (Color figure can be viewed at wileyonlinelibrary.com.)




Fig. 5. Sr versus Ba logarithmic plot showing OCs from various areas. The data of OCs from Oman (Al-Kathiri et al. 2005), Sahara desert (Folco et al. 2007; Saunier et al. 2010), and Europe (Folco et al. 2007) are shown along the data of this work. Some samples from Lut with Ba <10 ppm are also included. (Color figure can be viewed at wileyonlinelibrary.com.)

formation introduce terrestrial minerals containing REE, or cause the modification of REE patterns of primary minerals (Crozas et al. 2003). We suggest that the notable increase in  $\Sigma$ REE contents of the Atacama OCs is caused by these kind of implementations.

Is the modification in REE concentration controlled by the weathering degree of a meteorite?

The data of Lut OCs (Tables 1 and 3) show that despite having high degrees of weathering, their  $\Sigma$ REE do not change as much as the Atacama samples. For example, Kerman 001 (W4), which was half-buried in a valley in the Kalout formation with a soil composed of clay, sand, evaporites, and carbonates (Farpoor and Krouse 2008) at the time of recovery

Table 4. Major and trace element chemical composition of soil and gravel sample from Estacion Catalina in the Atacama Desert. Trace element concentration unit is ppm.

	SV2 < 2 mm soil				SV1 gravels			
	CS-Surface-01	CS-Surface-02	CS-1-01	CS-1-02	CS-Surface	CS-1-01	CS-1-02	CS-5
Depth (cm)	0–0.5	0–0.5	0.5–5	0.5–5	0–0.5	0–0.5	0.5–5	>34
SiO <sub>2</sub>	59.8	60.4	58	58	53	52.5	57.8	52.5
Al <sub>2</sub> O <sub>3</sub>	14.65	14.65	14.6	14.65	17.35	17	16.6	17.15
Fe <sub>2</sub> O <sub>3</sub>	8.09	7.88	7.56	7.62	8.13	8.29	6.99	8.21
CaO	4.42	4.11	3.9	3.97	8.88	8.72	7.46	9.44
MgO	2.11	1.98	2.19	2.23	5.52	5.51	3.82	5.55
Na <sub>2</sub> O	2.96	2.84	2.48	2.53	2.94	2.8	2.75	3.04
K <sub>2</sub> O	2.42	2.37	2.31	2.31	1.17	1.12	1.3	1.11
Cr <sub>2</sub> O <sub>3</sub>	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
TiO <sub>2</sub>	1.19	1.16	1.08	1.1	1.05	1.06	0.98	1.07
MnO	0.14	0.16	0.12	0.13	0.13	0.14	0.15	0.13
P <sub>2</sub> O <sub>5</sub>	0.19	0.2	0.21	0.23	0.27	0.23	0.23	0.23
Total %	96.14	95.92	92.6	92.92	98.6	97.53	98.27	98.58
Rb	84.1	85.4	95	91.6	30.5	25.4	35.6	19.2
Sr	525	483	423	421	931	865	878	921
Ba	931	954	789	782	412	384	779	360
La	25.8	26.1	26.1	25.8	19.6	18.8	20.1	17.9
Ce	52	53	52.6	52.4	42.2	40.4	45.6	38.7
Pr	6.3	6.4	6.4	6.4	5.3	5.1	5.3	4.9
Nd	24.3	24.3	25	24.4	21.8	21.1	21.5	20.5
Sm	4.9	4.6	4.9	4.8	4.3	4.3	4.3	4.1
Eu	1.2	1.3	1.2	1.2	1.4	1.4	1.4	1.4
Gd	5	4.7	5	4.7	4.8	4.4	4.8	4.4
Tb	0.7	0.7	0.7	0.6	0.7	0.6	0.7	0.6
Dy	3.7	3.7	3.8	3.8	3.8	3.8	3.9	3.6
Ho	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
Er	2.3	2.2	2.3	2.2	2.2	2.2	2.3	2.1
Tm	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3
Yb	2.1	2.1	2.1	2	2	1.9	2.1	1.9
Lu	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
La <sub>N</sub> /Lu <sub>N</sub>	9.1	9.2	9.2	9.1	6.9	6.6	7.1	6.3

and exhibits well-developed parallel weathering veins (Fig. 2a), does not show any noticeable deviation from the mean composition. In comparison, the Atacama samples that are generally less weathered show higher  $\Sigma$ REE. For example, El Médano 049, which contains the highest  $\Sigma$ REE among the studied H OCs and was recovered from a deflated soil surface on granitoids, is also highly weathered (W3, Fig. 2b), although less than Kerman 001. Compared to samples from other regions, we see that three highly weathered H OCs from UAE (Hezel et al. 2011) and even weathered samples from Europe (Folco et al. 2007) show  $\Sigma$ REE contents similar to Lut samples. However,  $\Sigma$ REE is elevated in highly weathered Saharan OCs (analyzed by Saunier et al. 2010) and Aridal 006 (W4). Therefore, the meteorite weathering as shown by weathering degree (Wlotzka 1993) and veins filled with secondary products is not the only factor responsible for the  $\Sigma$ REE increase.

As for REEs, different chemical trends in Sr and Ba are also evident in meteorites from different regions (Figs. 5 and 7). Al-Kathiri et al. (2005) and Folco et al. (2007) showed that the weathering of meteorites in Oman and Sahara deserts (respectively) increases the Sr and Ba contents of meteorites. The carbonate basement and the Sr- and Ba-rich soils in Oman deserts are responsible for the further enrichments of these elements in the recovered meteorites from the region. In some cases from Oman, high concentrations of Sr and Ba in the soil lead to the formation of secondary celestine and barite inside the meteorites (Al-Kathiri et al. 2005). Meteorites from Atacama exhibit a different trend compared to Oman meteorites (Fig. 5). The majority of collecting site of the meteorites from Atacama is mostly made of volcanics and evaporites (Valenzuela 2011), which usually have lower concentrations of Sr and Ba than carbonates. The difference in the soil type can explain different trends in the Sr budget. Compared to




Fig. 6. REE chemical composition of Estacion Catalina soil composition compared to the mean terrestrial soil compositions (Govindaraju 1994; Kabata-Pendias and Pendias 2001). The Atacama H and L OCs show LREE enrichment in relation to falls in an attempt to reach equilibrium during the hot desert terrestrial weathering. (Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).)

the Omani and Atacama OCs, the samples from Lut desert have very low Ba contents (<10 ppm). On the other hand, Sr concentrations of Lut OCs are in the range of Omani meteorites. Considering Sr and La contents (Fig. 7), again we see difference in the chemical composition of meteorites from different regions. These differences can be used as proxies which allow distinguishing between Atacama meteorites from Oman and Lut. Al-Kathiri et al. (2005) reported a positive correlation between weathering degree and Sr, Ba concentration for Omani OCs; however, we do not see any correlation between these parameters for the meteorites from Atacama and Lut samples.

The Atacama Desert has exceptionally high meteorite concentrations compared to other hot deserts. In a study of 22 meteorites, Gattacceca et al. (2011) found half of the meteorites with terrestrial ages older than 20 ka, which is much older than the age ranges reported from other hot deserts. We may hypothesize that the generally high terrestrial age of the Atacama meteorites is the main reason for their high  $\Sigma$ REE amounts. A similar effect of terrestrial age on REE

content can be seen in higher amounts of La in very old meteorites from Atacama and in samples with terrestrial ages of more than >30 kyr from Oman (Al-Kathiri et al. 2005) (Fig. 3a).

Most of the Atacama and some highly weathered Saharan OCs show differentiated REE patterns (Figs. 3 and 4). As the soil and sedimentary materials typically exhibit higher LREE contents (Aide and Aide 2012), as observed in our analyzed samples from the Atacama, the residence of a meteorite on a soil surface increases the LREE content of the meteorite (Dreibus et al. 2001; Crozaz et al. 2003; Al-Kathiri et al. 2005). The studied meteorites and their REE patterns, which tend to converge toward that of the surface material, are consistent with the assumption that soil has affected their chemical composition (Fig. 6). Taking into account the different soil types of the meteorite collecting sites (carbonate in Oman versus volcanic/evaporitic in Atacama and clay/sand in Lut), it can be deduced that nearly all of the meteorites represent approximately similar LREE-enriched patterns which form as a result of both (1) higher abundance of LREE




Fig. 7. Different chemical trends of OCs from different regions can be seen in the Sr versus  $La_N$  plot. The data of L and H OCs from Oman, in different age classes (Al-Kathiri et al. 2005), Sahara desert (Saunier et al. 2010; Folco et al. 2007), and Europe (Folco et al. 2007) are shown along the data of this work. (Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).)

in soils and (2) higher solubility of LREE (Aide and Aide [2012] and references therein).

Different clay and iron oxide/oxyhydroxide minerals display specific Ce and Eu anomalies. These minerals, which are the main components of soils, occur in different proportions based on the soil type; as a result, different Ce and Eu anomalies are exhibited in different soils (Laveuf and Cornu 2009). Although most of the Ce anomalies in our data are small and hardly significant, two slightly weathered (W1) meteorites (Cobija and Lutschaunig's Stone) display notable positive Ce anomalies (Figs. 3a and 3b) that might reveal the incipient phase of REE modification during weathering. Cerium is the most abundant REE in the average soil composition (Aide and Aide 2012). Once the meteorite falls, the initial physical soil contamination through cracks and fractures gives it a positive Ce anomaly. By developing the chemical weathering, trivalent REEs of the soil with higher solubility rates than  $Ce^{+4}$  infiltrate the meteorite, and most of the Ce oxides to  $Ce^{+4}$  and is preferentially retained in the soil. Progressive alteration of the meteorite and the addition of trivalent REEs remove the initial positive Ce anomaly. At higher residence times and more interaction of terrestrial contaminants with meteorite, the amount of La and Pr exceeds the Ce concentration and creates a negative Ce anomaly that is evident in the REE patterns of some meteorites with high terrestrial ages such as La Yesera

003, Caleta el Cobre 006, Paposo, and El Médano 049 (Fig. 3). Variation in Eu anomalies among studied meteorites can result from contamination by different soil components, although the presence of possible primary plagioclase should not be excluded as an explanation.

Putting these data together, we hypothesize that terrestrial age rather than weathering degree is the main governing factor of REE content and its modification in hot desert OCs, even though degree of weathering and terrestrial age at a given recovery site show a positive correlation (e.g., Bland et al. 2006). Indeed, the formation of secondary oxidation products decreases the porosity of the meteorite by filling the fractures and cracks which are the pathways for REE containing terrestrial fluid to circulate and solid materials to penetrate (Bland et al. 2006). The earlier this process initiates, the lower the possibility that a terrestrial material is able to affect the REE composition of the meteorite. However, in areas with low weathering rates (like Atacama), greater amounts of soil and fluid penetrate the meteorite and modify the composition readily.

We observe that as the meteorites from Atacama are older than those from Lut and other regions (Sahara, Oman), and also owing to low weathering rates in the former, more time has been available for the terrestrial environment to affect the chemistry of Atacama OCs by means of different interactions. The

terrestrial ages of Saharan meteorites used in this study are not determined; however, by comparing their weathering degree with terrestrial age of other meteorites collected from this desert (e.g., Welten et al. 2004), their terrestrial ages are expected to be high. The prevailing humid environment for UAE and Europe and thus higher weathering rates (Folco et al. 2007; Hezel et al. 2011) might have prevented the formation of a pronounced  $\Sigma$ REE increase in meteorites from these areas. We believe that the terrestrial age range of highly weathered Lut meteorites shall be in the range of UAE meteorites. The occurrence of archaeological sites in the surroundings of Lut (e.g., Muscarella 2001) points to wetter climatic conditions, at least periodically over the last 10 kyr compared to today. This is in agreement with the high weathering degree and the lack of a pronounced  $\Sigma$ REE change in the chemical composition of Lut OCs.

A large number of meteorites from Atacama are expected to be recovered in the future (Gattacceca et al. 2011; Hutzler et al. 2016). Hence, care must be taken into account while working on the REE composition of even “fresh-looking” Atacama meteorites.

## CONCLUSIONS

The chemical compositions of OCs from Atacama and Lut along with the published data from other hot desert areas reveal a difference in the effects of terrestrial weathering on REE distribution between these populations. Optimal conditions for meteorite preservation in the Atacama Desert (i.e., high stability of the surface and low meteorite weathering rates) provide enough time for the meteorites to interact efficiently with terrestrial materials. In contrast, higher weathering rates of the meteorites from other hot deserts (specifically Lut) prevent such interactions. As a consequence, the highly weathered meteorites from Lut desert have lower  $\Sigma$ REE contents than those of the moderately weathered meteorite from Atacama. In some cases, even the OCs with minor weathering from Atacama show modified REE patterns with notable positive Ce anomaly. These anomalies may have been created by the initial physical contamination (the addition of soil or dust directly into the fractures, rather than more time-consuming chemical weathering) and then by the addition of trivalent REEs. Notwithstanding the lack of terrestrial age data for Lut meteorites, it is envisaged on the basis of almost unaffected REE compositions and high weathering degrees that this area was wetter than today over the last tens of thousands of years and generally had a climate similar to the UAE and Oman deserts.

Future investigations on the mobilization of REE isotopes, which are being used frequently in cosmochemistry, are needed to estimate their behavior during weathering.

*Acknowledgments*—We thank the staff of the Cultural Office of the French Embassy in Tehran who supported the field trips in Iran and a Ph.D. thesis grant for the first author. Providing samples by Rodrigo Martínez, Edmundo Martínez, Enrique Stucken, and the Natural History Museum of London is appreciated. M.V. thanks the support by CONICYT-FONDECYT project No 3140562. H. P. thanks Morteza Djamali for his suggestions during preparation of the manuscript. We thank A. J. T. Jull for editorial handling, and P. Bland and an anonymous referee for their constructive reviews.

*Editorial Handling*—Dr. A. J. Timothy Jull

## REFERENCES

- Aide M. T. and Aide C. 2012. Rare earth elements: Their importance in understanding soil genesis. *ISRN Soil Science* 2012:1–11.
- Al-Kathiri A., Hofmann B. A., Jull A. J. T., and Gnos E. 2005. Weathering of meteorites from Oman: Correlation of chemical and mineralogical weathering proxies with 14 C terrestrial ages and the influence of soil chemistry. *Meteoritics & Planetary Science* 1239:1215–1239.
- Barrat J. A., Gillet P., Lesourd M., Blichert-Toft J., and Poupeau G. R. 1999. The Tatahouine diogenite: Mineralogical and chemical effects of sixty-three years of terrestrial residence. *Meteoritics & Planetary Science* 34:91–97.
- Barrat J. A., Jambon A., Bohn M., Blichert-Toft J., Sautter V., Göpel C., Gillet P., Boudouma O., and Keller F. 2003. Petrology and geochemistry of the unbrecciated achondrite Northwest Africa 1240 (NWA 1240): An HED parent body impact melt. *Geochimica et Cosmochimica Acta* 67:3959–3970.
- Bischoff A., Jersek M., Grau T., Mirtic B., Ott U., Kučera J., Horstmann A., Laubenstein M., Herrmann S., Randa Z., Weber M., and Heusser G. 2011. Jesenice-A new meteorite fall from Slovenia. *Meteoritics & Planetary Science* 46:793–804.
- Bland P. A., Berry F. J., and Pillinger C. T. 1998. Rapid weathering in Holbrook; an iron-57 Mossbauer spectroscopy study. *Meteoritics & Planetary Science* 33:127–129.
- Bland P. A., Zolensky M. E., Benedix G. K., and Sephton M. A. 2006. Weathering of chondritic meteorites. In *Meteorites and the early solar system II*, edited by Lauretta D. and McSween H. Y. Tucson, Arizona: University of Arizona Press. pp. 853–867. <http://www.lpi.usra.edu/books/MESSII/9041.pdf>.
- Boynton W. V. 1975. Fractionation in the solar nebula: Condensation of yttrium and the rare earth elements. *Geochimica et Cosmochimica Acta* 39:569–584.
- Boynton W., Starzyk P., and Schmitt R. 1976. Chemical evidence for the genesis of the ureilites, the achondrite

- Chassigny and the nakhlites. *Geochimica et Cosmochimica Acta* 40:1439–1447.
- Buddhue J. D. 1939. The oxidation of meteorites. *Contributions of the Society for Research on Meteorites* 2:75–79.
- Crozaz G. and Wadhwa M. 2001. The terrestrial alteration of saharan shergottites Dar al Ganid 476 and 489: A case study of weathering in a hot desert environment. *Geochimica et Cosmochimica Acta* 65:971–978.
- Crozaz G., Floss C., and Wadhwa M. 2003. Chemical alteration and REE mobilization in meteorites from hot and cold deserts. *Geochimica et Cosmochimica Acta* 67:4727–4741.
- Dauphas N. and Pourmand A. 2015. Thulium anomalies and rare earth element patterns in meteorites and Earth: Nebular fractionation and the nugget effect. *Geochimica et Cosmochimica Acta* 163:234–261.
- Djamali M., Akhani H., Khoshravesh R., Ponel P., and Brewer S. 2011. Application of the global bioclimatic classification to Iran: Implications for understanding the modern vegetation and biogeography. *Ecologia Mediterranea* 37:91–114.
- Dreibus G., Huisl W., Haubold R., and Jagoutz E. 2001. Influence of terrestrial desert weathering in Martian meteorites (abstract 50). *Meteoritics & Planetary Science* 36.
- Ebihara M. and Honda M. 1984. Distribution of rare earth elements and uranium in various components of ordinary chondrites. *Meteoritics* 19:69–77.
- Farpoor M. H. and Krouse H. R. 2008. Stable isotope geochemistry of sulfur bearing minerals and clay mineralogy of some soils and sediments in Loot Desert, central Iran. *Geoderma* 146:283–290.
- Floss C. and Crozaz G. 1991. Ce anomalies in the LEW85300 eucrite: Evidence for REE mobilization during Antarctic weathering. *Earth and Planetary Science Letters* 107:13–24.
- Folco L., D’Orazio M., and Perchiazzi N. 2007. Authenticating the recovery location of meteorites: The case of Castenaso. *Meteoritics & Planetary Science* 42:321–330.
- Gattacceca J., Valenzuela M., Uehara M., Jull A. J. T., Giscard M., Rochette P., Braucher R., Suavet C., Gounelle M., Morata D., Munayco P., Bourrot-Denis M., Bourles D., and Demory F. 2011. The densest meteorite collection area in hot deserts: The San Juan meteorite field (Atacama Desert, Chile). *Meteoritics & Planetary Science* 46:1276–1287.
- Goldsmith Y., Stein M., and Enzel Y. 2014. From dust to varnish: Geochemical constraints on rock varnish formation in the Negev Desert, Israel. *Geochimica et Cosmochimica Acta* 126:97–111.
- Gooding J. L. 1982. Mineralogical aspects of terrestrial weathering effects in chondrites from Allan Hills, Antarctica. *Lunar and Planetary Science* 12B:1105–1122.
- Govindaraju K. 1994. 1994 compilation of working values and sample description for 383 geostandards. *Geostandards and Geoanalytical Research* 18:1–158.
- Govindaraju K. 1995. 1995 working values with confidence limits for twenty-six CRPG, ANRT and IWG-GIT geostandards. *Geostandards and Geoanalytical Research* 19:1–32.
- Hartley A. J., Chong G., Houston J., and Mather A. E. 2005. 150 million years of climatic stability: Evidence from the Atacama Desert, northern Chile. *Journal of the Geological Society* 162:421–424.
- Hezel D. C., Schlüter J., Kallweit H., Jull A. J. T., Al Fakeer O. Y., Al Shamsi M., and Strekopytov S. 2011. Meteorites from the United Arab Emirates: Description, weathering, and terrestrial ages. *Meteoritics & Planetary Science* 46:327–336.
- Hutzler A. 2015. The flux of meteorites on Earth: Contribution of measuring the concentration of multiple cosmogenic nuclides, and collections in arid areas. Aix-Marseille Université, Marseille, France.
- Hutzler A., Gattacceca J., Rochette P., Braucher R., Carro B., Christensen E.J., Cournede C., Gounelle M., Laridhi Ouazaa N., Martinez R., Valenzuela M., Warner M., and Bourles D. 2016. Description of a very dense meteorite collection area in western Atacama: Insight into the long-term composition of the meteorite flux to Earth. *Meteoritics & Planetary Science* 51:468–482.
- Hyde B. C., Day J. M. D., Tait K. T., Ash R. D., Holdsworth D. W., and Moser D. E. 2014. Characterization of weathering and heterogeneous mineral phase distribution in brachinite Northwest Africa 4872. *Meteoritics & Planetary Science* 49:1141–1156.
- Inoue M., Nakamura N., and Kimura M. 2009. Tetrad effects in REE abundance patterns of chondrules from CM meteorites: Implications for aqueous alteration on the CM parent asteroid. *Geochimica et Cosmochimica Acta* 73:5224–5239.
- Jarosewich E., Clarke C. R. S., and Barrows J. N. 1987. Allende meteorite reference sample. *Smithsonian Contributions to the Earth Sciences* 27:1–49.
- Kabata-Pendias A. and Pendias H. 2001. Trace elements in soils and plants. Boca Raton, Florida: CRC Press. 23 p.
- Kagi H. and Takahashi K. 1998. Relationship between positive cerium anomaly and adsorbed water in Antarctic lunar meteorites. *Meteoritics & Planetary Science* 33:1033–1040.
- Kemp A. I. S. and Hawkesworth C. J. 2004. Granites and differentiation of the continental crust. *Geochimica et Cosmochimica Acta* 68:A667.
- Koeberl C. and Cassidy W. A. 1989. Differences between Antarctic and non-Antarctic meteorites. *An assessment. Geochimica et Cosmochimica Acta*. 55:3–18.
- Laveuf C. and Cornu S. 2009. A review on the potentiality of rare earth elements to trace pedogenetic processes. *Geoderma* 154:1–12.
- Lipin B. R. and McKay G. A. 1989. Geochemistry and mineralogy of rare earth elements. In *rare earth element chemistry*, edited by Lipin B. R. and McKay G. A. Chantilly, Virginia: Mineralogical Society of America. pp. 345–373.
- Llorca J., Roszjar J., Cartwright J. A., Bischoff A., Ott U., Pack A., Merchel S., Rugel G., Fimiani L., Ludwig P., Casado J. V., and Allepuz D. 2013. The Ksar Ghilane 002 shergottite—The 100th registered Martian meteorite fragment. *Meteoritics & Planetary Science* 48:493–513.
- Masuda A. and Tanaka T. 1978. REE, Ba, Sr and Rb in the Yamato meteorites, with special reference to Yamato-691 (a), -692(b) and -693(c). *Memoirs of National Institute of Polar Research*. Special issue8:229–232.
- Masuda A., Tanaka T., Asakura J., and Shimizu H. 1977. REE, Rb, Sr and Ba abundances in Yamato (j),(k) and (m) meteorites *Antarctic Record* 58:197–203.
- Mildrexler D. J., Zhao M., and Running S. W. 2011. Satellite finds highest land skin temperatures on Earth. *Bulletin of the American Meteorological Society* 92:855–860.
- Mittlefehldt D. W. and Lindstrom M. M. 1991. Generation of abnormal trace element abundances in Antarctic eucrites

- by weathering processes. *Geochimica et Cosmochimica Acta* 55:77–87.
- Murrell M. and Burnett D. 1983. The behavior of actinides, phosphorus, and rare earth elements during chondrite metamorphism. *Geochimica et Cosmochimica Acta* 47:1999–2014.
- Muscarella O. W. 2001. Jiroft and “Jiroft-Aratta”: A review article of Yousef Madjidzadeh, “Jiroft: The Earliest Oriental Civilization.” *Bulletin of the Asia Institute* 15:173–198.
- Olsen E. and Fuchs L. H. 1967. The state of oxidation of some iron meteorites. *Icarus* 6:242–253.
- Pillinger C. T., Greenwood R. C., Gibson J. M., Pillinger J. M., and Gibson E. K. 2013. The Holbrook Meteorite—99 years out in the weather. 44th Lunar and Planetary Science Conference. LPI Contribution No. 1719. Houston, Texas: Lunar and Planetary Institute. 2883 p.
- Pourkhorsandi H. and Mirnejad H. 2013. Lut Desert (Iran): A high-potential area for finding meteorites. 44th Lunar and Planetary Science Conference. CD-ROM.
- Pourkhorsandi H., Mirnejad H., Rochette P., and Hassanzadeh J. 2016. Lut 009, an H4 (S2, W4) ordinary chondrite meteorite from Lut Desert of Iran. *Journal of the Earth and Space Physics* 41:125–130.
- Saunier G., Poitrasson F., Moine B., Gregoire M., and Seddiki A. 2010. Effect of hot desert weathering on the bulk-rock iron isotope composition of L6 and H5 ordinary chondrites. *Meteoritics & Planetary Science* 45:195–209.
- Shimizu H., Masuda A., and Tanaka T. 1983. Cerium anomaly in REE pattern of Antarctic eucrite. *Memoirs of National Institute of Polar Research. Special issue* 30:341–348.
- Shinonaga T., Endo K., Ebihara M., Heumann K. G., and Nakahara H. 1994. Weathering of Antarctic meteorites investigated from contents of Fe<sup>3+</sup>, chlorine, and iodine. *Geochimica et Cosmochimica Acta* 58:3735–3740.
- Socki R. A., Gibson E. K., Jull A. J. T., and Karlsson H. R. 1991. Carbon and oxygen isotope composition of carbonates from an L6 chondrite: Evidence for terrestrial weathering from the Holbrook meteorite. *Meteoritics* 26:396.
- Swindle T. D., Kring D. A., Burkland M. K., Hill D. H., and Boynton W. V. 1998. Noble gases, bulk chemistry, and petrography of olivine-rich achondrites Eagles Nest and Lewis Cliff 88763: Comparison to brachinites. *Meteoritics & Planetary Science* 33:31–48.
- Thiagarajan N. and Aeolus Lee C.-T. 2004. Trace-element evidence for the origin of desert varnish by direct aqueous atmospheric deposition. *Earth and Planetary Science Letters* 224:131–141.
- Torigoye-Kita N., Misawa K., and Tatsumoto M. 1995. U-Th-Pb and Sm-Nd isotopic systematics of the Goalpara ureilite: Resolution of terrestrial contamination. *Geochimica et Cosmochimica Acta* 59:381–390.
- Valenzuela E. M. 2011. Procesos de Meteorización en Condritos Ordinarios del Desierto de Atacama, Norte de Chile: Nuevos Antecedentes sobre Meteorización de Material Extraterrestre en Ambientes Desérticos. Universidad de Chile.
- Velbel M. A. 2014. Terrestrial weathering of ordinary chondrites in nature and continuing during laboratory storage and processing: Review and implications for Hayabusa sample integrity. *Meteoritics & Planetary Science* 49:154–171.
- Wasson J. T. and Kallemeyn G. W. 1988. Compositions of chondrites. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 325:535–544.
- Welten K. C., Nishiizumi K., Finkel R. C., Hillegonds D. J., Jull A. J. T., Franke L., and Schultz L. 2004. Exposure history and terrestrial ages of ordinary chondrites from the Dar al Gani region, Libya. *Meteoritics & Planetary Science* 39:481–498.
- White J. S., Henderson E. P., and Mason B. 1967. Secondary minerals produced by weathering of Wolf Creek meteorite. *The American Mineralogist* 52:1190–1197.
- Wlotzka F. 1993. A weathering scale for the ordinary chondrites. *Meteoritics* 28:460.
- Zurfluh F. J. 2012. Quantification of terrestrial weathering and contamination in meteorites recovered in the Sultanate of Oman. Universität Bern.
- Zurfluh F. J., Hofmann B. A., Gnos E., Eggenberger U., Greber N. D., and Villa I. M. 2012. Weathering and strontium contamination of meteorites recovered in the sultanate of Oman. *Meteorite* 2000:34–38.